

Yersinia pestis YbtU and YbtT Are Involved in Synthesis of the Siderophore Yersiniabactin but Have Different Effects on Regulation

VALERIE A. GEOFFROY,[†] JACQUELINE D. FETHERSTON, AND ROBERT D. PERRY*

Department of Microbiology and Immunology, University of Kentucky, Lexington, Kentucky

Received 18 February 2000/Returned for modification 17 April 2000/Accepted 1 May 2000

One prerequisite for the virulence of *Yersinia pestis*, causative agent of bubonic plague, is the yersiniabactin (Ybt) siderophore-dependent iron transport system that is encoded within a high-pathogenicity island (HPI) within the *pgm* locus of the *Y. pestis* chromosome. Several gene products within the HPI have demonstrated functions in the synthesis or transport of Ybt. Here we examine the roles of *ybtU* and *ybtT*. In-frame mutations in *ybtT* or *ybtU* yielded strains defective in siderophore production. Mutant strains were unable to grow on iron-deficient media at 37°C but could be cross-fed by culture supernatants from a Ybt-producing strain of *Y. pestis*. The *ybtU* mutant failed to express four indicator Ybt proteins (HMWP1, HMWP2, YbtE, and Psn), a pattern similar to those for other *ybt* biosynthetic mutants. In contrast, strains carrying mutations in *ybtT* or *ybtS* (a previously identified gene required for Ybt biosynthesis) produced all four proteins at wild-type levels under iron-deprived conditions. To assess the effects of *ybtT*, *-U*, and *-S* mutations on transcription of *ybt* genes, reporter plasmids with *ybtP* or *psn* promoters controlling *lacZ* expression were introduced into these mutants. Normal iron-regulated β -galactosidase activity was observed in the *ybtT* and *ybtS* mutants, whereas a significant loss of expression occurred in the Δ *ybtU* strain. These results show that *ybtT* and *ybtU* genes are involved in the biosynthesis of the Ybt siderophore and that a *ybtU* mutation but not *ybtT* or *ybtS* mutations affects transcription from the *ybtP* and *psn* promoters.

To cause infections, pathogenic bacteria must be able to remove iron, an essential trace nutrient, from host iron- and/or heme-chelating proteins (12, 43, 65). *Yersinia pestis*, the causative agent of bubonic and pneumonic plague, possesses an ATP-binding cassette (ABC) hemoprotein transport system (Hmu) that allows it to use a variety of host hemoproteins (36, 64). The organism also contains one putative and two known inorganic iron transport systems. The Yfu system was identified by a search of the *Y. pestis* KIM10+ genome database (www.genome.wisc.edu) and belongs to a family of ABC iron transporters present in *Yersinia enterocolitica* (Yfu), *Neisseria* spp. (Fbp), *Haemophilus influenzae* (Hit), *Actinobacillus pleuropneumoniae* (Afu), and *Serratia marcescens* (Sfu) (7). Whether this system is functional in *Y. pestis* is currently unknown. The *Y. pestis* Yfe system belongs to a family of cation-transporting ABC systems and transports both iron and manganese. This system appears to function to acquire iron during the later stages of plague (3, 4).

The third inorganic iron transport system synthesizes the siderophore yersiniabactin (Ybt), which is composed of a phenolate, a thiazoline, and a thiazolidine ring (15, 19, 47) and which has considerable similarity to the siderophores pyochelin and anguibactin produced by *Pseudomonas aeruginosa* and *Vibrio anguillarum*, respectively (18, 38). The Ybt iron acquisition system is essential for the virulence of *Y. pestis* during the early stages of infection in mice (3). Ybt biosynthetic, regulatory, and transport genes are encoded within a high-pathogenicity island (HPI) that is present in highly pathogenic isolates of *Y. pestis*, *Yersinia pseudotuberculosis*, and *Y. enterocolitica*, as

well as several types of pathogenic *Escherichia coli* (8, 9, 13, 29, 33, 50, 56). In *Y. pestis*, HPI resides within the *pgm* locus, a 102-kb region of chromosomal DNA subject to high-frequency deletion (9, 10, 26, 29, 32, 41).

The genes encoding the Ybt systems of *Y. pestis* (see Fig. 1) and *Y. enterocolitica* have been completely sequenced and show >97% sequence identity (10, 29, 46, 50). One notable exception to this sequence identity is the unique insertion of a 125-bp ERIC sequence (enterobacterial repetitive intergenic consensus sequence; also called an intergenic repeated unit) within the promoter region of *ybtA* of *Y. enterocolitica* (50). *ybtA* encodes a transcriptional activator of *ybt* genes (22). Although this suggests possible differences in regulatory responses, some of the Ybt biosynthetic genes appear to be functionally interchangeable among the three pathogenic yersiniae: *Y. pestis*, *Y. pseudotuberculosis*, and *Y. enterocolitica* (14, 47).

Iron from Fe-Ybt is transported into the cell via an outer membrane (OM) receptor (termed Psn in *Y. pestis* and FyuA in *Y. enterocolitica*) in conjunction with an ABC transport system encoded by *ybtP* and *ybtQ*. Psn, which likely binds the Fe-Ybt complex, also binds the bacteriocin pesticin (23–25, 34, 40, 51). Translocation of the substrate across the OM is TonB dependent (21, 34). YbtP and YbtQ resemble inner-membrane permeases that are each fused to an ATP-binding domain. Both proteins are required for use of iron from Ybt. No periplasmic binding protein has been identified for the Ybt system (23, 29).

Ybt production occurs via a mixed polyketide synthase-nonribosomal peptide synthetase (NRPS) strategy that assembles the siderophore in modular fashion from salicylate, a linker group derived from malonyl-coenzyme A, three molecules of cysteine, and three methyl groups donated by *S*-adenosylmethionine (29). The requirement of three gene products (high-molecular-weight protein 2 [HMWP2], YbtE, and YbtS) for Ybt synthesis has been clearly demonstrated genetically. YbtS

* Corresponding author. Mailing address: Department of Microbiology and Immunology, MS415 Medical Center, University of Kentucky, Lexington, KY 40536-0084. Phone: (859) 323-6341. Fax: (859) 257-8994. E-mail: rperry@pop.uku.edu.

[†] Formerly Valérie A. Coulanges.

is likely required for the final steps in salicylate biosynthesis (2, 29). YbtE adenylates salicylate and transfers this activated compound to HMWP2 (30). HMWP2, encoded by *irp2*, possesses domains involved in nonribosomal peptide synthesis and likely participates in the initial cyclization and condensation reactions involving salicylate and two cysteine molecules (29–31, 62). HMWP1, encoded by *irp1*, contains polyketide/fatty acid synthase and modified NRPS domains that add the branched isobuteryl-alcohol linker and the last thiazoline moiety. Phosphopantetheinylation of a peptidyl carrier protein domain of HMWP1 (PCP3) has been demonstrated (29). The roles of the remaining *ybt* genes (*ybtX*, *ybtT*, and *ybtU*) contained within the HPI were undetermined or uncertain. The product encoded by *ybtX* is predicted to be extremely hydrophobic. However, a strain carrying a deletion in *ybtX* had no discernible *in vitro* phenotype (23). While YbtT contains a thioesterase domain (2, 29), a definitive role for YbtT in Ybt biosynthesis has not been demonstrated. YbtU has no significant homology to proteins in the database that have defined enzymatic or regulatory functions (29), nor has a function for YbtU in Ybt production or utilization been determined. In this study, we report that in-frame deletions in *ybtU* and *ybtT* abolish Ybt siderophore synthesis. Similar to mutations in other Ybt biosynthetic genes, the *ybtU* mutation downregulates expression of *ybt* genes; however, mutations in *ybtT* and *ybtS* did not have this regulatory effect.

MATERIALS AND METHODS

Bacterial strains, media, and culture conditions. All relevant characteristics of strains used in this study are presented in Table 1. All the *Y. pestis* strains used in this study were derived from KIM6+, an avirulent strain that possesses all of the known *Y. pestis* virulence determinants except for pCD1, a 70.5-kb plasmid encoding the low-calcium response (Lcr) stimulon (26, 59). The Lcr virulence regulon is unrelated to the Pgm⁺ phenotype and has no demonstrable role in iron metabolism (48, 49).

All strains were stored at -20°C in phosphate-buffered glycerol. *Y. pestis* cells were grown routinely at 30°C on Congo red agar (63) from glycerol stocks and then grown in heart infusion broth (Difco Laboratories) or on tryptone-blood agar base (Difco). For iron-deficient growth, *Y. pestis* cells were grown in the chemically defined medium PMH, which had been extracted prior to use with Chelex 100 resin (Bio-Rad Laboratories) (61). The residual iron that was not removed from deferrated PMH by the resin was precipitated by the addition of 0.5 mM NaCO₃-0.01 mM MnCl₂-4.0 mM CaCl₂ (PMH-S) or chelated by supplementation with 2,2'-dipyridyl (PMH-DIP) at a concentration of 100 μM. PMH-S and PMH-DIP plates were solidified with 1% agarose. PMH-S and PMH-DIP plates were subsequently used in cross-feeding experiments or to determine the growth characteristics of the *ybt* mutants at 37°C as previously described (24). For iron-replete growth, *Y. pestis* strains were cultivated in PMH supplemented with 10 μM FeCl₃.

All glassware used for iron-restricted studies was soaked overnight in chromic/sulfuric acid (46.3 g of K₂Cr₂O₇ per liter of 12 M sulfuric acid) or ScotClean (OWL Scientific, Inc.) to remove contaminating iron and copiously rinsed in deionized water. *E. coli* cells were grown in Luria broth. Where appropriate, ampicillin (100 μg/ml), spectinomycin (100 μg/ml), tetracycline (6.25 μg/ml), streptomycin (50 μg/ml), and kanamycin (50 μg/ml) were added to cultures.

Plasmids, sequencing, and recombinant DNA techniques. All the plasmids used in this study are listed in Table 1. Plasmids were purified from overnight cultures by alkaline lysis (6) and further purified when necessary by polyethylene glycol precipitation (37). Standard cloning and recombinant DNA methods (53) were used to construct the various plasmids in Table 1. A standard CaCl₂ procedure was used to introduce plasmids into *E. coli* (53). *Y. pestis* cells were transformed by electroporation as previously described (24). Plasmid DNA was sequenced by the dideoxynucleotide chain termination method (54) using Sequenase, version 2.0 (Amersham Pharmacia Biotech), ³⁵S-dATP (New England Nuclear/Dupont), and 7-deaza-dGTP. Synthetic oligonucleotide primers purchased from Integrated DNA Technologies were used to extend the sequence. Samples were electrophoresed at 70 W on 6% polyacrylamide gels containing Tris-borate-EDTA buffer and 8.3 M urea. Gels were fixed in a 10% ethanol-10% acetic acid solution, dried, and exposed to Kodak BioMax MR film at room temperature.

Generating *ybtU* and *ybtT* mutant strains and expression plasmids. All *Y. pestis* mutant strains were generated by homologous recombination using mutated DNA fragments cloned into suicide vectors carrying the *sacB* gene and an R6K origin of replication. For construction of an in-frame *ΔybtT* strain (KIM6-2072), we first subcloned an ~3.9-kb *SphI* fragment of plasmid pPSN3 into the

SphI site of pACYC184 to yield pYbtTU1. Removal of a 438-bp *PvuI* fragment from pYbtTU1 resulted in plasmid pYbtT1. Subcloning an ~3.5-kb *SphI* fragment from pYbtT1 into the *SphI* site of the suicide plasmid pSUC1 generated pYbtT1.1.

To generate the in-frame *ΔybtU* mutant (strain KIM6-2071), a 526-bp *EagI/EcoRV* fragment of plasmid pYbtTU1 was subcloned into the corresponding sites in pBluescript II KS+ (Stratagene) to yield pYbtU1. A 467-bp *PvuII* fragment was removed from pYbtTU1 and ligated into the *EcoRV* site of the plasmid pYbtU1, generating pYbtU2. The orientation of the insert was determined by DNA sequencing. We then subcloned an ~1-kb *SacI/SalI* fragment of pYbtU2 into the *SacI/SalI* sites in pSUC1 to yield pYbtU2.1. Gene replacement attempts with pYbtU2.1 did not yield the desired chromosomal integrants. Therefore, we constructed an alternate suicide plasmid, pYbtU2.2, by subcloning a *SalI/SmaI* fragment of pYbtU2.1 into the *SalI/SmaI* sites in pKNG101.

To generate hexahistidine fusion proteins, the *ybtT* and *ybtU* gene coding regions were amplified with *Pfu* polymerase (Stratagene) from pPSN345 by PCR using primers ybtT-1 (5'-TGATGGCGCCTCTGTGACGCAATCTGCAATG-3') and M13 reverse (5'-AGCGGATAACAATTTCAC-3') and ybtU-1 (GGAATCTCTATGATGCCGTCCGCTCC) and ybtU-2 (CGGGATCTCACAGCCCTCTTATC), respectively. Reaction mixtures contained 0.2 mM deoxy nucleoside triphosphates and 0.2 μM primers, and reactions consisted of 20 s at 94°C, 20 s at 50°C, and 90 s at 72°C for 30 cycles, followed by a single cycle at 72°C for 10 min for *ybtT*. Amplification conditions for *ybtU* were 45 s at 94°C, 30 s at 50°C, and 2 min at 72°C for 30 cycles followed by a single cycle at 72°C for 10 min. The *ybtS* coding region was amplified with *Pfu* polymerase (Stratagene) from pSDR498.1 using primers ybtS-1 (5'-GGAATCTCTATGAAAATCAGTGAATT-3') and ybtS-2 (5'-CGGGATCCCTACCAATTAATAGG-3'). Amplification conditions for *ybtS* consisted of 45 s at 94°C, 30 s at 45°C, and 2 min at 72°C for 30 cycles followed by a single cycle at 72°C for 10 min. The 827-, 1,305-, and 1,100-bp products of *ybtT*, *ybtS*, and *ybtU*, respectively, were purified from low-melting-point agarose and digested with *KasI/EcoRI* (for *ybtT*) or with *BamHI/EcoRI* (for *ybtS* and *ybtU*), followed by ligation into the corresponding sites of pPROEX1, yielding pYbtT-H6, pYbtS-H6, and pYbtU-H6. Ligated products were transformed into DH5α cells. Positive clones containing the desired inserts were identified by restriction enzyme digests and verified by PCR using a set of nested primers specific for the *ybtS*, *ybtT*, and *ybtU* coding regions. Expression of all three gene products was verified in the positive clones by IPTG (isopropyl-β-D-thiogalactopyranoside) induction of Luria broth mini-cultures followed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) analysis (data not shown). pYbtT-H6, pYbtS-H6, and pYbtU-H6 were electroporated into *ΔybtT*, *ΔybtS*, and *ΔybtU* mutants, respectively. Growth stimulation in these three strains was tested by bioassay on PMH-DIP as described above.

Protein analyses. To label cellular proteins, whole cells of *Y. pestis* strains acclimated to growth under iron-deficient or iron-sufficient conditions by serial passage in PMH, with or without FeCl₃ (10 μM), for a total of approximately six generations, were labeled with ³⁵S-amino acids (DuPont NEN Research Products) for 1 h as previously described (25). To analyze the effect of Ybt on protein synthesis, purified Ybt from *Y. pestis* KIM6+ was added to cells, acclimated to iron starvation, at the same time as ³⁵S-amino acids. An equivalent number of counts was electrophoresed on 7.5% polyacrylamide gels containing SDS. Dried gels were exposed to Kodak BioMax MR film at room temperature.

Ybt bioassay. Culture supernatants were obtained from *ybt* mutants inoculated into deferrated PMH and grown for a total of six to nine generations at 37°C as previously described (24). The cells were pelleted by centrifugation, and the supernatant was filtered through a 0.45-μm-pore-size filter. For growth responses, PMH-S or PMH-DIP plates were overlaid with 0.04 optical density (at 620 nm) units of KIM6-2046.1 (*irp2::kan2046.1*) cells grown in deferrated PMH and 25 μl of filtered supernatants from iron-deficient *ybt* mutant cultures were added to wells in the plates.

The *ybtU* and *ybtT* mutants were also tested for their ability to promote the growth of KIM6-2046.1 at 37°C by streaking the mutants adjacent to KIM6-2046.1 on PMH-S or PMH-DIP plates. Prior to streaking, the mutants were adapted to iron-deficient growth conditions as described above. *Y. pestis* strains that do not produce Ybt are unable to grow on PMH-S or PMH-DIP at 37°C but can be cross-fed by Ybt-producing strains.

β-Galactosidase assays. Lysates were prepared from cells carrying the *ybtP::lacZ* or *psn::lacZ* reporter plasmid and grown in PMH in the presence or absence of iron through two transfers for a total of approximately six generations, as previously described (60). β-Galactosidase activities were measured spectrophotometrically following cleavage of ONPG (4-nitro-phenyl-β-D-galactopyranoside). Activity is expressed in Miller units (44).

Ybt detection and purification. *Y. pestis* KIM6+, KIM6-2071, and KIM6-2072 were grown at 37°C in 100 ml of deferrated PMH for approximately eight generations. The supernatant of each culture was filtered through a 0.45-μm-pore-size filter. The presence of the Ybt siderophore was determined using methods modified from Chambers et al. (15) and Drechsel et al. (19) as previously described (47). Three C18-SEP-PAK cartridges were used in a preliminary purification step. The sample, in 50% methanol, was then applied to an analytical C-18 high-pressure liquid chromatography (HPLC) column and eluted with 100% methanol. The Ybt siderophore was detected by its absorbance maximum at 385 nm (19) and by bioassay.

TABLE 1. Bacterial strains and plasmids^a

Strain or plasmid	Relevant characteristic(s)	Reference or source
<i>E. coli</i>		
DH5 α	Cloning host	1
DH5 α (λ pir)	Strain for propagating plasmids with R6K origins, derived from DH5 α	S. C. Straley
<i>Y. pestis</i>		
KIM6+	Pgm ⁺ (Hms ⁺ Ybt ⁺) Lcr ⁻	24
KIM6-2046.1	Hms ⁺ Ybt ⁻ (<i>irp2::kan2046.1</i>) Lcr ⁻ Km ^r	47
KIM6-2046.3	Hms ⁺ Ybt ⁻ (Δ <i>irp2-2046.3</i>) Lcr ⁻	2
KIM6-2070.1	Hms ⁺ Ybt ⁻ (<i>ybtS::kan2070.1</i>) Lcr ⁻ Km ^r	29
KIM6-2071	Hms ⁺ Ybt ⁻ (Δ <i>ybtU2071</i>) Lcr ⁻	This study
KIM6-2072	Hms ⁺ Ybt ⁻ (Δ <i>ybtT2072</i>) Lcr ⁻	This study
Plasmids		
pACYC184	4.2-kb cloning vector; Cm ^r Tc ^r	16
pBluescript II KS+	2.9-kb cloning vector; Ap ^r	Stratagene
pEUPP1	15.4-kb low-copy-number <i>psn::lacZ</i> reporter plasmid; Spc ^r ; iron-, Fur-, and YbtA-regulated expression of β -galactosidase	22
pEUYbtP	15.4-kb low-copy-number <i>ybtP::lacZ</i> reporter plasmid; Spc ^r ; iron-, Fur-, and YbtA-regulated expression of β -galactosidase	23
pKNG101	6.8-kb suicide vector; <i>sacB</i> ⁺ ; R6K origin; Sm ^r	39
pPROEX-1	4.7-kb protein expression vector; Ap ^r	Gibco/BRL
pPSN3	9.9-kb <i>SalI</i> fragment from pSDR498.4 ligated into pBGL2; Ap ^r	25
pPSN345	3.45-kb <i>SphI/EcoRI</i> fragment from pPSN3 ligated into pUC18; <i>ybtU</i> ⁺ <i>ybtT</i> ⁺ ; Ap ^r	This study
pSDR498.1	~4.6-kb <i>BamHI/EcoRI</i> fragment from pSDR498 ligated into pHC79; <i>ybtS</i> ⁺ ; Ap ^r	29, 35
pSUC1	4.7-kb suicide vector; <i>sacB</i> ⁺ ; R6K origin; Ap ^r	23
pUC18	2.7-kb cloning vector; Ap ^r	68
pYbtS-H6	1,305-bp PCR fragment from pSDR498.1 ligated into pPROEX-1; IPTG-regulated expression of YbtS-H6; Ap ^r	This study
pYbtT-H6	800-bp PCR fragment from pPSN345 ligated into pPROEX-1; IPTG-regulated expression of YbtT-H6; Ap ^r	This study
pYbtU-H6	1,100-bp PCR fragment from pPSN345 ligated into pPROEX-1; IPTG-regulated expression of YbtU-H6; Ap ^r	This study
pYbtTU1	~3.9-kb <i>SphI</i> fragment containing <i>ybtT</i> and <i>ybtU</i> from pPSN3 ligated into pACYC184; Cm ^r	This study
pYbtT1	pYbtTU1 with 438-bp <i>PvuI</i> fragment deleted; Cm ^r	This study
pYbtT1.1	3.5-kb <i>SphI</i> fragment from pYbtT1 ligated into pSUC1; Δ <i>ybtT2072</i> (deletion of 808 bp), <i>sacB</i> ⁺ ; R6K origin; Ap ^r	This study
pYbtU1	526-bp <i>EagI/EcoRV</i> fragment from pYbtTU1 ligated into pBluescript II KS+; Ap ^r	This study
pYbtU2	467-bp <i>PvuII</i> fragment from pYbtTU1 ligated into the <i>EcoRV</i> site of pYbtU1; Cm ^r	This study
pYbtU2.1	~1.0-kb <i>SacI/SalI</i> fragment from pYbtU2 ligated into pSUC1; Δ <i>ybtU2071</i> (deletion of 225 bp), <i>sacB</i> ⁺ ; R6K origin; Ap ^r	This study
pYbtU2.2	~1.0-kb <i>SalI/SmaI</i> fragment from pYbtU2.1 ligated into pKNG101; Δ <i>ybtU</i> (deletion of 225 bp); <i>sacB</i> ⁺ ; R6K origin; Sm ^r	This study

^a *Y. pestis* strains with a plus sign possess an intact 102-kb *pgm* locus containing the genes for hemin storage (*hms*) and the Ybt system. All other *Y. pestis* strains contain a mutation within the *pgm* locus due to either a deletion or insertion of an antibiotic resistance cassette. Strains synthesizing the siderophore yersiniabactin are designated Ybt⁺, while those affected in yersiniabactin production are Ybt⁻. Lcr⁻ indicates the absence of the low-calcium response virulence plasmid pCD1. Ap^r, Km^r, Spc^r, Sm^r, Tc^r, and Cm^r, resistance to ampicillin, kanamycin, spectinomycin, streptomycin, tetracycline, and chloramphenicol, respectively.

RESULTS

Mutation of *ybtU* or *ybtT* causes loss of siderophore production. Most biosynthetic enzymes for the synthesis of Ybt are encoded within a large, putative operon of five genes: *irp2*, *irp1*, *ybtU*, *ybtT*, and *ybtE* (Fig. 1). The likely functions of HMWP1, HMWP2, and YbtE in the synthesis of Ybt have been described previously (29, 30, 62). To study the functions of YbtT and YbtU in Ybt production or utilization, we constructed in-frame deletions in both genes (Fig. 1). We have been unable to identify the YbtU and YbtT products by SDS-PAGE. However, when grown under appropriate conditions, both the YbtT⁻ and YbtU⁻ mutants express YbtE (Fig. 2), the product of the final, downstream gene of the operon, indicating that both mutations are in-frame deletions.

Two different media (PMH-S and PMH-DIP) were used to examine the ability of the mutants to grow at 37°C under iron-chelated conditions. Neither KIM6-2071 (Δ *ybtU*) nor

KIM6-2072 (Δ *ybtT*) grew at 37°C on the iron-chelated media (Table 2), indicating that the mutated strains lost the ability to either synthesize or utilize Ybt. Supernatants from iron-deficient cultures of KIM6-2071 and KIM6-2072 were unable to stimulate the growth of a *Y. pestis* strain (KIM6-2046.1) defective in Ybt synthesis (Table 2). However, culture supernatant from KIM6+, a yersiniabactin-producing strain of *Y. pestis*, allowed the growth of KIM6-2046.1 as well as the YbtU⁻ and YbtT⁻ mutants under these conditions. Similar results were obtained with KIM6-2070.1 (Δ *ybtS::kan2070.1*) (Table 2), a strain previously demonstrated to be defective in Ybt synthesis (29). These results suggest that the YbtU⁻ and YbtT⁻ mutants are defective in synthesis of the Ybt siderophore. To clearly demonstrate that Ybt was not produced at significant levels, we attempted to purify Ybt from 100 ml of iron-deficient culture supernatant of each mutant, as well as the KIM6+ parental strain. Although a significant amount of Ybt was produced by




FIG. 1. Genetic organization of the *Y. pestis* Ybt operons. Arrows, putative promoters for the operons. Gene designations are given below the lines, while nucleotide numbers are shown above the lines. Numbering corresponds to that used for GenBank accession no. AF091251 (29). An expanded version of the *irp1-irp2-ybtU-ybtT-ybtE* operon is shown; *irp1*, *irp2*, and *ybtE* genes are not drawn to scale in the expanded version. The size and location of the *ybtU* and *ybtT* deletions are indicated on the expanded map.

KIM6+ as determined by HPLC analysis, we were not able to detect any Ybt production by the YbtU⁻, YbtT⁻, and YbtS⁻ mutants (data not shown).

For complementation analyses, *ybtT*, *ybtU*, and *ybtS* were amplified by PCR and cloned into an IPTG-inducible His tag expression vector generating pYbtT-H6, pYbtU-H6, and pYbtS-H6. The recombinant plasmids restored the ability of the respective mutants to grow on iron-chelated plates at 37°C. In addition, culture supernatant from each of the complemented strains was able to promote the growth of KIM6-2046.1 (*irp2::kan2046.1*) on PMH-DIP plates (Table 2). Although the genes encoding the recombinant proteins are inducible by IPTG in *E. coli*, there is no direct evidence for IPTG induction in *Y. pestis*. SDS-PAGE analysis showed that YbtU-H6 expression was induced by IPTG in *Y. pestis* but to a much lesser extent than in *E. coli* (data not shown). However, in the experiments described above, IPTG addition was not necessary for complementation in *Y. pestis*, indicating that there was sufficient production of the fusion proteins in the absence of IPTG to restore Ybt synthesis. Furthermore, the additional amino acid residues on the His-tagged fusion proteins apparently did not perturb their ability to function in Ybt synthesis.

The $\Delta ybtU2071$ mutation results in decreased expression of ybt operons. Total ³⁵S-labeled proteins synthesized by cells grown under iron-sufficient and iron-deficient conditions were analyzed by SDS-PAGE to determine the effect, if any, of the $\Delta ybtU2071$ mutation on the expression of other iron-regulated proteins. Previously, we showed that mutations causing loss of siderophore production lower expression of other Ybt biosynthetic genes (those encoding HMWP1, HMWP2, YbtE, and YbtE) as well as the Ybt receptor (Psn) (2, 22). The analysis of whole-cell extracts of strains grown in iron-deficient media revealed that the YbtU⁻ mutant had a pattern of protein expression similar to that of a $\Delta irp2-2046.3$ mutant that is defective in siderophore biosynthesis. In the absence of iron, the levels of expression of HMWP1, HMWP2, YbtE, and Psn proteins were greatly reduced in KIM6-2071 ($\Delta ybtU2071$) and KIM6-2046.3 ($\Delta irp2-2046.3$) cells (Fig. 2, lanes 4 and 7) compared to those in the parental strain, KIM6+ (Fig. 2, lane 1). Cells grown in the presence of iron do not express detectable levels of these proteins (Fig. 2, lanes 3, 6, 9, 12, and 15).

We have previously shown that addition of purified Ybt or

supernatant containing Ybt to *Y. pestis* Ybt biosynthetic mutants restores expression of HMWP1, HMWP2, YbtE, and Psn (2, 22, 47). Similarly, when purified Ybt was added to iron-deficient cultures, both the YbtU⁻ and Δ HMWP2 mutants expressed wild-type levels of HMWP1, HMWP2, Psn, and YbtE (Fig. 2, lanes 5 and 8).

To test the effect of the *ybtU2071* mutation on gene transcription, we used two well-characterized reporter plasmids with promoters that have been demonstrated to be regulated by Fur, iron, YbtA, and the Ybt siderophore: pEUYbtP, a *ybtP* promoter fusion to *lacZ*, and pEUPP1, a *psn* promoter fusion to *lacZ*. Both reporter genes were cloned into the low-copy-number plasmid pEU730 (2, 22, 23, 27). Because our *Y. pestis* strains are phenotypically β -galactosidase negative, any β -galactosidase activity is due to the presence of the reporter plasmid (61). The β -galactosidase activities of cells bearing these plasmids and grown in deferrated PMH, in the presence or absence of added iron, are presented in Table 3. As expected for KIM6+, which contains all the genes needed for Ybt synthesis and utilization, expression of *lacZ* from the *ybtP* promoter is iron regulated; there is a 23-fold repression of β -galactosidase activity in cells grown in the presence of surplus iron compared to those cultured under iron-deficient conditions. Expression of the *ybtP::lacZ* reporter in the YbtU⁻ mutant, KIM6-2071, was still somewhat iron regulated (12-fold repression); however the overall expression was greatly reduced (Table 3). The *ybtP* promoter was more active than the *psn* promoter during iron deficiency in the Ybt⁺ strain: a 23-fold versus a 14-fold induction. However, the effects of the *ybtU* mutation on the two promoters were similar, reducing expression to ~2,000 Miller units, a 16-fold reduction for the *ybtP* promoter and an 11-fold reduction for the *psn* promoter (Table 3). These studies suggest that loss of expression of HMWP1, HMWP2, YbtE, and Psn in the YbtU⁻ mutant results from decreased transcription from the relevant *ybt* promoters.

The level of expression from the *psn::lacZ* reporter in KIM6+ is higher here (Table 3) than in previously reported results (22). Although several factors, including the level of residual iron remaining in different preparations of PMH, could help account for this, we have found that mutations occurred within the promoter region of pEUPP1 after its construction. In other reporter strains, we have identified a single




FIG. 2. SDS-PAGE analysis of whole-cell proteins from *Y. pestis* strains grown in PMH in the presence (+) or absence (-) of FeCl_3 (10 μM) or Ybt. Ybt was added since previous studies showed induction of Ybt protein expression when the siderophore was added to Ybt biosynthetic mutants (2, 22). Cultures from *Y. pestis* KIM6+ (lanes 1 to 3), KIM6-2071 ($\Delta ybtU2071$; lanes 4 to 6), KIM6-2046.3 ($\Delta irp2-2046.3$; lanes 7 to 9), KIM6-2072 ($\Delta ybtT2072$; lanes 10 to 12), and KIM6-2070.1 (*ybtS::kan2070.1*; lanes 13 to 15) were incubated with ^{35}S -labeled amino acids for 1 h. Total cellular proteins were separated on a 7.5% polyacrylamide gel and visualized by autoradiography. Arrows, iron-regulated proteins HMWP1 (240 kDa), HMWP2 (190 kDa), Psn (68 kDa), and YbtE (56 kDa) and the truncated *irp2* gene product (ΔHMWP2) expressed by the *Y. pestis* in-frame deletion mutant KIM6-2046.3 ($\Delta irp2-2046.3$).

base change in the putative -10 region that reduces its similarity to the consensus and could decrease overall transcriptional efficiency (data not shown). A similar mutation might have caused the lower expression values reported previously (22). The pEUPP1 plasmid used in this study does not contain this mutation.

The *ybtT2072* and *ybtS2070.1* mutations do not affect expression of *ybt* operons. We also performed analyses of iron-repressible protein expression by *YbtT*⁻ and *YbtS*⁻ mutants of *Y. pestis*. Expression of HMWP1, HMWP2, YbtE, and Psn by

these two mutants was iron repressible with patterns similar to that of the parental strain, KIM6+ (Fig. 2). In contrast to the *YbtU*⁻ and ΔHMWP2 mutants, the *YbtT*⁻ and *YbtS*⁻ mutants produced all four proteins at wild-type levels under iron-deprived conditions (Fig. 2, lanes 10 and 13) and the addition of purified Ybt during growth of these mutants did not increase expression of these four proteins (Fig. 2, lanes 11 and 14).

We also tested the effects of the *ybtT* and *ybtS* mutations on transcription of the *ybtP* and *psn* promoter fusions to *lacZ*. The β -galactosidase activities from strains KIM6-2072 ($\Delta ybtT$) and

TABLE 2. Growth of *Y. pestis* KIM6+ and *ybt* mutants on PMH-S and PMH-DIP

Strain	Relevant characteristics	Growth on PMH-DIP and PMH-S ^a	Growth stimulation on PMH-DIP and PMH-S	
			of KIM6-2046.1 ^b	by KIM6+ ^c
KIM6+	<i>ybt</i> ⁺	+	+	ND ^d
KIM6-2071	$\Delta ybtU2071$	-	-	+
KIM6-2072	$\Delta ybtT2072$	-	-	+
KIM6-2070.1	<i>ybtS::kan2070.1</i>	-	-	+
KIM6-2071(pYbtU-H6)	$\Delta ybtU2071 ybtU-H6$	+	+	ND
KIM6-2072(pYbtT-H6)	$\Delta ybtT2072 ybtT-H6$	+	+	ND
KIM6-2070.1(pYbtS-H6)	$\Delta ybtS::kan2070.1 ybtS-H6$	+	+	ND
KIM6-2046.1	<i>irp2::kan2046.1</i>	-	-	+

^a The presence (+) or absence (-) of growth on PMH-S and PMH-DIP plates at 37°C.

^b Each strain was tested for its ability to promote the growth of KIM6-2046.1 (*irp2::kan2046.1*) at 37°C on PMH-S and PMH-DIP either by streaking adjacent to KIM6-2046.1 or by spent-culture supernatants.

^c Each strain was tested for its ability to use the exogenous Ybt siderophore at 37°C by streaking adjacent to KIM6+ on PMH-S and PMH-DIP at 37°C.

^d ND, not determined.

KIM6-2070.1 ($\Delta ybtS$) carrying either pEUYbtP or pEUPP1 were essentially the same as that observed with KIM6+ cells carrying these reporter plasmids (Table 3). These results indicate that the $\Delta ybtT2072$ and $\Delta ybtS2070.1$ mutations do not affect the transcription of the *ybtPQXS* or *psn* operons or expression of Ybt proteins. This is in sharp contrast to the previous three Ybt biosynthetic mutants, which all decreased expression of the Ybt system (this study; 2, 22, 23, 47).

DISCUSSION

While there are some differences among the HPIs of the pathogenic yersiniae, all encode an essentially identical and interchangeable siderophore-iron transport system (Ybt). The *ybt* genes appear to be organized into four operons: (i) *psn*, (ii) *irp2-irp1-ybtU-ybtT-ybtE*, (iii) *ybtA*, and (iv) *ybtP-ybtQ-ybtX-ybtS* (10, 23, 24, 29, 46, 47) (Fig. 1). In *Y. pestis*, it has previously been shown that the OM receptor Psn and an ABC transporter composed of YbtP and YbtQ are essential for use of the Ybt siderophore (23, 24). YbtA is an AraC-type activator for the other *ybt* operons and represses its own transcription (22). The role of YbtX is unclear since a mutation disrupting *ybtX* had no demonstrable effect on Ybt synthesis or iron uptake (23).

HMWP1 and HMWP2 (encoded by *irp1* and *irp2*, respectively), YbtE, and YbtS have demonstrated roles in siderophore synthesis (2, 29, 30). Here we have determined that the two remaining genes, *ybtU* and *ybtT*, also appear to be required for Ybt siderophore synthesis. Thus, cells with mutations in *ybtU* or *ybtT* were unable to grow on iron-deficient medium at 37°C and culture supernatants from these mutants could not stimulate the growth of a Ybt biosynthetic mutant and did not contain detectable levels of siderophore. However, the *ybtU* and *ybtT* mutants could grow on iron-deficient medium at 37°C when provided with exogenous Ybt. These results suggest that the *ybtU* and *ybtT* mutants can use Ybt but are unable to produce the siderophore.

While our results indicate that YbtT and YbtU are required for Ybt synthesis, the precise role of these proteins in the production of the siderophore is unknown. YbtT has a thioesterase domain with the G(Y/W/H)SxG signature motif (amino acids [aa] 102 to 106 in YbtT) and a distal GxH conserved sequence (aa 134 to 136 in YbtT) (2, 29, 58). The four highest homologies to YbtT in the database (3 January 2000) are to external putative thioesterases; NrpT in *Proteus mirabilis* is part of a NRPS/polyketide synthase system involved in swarm-

TABLE 3. β -Galactosidase activities of *Y. pestis* strains containing either a *ybtP::lacZ* or a *psn::lacZ* reporter plasmid^a

Strain	β -Galactosidase activity ^b of cells grown:			Ratio	
	-Fe	+Fe	-Fe/+Fe	Wild type/mutant	
				-Fe	+Fe
With pEUYbtP (<i>ybtP::lacZ</i>)					
KIM6+ (<i>ybt</i> ⁺)	30,356 \pm 4,196	1,308 \pm 162	23.2 \pm 2.6		
KIM6-2071 ($\Delta ybtU2071$)	1,926 \pm 394	165 \pm 18	11.6 \pm 2.3	15.8 \pm 3.8	7.9 \pm 0.7
KIM6-2072 ($\Delta ybtT2072$)	29,541 \pm 8,474	1,324 \pm 317	22.3 \pm 8.4	1.0 \pm 0.2	1.0 \pm 0.4
KIM6-2070.1 (<i>ybtS::kan2070.1</i>)	29,670 \pm 2,637	1,106 \pm 105	29.2 \pm 3.5	1.0 \pm 0.2	1.2 \pm 0.1
With pEUPP1 (<i>psn::lacZ</i>)					
KIM6+ (<i>ybt</i> ⁺)	22,199 \pm 3,388	1,595 \pm 400	13.9 \pm 5.5		
KIM6-2071 ($\Delta ybtU2071$)	2,042 \pm 326	138 \pm 37	14.8 \pm 2.0	10.9 \pm 1.6	11.5 \pm 6.0
KIM6-2072 ($\Delta ybtT2072$)	23,125 \pm 3,351	949 \pm 311	24.4 \pm 9.1	1.0 \pm 0.1	1.7 \pm 0.5
KIM6-2070.1 (<i>ybtS::kan2070.1</i>)	22,349 \pm 3,370	1,386 \pm 585	16.1 \pm 5.5	1.0 \pm 0.2	1.2 \pm 0.3
With no reporter					
KIM6+ (<i>ybt</i> ⁺)	0	0			

^a Cells were harvested during exponential growth at 37°C after approximately six generations in PMH containing either no added iron (-Fe) or 10 μ M FeCl₃ (+Fe).

^b Enzyme activities are expressed in Miller units (44). The values are averages \pm standard deviations of three or four individual reactions from two or more independent cultures.

YbtU 1 -----MPSASPKORVLIIVGAKFGEMYLNAFMQPPEGLELVGLLAQGSARSRELAHAFGIP
 PchG 1 -----MSDVRSVVVAGSRFGQFYA-AGVAADPRFVLRGILGQSRRSRALAERLGVE
 NrpU 1 -----MKPAKVVIVGSKFGEVYLNAFIQODKWHLLGLFSKGSRSRELSKAFGIP
 consensus 1 msp rrVvivGskfGemYlnAfmqp erf LvGllaqGS RSReLahafGip

YbtU 56 LYTSPEQITRMPDIACIVVRSIVAGGTGTQLARHFLTRGVHVIQEHPLHPDDISSLOQLA
 PchG 52 LWCEVEALPDDVRLACVAVGGAARCEQGPALAEALMARGIDVLIIEHPLLPREWQDLLRSA
 NrpU 53 LE TRFDQLPEKIDLACIVVRSATVGGEGSOLAQAFLQRGISVWQEHVPVHPDEITRLQSLA
 consensus 61 lyt eqlpd vdlACivVrsav Gg GtqLA afl RGi ViqEHPlhPdeis LqtLA

trans-

YbtU 116 QEQGCCYWNNTFYPHTRAGRITWRDAQQLRECLAKTPPV-VHATTSRQLYSTLDLILL
 PchG 112 -ERIGRFCHLNTFYPLPAVARFIELGRQLHRRRCIRHLDAACGV---QVGFATLDILAA
 NrpU 113 -EKMHCHYVNSLYPHNKAGRLWIENT-QKIYQQIQQRPVWQIITTSRQLIYSALDIYCO
 consensus 121 ErlgcrylvNtfYPh rAgr wie a Qlrr a hpv a avtsrQllystLDil

membrane

YbtU 174 ALGVDAAAVECDVVGVSFSDFHCLRLFWPEGEACLLQRYLDPDDPDMHSLIMHRILLGWP
 PchG 168 LLEGVCPWSLESPSNOLSAMRGLSLVLAEVPLSLHVLNLEAAAADDGRMILLQRVSLTTDR
 NrpU 171 AMKIHFNDDITVILEKDNTPILQFLRLSNPTGDLLELCLQKHLSSNDPQHSVMHHMILLGWP
 consensus 181 al v a v sv d s mh LrL pegel L lqr L Dpd hsLimhrlllgwp

YbtU 234 ECHLSLEASYPVWSSSLFVADHQBNAHSIYRRPEITRDLPGLTRSA--PLSWRDCCE
 PchG 228 -GTLSSLSPHGPILWTPAVPAEDDDGLFALFDEIAGEPLPSAQLWYAE-PCSWAQVHC
 NrpU 231 AGYLHLASYPVWNNALYIHHQDSKKAMYQSPATWELDEPLFHSFHTPPNSWQDVME
 consensus 241 G LsL asyGPviWs alfv hqd ah ly p le lp l rsya P SW dv e

YbtU 292 TVGPECVSWLLHQLRSH--LAGHPAACOSVHOTALSRLWQOILRKTGNAEIRR--LTP
 PchG 286 RHWPAAAAALALADG----DEVRRNRORSLEVA--LWQIRGERLGFPEAPPASLAP
 NrpU 291 CEAPEAINYLLAEIDKCWQLPNDKKEMILOPHYQLALSQWLWIKTLQTAGKAITDGT--IAP
 consensus 301 vgPeav wLa l l dh p Q qiAls LWqril k G ae laP

YbtU 348 PTHDRLAGFYNDDEKRAL 365
 PchG 339 ASLEQVLEQAS----- 349
 NrpU 349 FK--RMN-FTKSSGRRK- 362
 consensus 361 h drl f k

FIG. 3. Amino acid sequence alignment of YbtU from *Y. pestis*, PchG from *P. aeruginosa*, and NrpU from *P. mirabilis*. Black background, residues with identity to YbtU; gray background, conservative amino acid substitutions. The consensus line shows identical residues in all three proteins (capital letters), identical residues in two proteins (lowercase letters), and conservative amino acids in two proteins (colons). A putative transmembrane region is labeled and overlined.

ing (39.7% identity and 56.6% similarity), AngT is encoded within the anguibactin siderophore gene cluster in *V. anguillarum* (36.7% identity and 57.7% similarity), PikAV is required for macrolide antibiotic biosynthesis in *Streptomyces venezuelae* (33% identity and 50.9% similarity), and Srf4 (or SrfA-TE; 26.2% identity and 50.6% similarity) of *Bacillus subtilis* is needed for surfactin biogenesis (17, 20, 28, 66, 67). PchC, a putative thioesterase of the pyochelin siderophore system of *P. aeruginosa* (57), has 29.6% identity and 47.2% homology to YbtT. Many bacterial NRPS systems possess a C-terminal thioesterase domain as part of the NRPS in addition to a separate

gene encoding an external thioesterase. In most systems, the biochemical function of the external thioesterase is unknown. However, it is thought to serve an editing function by removing aberrant structures on mischarged NRPSs caused by nonspecific thioesterification (11, 42, 55). Thus, loss of AngT expression reduced anguibactin synthesis to 30 to 40% of wild-type levels (66) and deletion of the gene encoding SrfA-TE caused a sixfold drop in synthesis of surfactin (55). Deletion of *tylO*, encoding an external putative thioesterase in *Streptomyces fradiae*, reduced tylosin synthesis by 85% (11). Our bioassay detects Ybt in iron-deficient culture supernatants diluted 1:16

(data not shown). Thus, if any Ybt siderophore is produced by the YbtT⁻ mutant, it is at less than 6% of wild-type levels. Therefore, loss of the YbtT putative external thioesterase appears to have a slightly greater effect on product synthesis than do mutations in external thioesterase genes in other systems. In this regard the YbtT⁻ mutant resembles the *S. venezuelae* *pikAV* mutant where biosynthesis of three macrolide antibiotics was reduced to less than 5%. The *S. venezuelae* macrolide biosynthesis gene cluster has no product containing an internal thioesterase domain (67). A thioesterase domain, present in the C terminus of HMWP1, is hypothesized to release the completed Ybt siderophore from the enzyme complex (29). Experiments to test the role of the HMWP1 thioesterase domain in Ybt production are in progress.

Our results show that a $\Delta ybtU$ mutation affects Ybt synthesis and regulation. The YbtU⁻ mutant had greatly reduced transcription from the *psn* and *ybtP* promoters and reduced levels of HMWP1, HMWP2, YbtE, and Psn proteins. However, it is unlikely that YbtU is a direct regulator; the regulatory effects of the $\Delta ybtU$ mutation and polar mutations in the upstream genes *irp1* and *irp2* are corrected by the addition of the Ybt siderophore (2, 47). This suggests that the regulatory effects in these mutants are caused by loss of siderophore synthesis. The role of YbtU in siderophore synthesis is unknown. Protein motif and homology searches have not provided insight into the function of this protein. TopPred2 and DAS (<http://www/biohemi.su.se>) predict that YbtU has one potential transmembrane domain with a significant score (Fig. 3). YbtU is predicted by PSORT (45) to be an inner membrane protein. The two highest homologies in the database (3 January 2000) are to PchG of *P. aeruginosa* (pyochelin biosynthesis) and NrpU of *P. mirabilis* (swarming phenotype) (28, 57) (Fig. 3 shows the alignment), whose functions are also unknown.

In *Y. pestis*, we proposed that Ybt functions as a signal molecule by binding to the AraC-type regulator YbtA. The Ybt-YbtA complex then activates transcription of other genes in the Ybt system and represses transcription of *ybtA*. Thus mutants in *ybtA* had reduced β -galactosidase activity from a *psn::lacZ* reporter plasmid but elevated expression from a *ybtA::lacZ* reporter (22). In addition, strains bearing mutations in *irp2* or *ybtE*, both genes encoding products involved in Ybt synthesis, had significantly reduced expression of HMWP1, HMWP2, and Psn (2). Curiously, mutations in *psn* (encoding the OM receptor for Ybt) and *ybtP* (encoding a putative ABC transporter involved in Ybt uptake) do not affect regulation of gene transcription or the levels of Ybt proteins (2, 22, 24, 47). This is in contrast to results obtained with *Y. enterocolitica*, where a strain with a mutant OM receptor, designated FyuA, exhibited reduced levels of HMWP2, suggesting the involvement of FyuA in regulation. However, despite a reduction in HMWP2, a key component in Ybt biosynthesis, this mutant has been described as a Ybt overproducer (46, 51). Psn and FyuA are 98% identical at the deduced amino acid level, making it unlikely that sequence differences would account for the observed regulatory discrepancies between *Y. enterocolitica* and *Y. pestis* (24, 51). An ERIC sequence is present in the *ybtA* promoter region of *Y. enterocolitica* IB strains (52) but absent from *Y. pestis* KIM6+ (29). The effect of this insertion on expression of YbtA has not been experimentally determined (52). Alternatively, the dissimilarity between *Y. pestis* and *Y. enterocolitica* Ybt receptor mutants could be due to differential permeation of Ybt through the OM. Bengoechea et al. (5) concluded that the OM of *Y. pestis* was more permeable to small hydrophobic molecules than the OM of *Y. enterocolitica*. Thus *Y. enterocolitica* FyuA may be required to translocate Ybt to the periplasm because of the increased OM impermeability.

The effects of a $\Delta ybtU2071$ mutation on expression of *ybt* genes are similar to those observed with other Ybt biosynthetic mutants (*irp1*, *irp2*, and *ybtE*) and support the hypothesis that the siderophore functions as a regulatory molecule. However, the lack of regulatory effects caused by $\Delta ybtT2072$ and *ybtS::kan2070.1* appears to contradict this since neither mutant made detectable levels of Ybt. YbtT is a potential proofreading thioesterase, while YbtS is hypothesized to synthesize salicylate, the first moiety activated to initiate chain elongation and Ybt synthesis (29). YbtE catalyzes the adenylation of salicylate and transfer of this activated group to the N-terminal aryl carrier protein domain of HMWP2. Albeit at a much lower efficiency, YbtE also adenylates 2,3-dihydroxybenzoate (30). Thus, in the YbtS⁻ mutant YbtE may activate 2,3-dihydroxybenzoate or another phenolate compound to initiate synthesis of an aberrant Ybt molecule. The low efficiency of YbtE-catalyzed activation or chain elongation from the altered phenolate moiety may lead to low levels of the altered siderophore that might interact with YbtA and thus allow normal regulation of the Ybt system. Similarly, YbtT⁻ mutants may produce an aberrant compound(s) that can function as an inducer(s) in concert with YbtA. The altered Ybt molecule must either be produced at low levels or be retained within the bacterial cells since this molecule was not detected by our HPLC analysis. Alternatively, it is possible that small amounts of Ybt are synthesized by the YbtT⁻ mutant and are sufficient for normal regulation but not for growth stimulation. Further experiments will be necessary to determine the nature of the signal molecule in Ybt⁺ cells as well as in YbtT⁻ and YbtS⁻ mutants.

ACKNOWLEDGMENTS

This work was supported by Public Health Service grant AI42738 from the National Institutes of Health.

We thank Scott W. Bearden for assistance with some sequence analyses and construction of pPSN345.

REFERENCES

- Ausubel, F. M., R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith, and K. Struhl (ed.). 1987. Current protocols in molecular biology. John Wiley & Sons, New York, N.Y.
- Bearden, S. W., J. D. Fetherston, and R. D. Perry. 1997. Genetic organization of the yersiniabactin biosynthetic region and construction of avirulent mutants in *Yersinia pestis*. Infect. Immun. **65**:1659-1668.
- Bearden, S. W., and R. D. Perry. 1999. The Yfe system of *Yersinia pestis* transports iron and manganese and is required for full virulence of plague. Mol. Microbiol. **32**:403-414.
- Bearden, S. W., T. M. Staggs, and R. D. Perry. 1998. An ABC transporter system of *Yersinia pestis* allows utilization of chelated iron by *Escherichia coli* SAB11. J. Bacteriol. **180**:1135-1147.
- Bengoechea, J.-A., K. Brandenburg, U. Seydel, R. Díaz, and I. Moriyón. 1998. *Yersinia pseudotuberculosis* and *Yersinia pestis* show increased outer membrane permeability to hydrophobic agents which correlates with lipopolysaccharide acyl-chain fluidity. Microbiology **144**:1517-1526.
- Birnboim, H. C., and J. Doly. 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. Nucleic Acids Res. **7**:1513-1523.
- Braun, V., K. Hantke, and W. Köster. 1998. Bacterial iron transport: mechanisms, genetics, and regulation. Met. Ions Biol. Syst. **35**:67-145.
- Buchrieser, C., R. Brosch, S. Bach, A. Guiyoule, and E. Carniel. 1998. The high-pathogenicity island of *Yersinia pseudotuberculosis* can be inserted into any of the three chromosomal *asn* tRNA genes. Mol. Microbiol. **30**:965-978.
- Buchrieser, C., M. Prentice, and E. Carniel. 1998. The 102-kilobase unstable region of *Yersinia pestis* comprises a high-pathogenicity island linked to a pigmentation segment which undergoes internal rearrangement. J. Bacteriol. **180**:2321-2329.
- Buchrieser, C., C. Rusniok, L. Frangeul, E. Couve, A. Billault, F. Kunst, E. Carniel, and P. Glaser. 1999. The 102-kilobase *pgm* locus of *Yersinia pestis*: sequence analysis and comparison of selected regions among different *Yersinia pestis* and *Yersinia pseudotuberculosis* strains. Infect. Immun. **67**:4851-4861.
- Butler, A. R., N. Bate, and E. Cundliffe. 1999. Impact of thioesterase activity on tylosin biosynthesis in *Streptomyces fradiae*. Chem. Biol. **6**:287-292.
- Byers, B. R., and E. L. Arceneaux. 1998. Microbial iron transport: iron acquisition by pathogenic microorganisms. Met. Ions Biol. Syst. **35**:37-66.

13. Carniel, E., I. Guilvout, and M. Prentice. 1996. Characterization of a large chromosomal "high-pathogenicity island" in biotype 1B *Yersinia enterocolitica*. J. Bacteriol. **178**:6743–6751.
14. Carniel, E., A. Guiyoule, I. Guilvout, and O. Mercereau-Puijalon. 1992. Molecular cloning, iron-regulation and mutagenesis of the *irp2* gene encoding HMWP2, a protein specific for the highly pathogenic *Yersinia*. Mol. Microbiol. **6**:379–388.
15. Chambers, C. E., D. D. McIntyre, M. Mouck, and P. A. Sokol. 1996. Physical and structural characterization of yersiniophore, a siderophore produced by clinical isolates of *Yersinia enterocolitica*. Biometals. **9**:157–167.
16. Chang, A. C. Y., and S. N. Cohen. 1978. Construction and characterization of amplifiable multicopy DNA cloning vehicles derived from the P15A cryptic miniplasmid. J. Bacteriol. **134**:1141–1156.
17. Cosmina, P., R. Rodriguez, F. de Ferrá, G. Grandi, M. Perego, G. Venema, and D. van Sinderen. 1993. Sequence and analysis of the genetic locus responsible for surfactin synthesis in *Bacillus subtilis*. Mol. Microbiol. **8**:821–831.
18. Cox, C. D., K. L. Rinehart, Jr., M. L. Moore, and C. J. Cook, Jr. 1981. Pyochelin: novel structure of an iron-chelating growth promoter for *Pseudomonas aeruginosa*. Proc. Natl. Acad. Sci. USA **78**:4256–4260.
19. Drechsel, H., H. Stephan, R. Lotz, H. Haag, H. Zähler, K. Hantke, and G. Jung. 1995. Structure elucidation of yersiniabactin, a siderophore from highly virulent *Yersinia* strains. Liebig's Ann. **1995**:1727–1733.
20. Farrell, D. H., P. Mikesell, L. A. Actis, and J. H. Crosa. 1990. A regulatory gene, *angR*, of the iron uptake system of *Vibrio anguillarum*: similarity with phage P22 *cro* and regulation by iron. Gene **86**:45–51.
21. Ferber, D. M., J. M. Fowler, and R. R. Brubaker. 1981. Mutations to tolerance and resistance to pesticin and colicins in *Escherichia coli*. J. Bacteriol. **146**:506–511.
22. Fetherston, J. D., S. W. Bearden, and R. D. Perry. 1996. YbtA, an AraC-type regulator of the *Yersinia pestis* pesticin/yersiniabactin receptor. Mol. Microbiol. **22**:315–325.
23. Fetherston, J. D., V. J. Bertolino, and R. D. Perry. 1999. YbtP and YbtQ: two ABC transporters required for iron uptake in *Yersinia pestis*. Mol. Microbiol. **32**:289–299.
24. Fetherston, J. D., J. W. Lillard, Jr., and R. D. Perry. 1995. Analysis of the pesticin receptor from *Yersinia pestis*: role in iron-deficient growth and possible regulation by its siderophore. J. Bacteriol. **177**:1824–1833.
25. Fetherston, J. D., and R. D. Perry. 1994. The pigmentation locus of *Yersinia pestis* KIM6+ is flanked by an insertion sequence and includes the structural genes for pesticin sensitivity and HMWP2. Mol. Microbiol. **13**:697–708.
26. Fetherston, J. D., P. Schuetze, and R. D. Perry. 1992. Loss of the pigmentation phenotype in *Yersinia pestis* is due to the spontaneous deletion of 102 kb of chromosomal DNA which is flanked by a repetitive element. Mol. Microbiol. **6**:2693–2704.
27. Froehlich, B., L. Husmann, J. Caron, and J. R. Scott. 1994. Regulation of *ms*, a positive regulatory factor for pili of enterotoxigenic *Escherichia coli*. J. Bacteriol. **176**:5385–5392.
28. Gaisser, S., and C. Hughes. 1997. A locus coding for putative non-ribosomal peptide/polyketide synthase functions is mutated in a swarming-defective *Proteus mirabilis* strain. Mol. Gen. Genet. **253**:415–427.
29. Gehring, A. M., E. DeMoll, J. D. Fetherston, I. Mori, G. F. Mayhew, F. R. Blattner, C. T. Walsh, and R. D. Perry. 1998. Iron acquisition in plague: modular logic in enzymatic biogenesis of yersiniabactin by *Yersinia pestis*. Chem. Biol. **5**:573–586.
30. Gehring, A. M., I. Mori, R. D. Perry, and C. T. Walsh. 1998. The nonribosomal peptide synthetase HMWP2 forms a thiazoline ring during biogenesis of yersiniabactin, an iron-chelating virulence factor of *Yersinia pestis*. Biochemistry **37**:11637–11650.
31. Guilvout, I., O. Mercereau-Puijalon, S. Bonnefoy, A. P. Pugsley, and E. Carniel. 1993. High-molecular-weight protein 2 of *Yersinia enterocolitica* is homologous to AngR of *Vibrio anguillarum* and belongs to a family of proteins involved in nonribosomal peptide synthesis. J. Bacteriol. **175**:5488–5504.
32. Hare, J. M., and K. A. McDonough. 1999. High-frequency RecA-dependent and -independent mechanisms of Congo red binding mutations in *Yersinia pestis*. J. Bacteriol. **181**:4896–4904.
33. Hare, J. M., A. K. Wagner, and K. A. McDonough. 1999. Independent acquisition and insertion into different chromosomal locations of the same pathogenicity island in *Yersinia pestis* and *Yersinia pseudotuberculosis*. Mol. Microbiol. **31**:291–303.
34. Heesemann, J., K. Hantke, T. Vocke, E. Saken, A. Rakin, I. Stojiljkovic, and R. Berner. 1993. Virulence of *Yersinia enterocolitica* is closely associated with siderophore production, expression of an iron-repressible outer membrane polypeptide of 65 000 Da and pesticin sensitivity. Mol. Microbiol. **8**:397–408.
35. Hohn, B., and J. Collins. 1980. A small cosmid for efficient cloning of large DNA fragments. Gene **11**:291–298.
36. Hornung, J. M., H. A. Jones, and R. D. Perry. 1996. The *hmu* locus of *Yersinia pestis* is essential for utilization of free haemin and haem-protein complexes as iron sources. Mol. Microbiol. **20**:725–739.
37. Humphreys, G. O., G. A. Willshaw, and E. S. Anderson. 1975. A simple method for the preparation of large quantities of pure plasmid DNA. Biochim. Biophys. Acta **383**:457–463.
38. Jalal, M. A. F., M. B. Hossain, D. van der Helm, J. Sanders-Loehr, L. A. Actis, and J. H. Crosa. 1989. Structure of anguibactin, a unique plasmid-related bacterial siderophore from the fish pathogen *Vibrio anguillarum*. J. Am. Chem. Soc. **111**:292–296.
39. Kaniga, K., I. Delor, and G. R. Cornelis. 1991. A wide-host-range suicide vector for improving reverse genetics in Gram-negative bacteria: inactivation of the *blaA* gene of *Yersinia enterocolitica*. Gene **109**:137–141.
40. Kutnyrev, V. V., A. A. Filippov, O. S. Oparina, and O. A. Protchenko. 1992. Analysis of *Yersinia pestis* chromosomal determinants Pgm⁺ and Pst^S associated with virulence. Microb. Pathog. **12**:177–186.
41. Lucier, T. S., and R. R. Brubaker. 1992. Determination of genome size, macrorestriction pattern polymorphism, and nonpigmentation-specific deletion in *Yersinia pestis* by pulsed-field gel electrophoresis. J. Bacteriol. **174**:2078–2086.
42. Marahiel, M. A., T. Stachelhaus, and H. D. Mootz. 1997. Modular peptide synthetases involved in nonribosomal peptide synthesis. Chem. Rev. **97**:2651–2673.
43. Mietzner, T. A., and S. A. Morse. 1994. The role of iron-binding proteins in the survival of pathogenic bacteria. Annu. Rev. Nutr. **14**:471–493.
44. Miller, J. H. 1992. A short course in bacterial genetics. A Laboratory Manual and Handbook for *Escherichia coli* and Related Bacteria. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
45. Nakai, K., and M. Kanehisa. 1991. Expert system for predicting protein localization sites in Gram-negative bacteria. Proteins **11**:95–110.
46. Pelludat, C., A. Rakin, C. A. Jacobi, S. Schubert, and J. Heesemann. 1998. The yersiniabactin biosynthetic gene cluster of *Yersinia enterocolitica*: organization and siderophore-dependent regulation. J. Bacteriol. **180**:538–546.
47. Perry, R. D., P. B. Balbo, H. A. Jones, J. D. Fetherston, and E. DeMoll. 1999. Yersiniabactin from *Yersinia pestis*: biochemical characterization of the siderophore and its role in iron transport and regulation. Microbiology **145**:1181–1190.
48. Perry, R. D., and J. D. Fetherston. 1997. *Yersinia pestis*—etiologic agent of plague. Clin. Microbiol. Rev. **10**:35–66.
49. Perry, R. D., S. C. Straley, J. D. Fetherston, D. J. Rose, J. Gregor, and F. R. Blattner. 1998. DNA sequencing and analysis of the low-Ca²⁺-response plasmid pCD1 of *Yersinia pestis* KIM5. Infect. Immun. **66**:4611–4623.
50. Rakin, A., C. Noeltgen, S. Schubert, and J. Heesemann. 1999. Common and specific characteristics of the high-pathogenicity island of *Yersinia enterocolitica*. Infect. Immun. **67**:5265–5274.
51. Rakin, A., E. Saken, D. Harmsen, and J. Heesemann. 1994. The pesticin receptor of *Yersinia enterocolitica*: a novel virulence factor with dual function. Mol. Microbiol. **13**:253–263.
52. Rakin, A., S. Schubert, C. Pelludat, D. Brem, and J. Heesemann. 1999. The high-pathogenicity island of yersiniae, p. 77–90. In J. B. Kaper and J. Hacker (ed.), Pathogenicity islands and other mobile virulence elements. American Society for Microbiology, Washington, D.C.
53. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
54. Sanger, F., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. Proc. Natl. Acad. Sci. USA **74**:5463–5467.
55. Schneider, A., and M. A. Marahiel. 1998. Genetic evidence for a role of thioesterase domains, integrated in or associated with peptide synthetases, in non-ribosomal peptide biosynthesis in *Bacillus subtilis*. Arch. Microbiol. **169**:404–410.
56. Schubert, S., A. Rakin, H. Karch, E. Carniel, and J. Heesemann. 1998. Prevalence of the "high-pathogenicity island" of *Yersinia* species among *Escherichia coli* strains that are pathogenic to humans. Infect. Immun. **66**:480–485.
57. Serino, L., C. Reimann, P. Visca, M. Beyeler, V. Della Chiesa, and D. Haas. 1997. Biosynthesis of pyochelin and dihydroaeruginic acid requires the iron-regulated *pchDCBA* operon in *Pseudomonas aeruginosa*. J. Bacteriol. **179**:248–257.
58. Shaw-Reid, C., N. Kelleher, H. C. Losey, A. M. Gehring, C. Berg, and C. T. Walsh. 1999. Assembly line enzymology by multimodular nonribosomal peptide synthetases: the thioesterase domain of *Escherichia coli* EntF catalyzes both elongation and cyclolactonization. Chem. Biol. **6**:385–400.
59. Sikkema, D. J., and R. R. Brubaker. 1987. Resistance to pesticin, storage of iron, and invasion of HeLa cells by yersiniae. Infect. Immun. **55**:572–578.
60. Staggs, T. M., J. D. Fetherston, and R. D. Perry. 1994. Pleiotropic effects of a *Yersinia pestis fur* mutation. J. Bacteriol. **176**:7614–7624.
61. Staggs, T. M., and R. D. Perry. 1991. Identification and cloning of a *fur* regulatory gene in *Yersinia pestis*. J. Bacteriol. **173**:417–425.
62. Suo, Z., C. T. Walsh, and D. A. Miller. 1999. Tandem heterocyclization activity of the multidomain 230 kDa HMWP2 subunit of *Yersinia pestis* yersiniabactin synthetase: interaction of the 1-1382 and 1383-2035 fragments. Biochemistry **38**:14023–14035.
63. Surgalla, M. J., and E. D. Beesley. 1969. Congo red-agar plating medium for detecting pigmentation in *Pasteurella pestis*. Appl. Microbiol. **18**:834–837.

64. **Thompson, J. M., H. A. Jones, and R. D. Perry.** 1999. Molecular characterization of the hemin uptake locus (*hmu*) from *Yersinia pestis* and analysis of *hmu* mutants for hemin and hemoprotein utilization. *Infect. Immun.* **67**: 3879–3892.
65. **Weinberg, E. D., and G. A. Weinberg.** 1995. The role of iron in infection. *Curr. Opin. Infect. Dis.* **8**:164–169.
66. **Wertheimer, A. M., W. Verweij, Q. Chen, L. M. Crosa, M. Nagasawa, M. E. Tolmasky, L. A. Actis, and J. H. Crosa.** 1999. Characterization of the *angR* gene of *Vibrio anguillarum*: essential role in virulence. *Infect. Immun.* **67**: 6496–6509.
67. **Xue, Y., L. Zhao, H. Liu, and D. H. Sherman.** 1998. A gene cluster for macrolide antibiotic biosynthesis in *Streptomyces venezuelae*: architecture of metabolic diversity. *Proc. Natl. Acad. Sci. USA* **95**:12111–12116.
68. **Yanisch-Perron, C., J. Vieira, and J. Messing.** 1985. Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**:103–119.

Editor: J. T. Barbieri