

Experimental Q Fever Infection in Congenitally Athymic Nude Mice

R. A. KISHIMOTO,* H. ROZMIAREK, AND E. W. LARSON

U.S. Army Medical Research Institute of Infectious Diseases, Fort Detrick, Frederick, Maryland 21701

Received for publication 14 April 1978

Congenitally athymic nude (nu/nu) mice and their phenotypically normal (nu/+) euthymic littermates were exposed to *Coxiella burnetii* administered as small-particle aerosols. After challenge, both strains of mice became infected, as characterized by rickettsemia, viable rickettsiae in the spleen, and serological conversion. The major difference noted was that euthymic animals had cleared rickettsiae from peripheral circulation and the spleen within 14 days. In contrast, rickettsiae were detected and isolated from spleen and blood of athymic mice through 60 days.

The host defense mechanisms involved in successful elimination of *Coxiella burnetii* from experimentally infected animals have not been fully elucidated. Previous *in vivo* studies suggest a role for immune serum. Abinanti and Marmion (1) showed that although immune serum is not known to have a direct rickettsicidal action on *C. burnetii*, passive transfer of serum from hyperimmune donors modified the infection in nonimmune animals. Although serum transfer can abrogate an infective challenge, the importance of specific antibody to an active Q fever infection is not known.

In addition to antibody, a role for cell-mediated immunity in experimental Q fever had also been suggested. Peritoneal macrophages obtained from guinea pigs previously vaccinated with killed phase I *C. burnetii* were capable of phagocytizing and killing homologous rickettsiae in the absence of immune serum *in vitro* (13). Kazár et al. (10) demonstrated that spleen cells taken from mice immunized with live phase I *C. burnetii* transferred resistance.

Whereas recent studies have shown thymus rudiment in athymic mice (15) and high frequency of T-lineage lymphocytes in spleen (9), these animals have been shown to be devoid of functional T lymphocytes, which are required for the expression of cell-mediated immunity (16-18). Thus, the nude mouse provides a means for studying the role of cell-mediated immunity in host defense against Q fever.

This study was undertaken to assess further the relative importance of cellular and humoral immune systems in experimental Q fever infections. Congenitally athymic nude (nu/nu) mice and their phenotypically normal (nu/+) euthymic littermates were infected with *C. burnetii*

to elucidate the role of humoral and thymus-dependent cell-mediated immunity resistance to Q fever infection.

MATERIALS AND METHODS

Preparation of rickettsial stock suspension. The third egg passage of the Henzerling strain of *C. burnetii* in phase I was grown in chicken embryo cells as previously described (13). The infectivity of the rickettsial suspension was estimated to be $10^{9.5}$ mouse median intraperitoneal infectious doses.

Animals. Outbred Swiss (nu/nu) athymic mice and their phenotypically normal euthymic (nu/+) littermates, weighing 18 to 22 g, were obtained from the animal breeding facilities, Frederick Cancer Research Center. These animals had been bred and raised in a pathogen-free environment and, during the experimental period, were housed in gas-tight, total-containment isolation facilities. All food (Purina 5010, Ralston Purina Co., St. Louis, Mo.), bedding, and water acidified to pH 2.5 were sterilized prior to use.

Infection of mice. Mice were exposed to 10^4 mouse median intraperitoneal infectious doses of the phase I Henzerling strain of *C. burnetii* presented in small-particle aerosols as previously described (11). Mice exposed to aerosols of sterile Earle medium 199 served as controls.

Serological assays. Serum antibody activity against phase I and II *C. burnetii* was evaluated at weekly intervals by the indirect immunofluorescent technique of Bozeman and Elisberg (2).

Rickettsemia. Blood from both strains of mice was collected at selected intervals and inoculated intraperitoneally into Swiss-Webster mice. The animals were bled 21 days later, and the presence of specific antibodies in serum was determined.

Spleen impression smears. The presence of *C. burnetii* in spleens of three to five mice was determined microscopically after staining two impression smears with Giménez stain (6). Examination of two separate impression smears by the direct immunoflu-

orescent antibody technique (7) was used as a confirmatory test. The amount of rickettsiae in spleens was estimated in 10 different fields of view ($\times 1,000$) and scored according to the method of Kazár et al. (10).

Viability of rickettsiae. The viability of rickettsiae in spleens was determined by inoculation of various dilutions of spleen homogenates into the yolk sac of 5-day-old chicken embryonated eggs. The surviving eggs were harvested after 10 days of incubation, and yolk sac smears were stained with Giménez stain and fluorescein isothiocyanate-conjugated phase I and II *C. burnetii* antiserum.

RESULTS

Clinical signs and gross pathology. No clinical signs of illness (fever, weight loss, lethargy, coryza) or mortality were observed in any of the athymic or euthymic mice after exposure to *C. burnetii*. Gross lung lesions were observed from 7 through 14 days. These lesions appeared to be resolving by 21 days in both strains of mice. The spleen weight began to increase on day 5 and continued to increase until day 14 in both types of mice (Fig. 1). However, the spleen weight in euthymic mice declined to normal levels by day 21, whereas the spleen weights of athymic mice continued to increase; by day 30 the spleen weights were approximately 11 times heavier than those of control animals. On day 60 the weight of the spleen of athymic mice was approximately five times that of controls.

Isolation of rickettsiae from spleen and blood. The spleens from both strains of mice contained rickettsiae between 7 and 14 days, as shown by the impression smears (Fig. 1). Rickettsiae were no longer detected in euthymic animals on day 21, but uncountable numbers of organisms per each field of view were observed in athymic mice for 60 days, at which time the experiment was terminated. Titration of infected spleens in yolk sacs of embryonated eggs indicated that the spleens of athymic mice contained

approximately 10^5 to 10^6 rickettsiae at 30 and 60 days, whereas no rickettsiae were recovered from euthymic animals at these times.

Rickettsiae were isolated from the blood of both athymic and euthymic mice on day 4, but organisms were no longer detected in euthymic mice by 21 days (Fig. 2). In contrast, rickettsemia persisted in athymic mice through day 60.

Antibody response. Antibody to phase I and II *C. burnetii* developed at the same rate in both strains of mice (Fig. 2). Antibody against phase II antigen was initially detected on day 14 and persisted at elevated titers through day 60. Antibody to phase I antigen was demonstrable by 21 days and persisted at high levels. Uninfected control athymic and euthymic mice did not develop antibodies to either antigen.

DISCUSSION

These data demonstrate that both athymic and euthymic mice are infected after an aerosol exposure to 10^4 mouse intraperitoneal infectious doses of *C. burnetii*. The major difference noted was that euthymic animals were capable of clearing rickettsiae from peripheral blood and the spleen by 14 days, but athymic mice became chronically infected and did not eliminate the organisms. Studies by other investigators have also shown that infection of athymic nude mice with other intracellular microorganisms resulted in chronic infections (4, 5).

Although it is not known at this time what mechanism(s) underlies the observed phenomena, it appears that the T lymphocytes may play a role in terminating Q fever infection in normal animals. Efforts are currently underway in our laboratory to determine the specific requirement for T lymphocytes by means of adoptive transfer studies. It is possible that soluble products of T




FIG. 1. Spleen weights and impression smears of athymic and euthymic mice after challenge with *C. burnetii*. Each point represents the mean of three to five animals.




FIG. 2. Onset and duration of rickettsemia and indirect fluorescent antibody titers of athymic and euthymic mice after challenge with *C. burnetii*. Each point represents the mean of three to five animals.

cells (14) or macrophage inhibition factor can activate macrophages and remove rickettsiae. Studies by Hinrichs and Jerrells (8) provide evidence that normal guinea pig peritoneal macrophages cultured in vitro with immune lymphocytes from Q fever-infected guinea pigs or with supernatant fluid rich in macrophage inhibition factor inhibited the growth of ingested *C. burnetii*.

It appears that active, nonspecific defense mechanisms were present in Q fever-infected athymic mice, since none of the animals died after challenge. However, this response was not sufficient to terminate infection, because the mice were chronically infected. Cheers and Waller (3) and Emmerling et al. (5) noted that macrophages of nude mice showed increased resistance during the early course of *Listeria monocytogenes* infection. However, these animals also were not able to control and terminate infection, as did the phenotypically normal control mice.

Our results demonstrate that both athymic and euthymic mice are capable of producing antibodies against phase I and II *C. burnetii*. Since athymic mice were able to produce specific antibodies, it can be inferred that *C. burnetii* antigen elicits a T-cell-independent humoral response. This was not totally unexpected, since the cell envelope of phase I *C. burnetii* is composed of polysaccharide, fatty acids, and protein. This antibody may act as an opsonin which then potentiates the destruction of rickettsiae by macrophages (12, 13). However, antibody in concert with macrophages are not sufficient to terminate infection.

We believe that the athymic mouse would be a useful model for chronic and persistent Q fever studies.

ACKNOWLEDGMENTS

We thank Robert Stockman and Chester Redmond for technical assistance.

LITERATURE CITED

1. Abinanti, F. R., and B. P. Marmion. 1957. Protective or neutralizing antibody in Q fever. *Am. J. Hyg.* **66**:173-195.
2. Bozeman, F. M., and B. L. Elisberg. 1963. Serological diagnosis of scrub typhus by indirect immunofluorescence. *Proc. Soc. Exp. Biol. Med.* **112**:568-573.
3. Cheers, C., and R. Waller. 1975. Activated macrophages in congenitally athymic "nude" mice and in lethally irradiated mice. *J. Immunol.* **115**:844-847.
4. Clark, I. A., and A. C. Allison. 1974. *Babesia microti* and *Plasmodium berghei yoelii* infections in nude mice. *Nature (London)* **252**:328-329.
5. Emmerling, P., H. Finger, and J. Bockenmühl. 1975. *Listeria monocytogenes* infection in nude mice. *Infect. Immun.* **12**:437-439.
6. Giménez, D. F. 1964. Staining rickettsiae in yolk-sac cultures. *Stain Technol.* **39**:135-140.
7. Hahon, N., and K. O. Cooke. 1966. Assay of *Coxiella burnetii* by enumeration of immunofluorescent infected cells. *J. Immunol.* **97**:492-497.
8. Hinrichs, D. J., and T. R. Jerrells. 1976. In vitro evaluation of immunity to *Coxiella burnetii*. *J. Immunol.* **117**:996-1003.
9. Holub, M., P. Rossman, H. Tlaskalova, and H. Vidmarova. 1975. Thymus rudiment of the athymic nude mouse. *Nature (London)* **256**:491-493.
10. Kazár, J., E. El-Najdawi, R. Brezina, and S. Schrammek. 1977. Search for correlates of resistance to virulent challenge in mice immunized with *Coxiella burnetii*. *Acta Virol.* **21**:422-430.
11. Kishimoto, R. A., and G. T. Burger. 1977. Appearance of cellular and humoral immunity in guinea pigs after infection with *Coxiella burnetii* administered in small-particle aerosols. *Infect. Immun.* **16**:518-521.
12. Kishimoto, R. A., B. J. Veltri, P. G. Canonico, F. G. Shirey, and J. S. Walker. 1976. Electron microscopic study on the interaction between normal guinea pig peritoneal macrophages and *Coxiella burnetii*. *Infect. Immun.* **14**:1087-1096.
13. Kishimoto, R. A., and J. S. Walker. 1976. Interaction between *Coxiella burnetii* and guinea pig peritoneal macrophages. *Infect. Immun.* **14**:416-421.
14. Krahenbuhl, J. L., and J. S. Remington. 1971. In vitro induction of nonspecific resistance in macrophages by specifically sensitized lymphocytes. *Infect. Immun.* **4**:337-343.
15. Loor, F., and G. E. Roelants. 1974. High frequency of T lineage lymphocytes in nude mouse spleen. *Nature (London)* **251**:229-230.
16. Pantelouris, E. M. 1968. Absence of thymus in a mouse mutant. *Nature (London)* **217**:370-371.
17. Pantelouris, E. M. 1971. Observations on the immunobiology of 'nude' mice. *Immunology* **20**:247-252.
18. Thurman, G. B., B. B. Silver, J. A. Hooper, B. C. Giovanella, and A. L. Goldstein. 1974. In vitro mitogenic responses of spleen cells and ultrastructural studies of lymph nodes from nude mice following thymosin administration *in vivo*, p. 105-117. *In Proceedings, First International Workshop on Nude Mice.* Gustav Fischer Verlag, Stuttgart.