

Effect of Estrogens on the Myeloperoxidase-Mediated Antimicrobial System

SEYMOUR J. KLEBANOFF

Department of Medicine, University of Washington, Seattle, Washington 98195

Received for publication 12 April 1979

Estradiol 17^β prevented the fall in the microbicidal activity of the myeloperoxidase-H₂O₂-halide system induced by high H₂O₂ concentrations. In contrast, when the H₂O₂ (and halide) concentrations were low, the myeloperoxidase-H₂O₂-halide antimicrobial system was inhibited by estradiol. These properties of estradiol 17^β were shared by estradiol 17^α, estrone, estriol, ethinyl estradiol, and phenol, but not by estradiol-3-benzoate, testosterone, progesterone, hydroxyprogesterone, cortisone, hydrocortisone, or deoxycorticosterone.

A number of peroxidase-catalyzed reactions are stimulated by estrogens (9-12, 20). An intact phenolic hydroxyl group is required, and it was proposed that the estrogen may act as an oxidation-reduction catalyst, being alternately oxidized by peroxidase and H₂O₂, possibly to the phenoxy radical, and then reduced to its original form by an electron donor which is in turn oxidized. In several, but not all, instances, the electron donor is oxidized more rapidly in the presence of the estrogen than in its absence. In the absence of an appropriate electron donor, oxidation of the estrogen can lead to its further degradation with a loss of biological activity (13, 17). In the presence of a protein, firm, presumably covalent binding of the estrogen occurs (6-8, 15), and firm binding of estrogens to peroxidase-positive sites in tissues has been reported (4, 15, 19).

Myeloperoxidase (MPO), H₂O₂, and a halide form a potent antimicrobial system effective against a variety of microorganisms, and this system appears to contribute significantly to the antimicrobial activity of the polymorphonuclear leukocyte (PMN) (for reviews see 14, 16). Estradiol is converted to an alcohol-precipitable form by PMNs during phagocytosis (15). This reaction is inhibited by hypoxia and does not occur in PMNs from patients with chronic granulomatous disease, suggesting a requirement for the phagocytosis-induced respiratory burst. Binding is reduced by catalase and is increased by superoxide dismutase, an enzyme which catalyzes the conversion of the superoxide anion to oxygen and H₂O₂, implicating H₂O₂ as the active product of the respiratory burst. The peroxidase inhibitors azide and cyanide decrease binding under certain experimental conditions, and binding is low when leukocytes which lack MPO are employed, suggesting that MPO also is required.

These findings suggest that estrogens are oxidized by intact PMNs during phagocytosis, in part, as a result of the action of MPO and H₂O₂. This prompted a study of the effect of estrogens on the MPO-H₂O₂-halide antimicrobial system. A dual effect was found: a stimulation of microbicidal activity at high H₂O₂ concentrations and an inhibition when the H₂O₂ and halide concentrations were low.

MATERIALS AND METHODS

Special reagents. *Escherichia coli* (no. 11775, American Type Culture Collection, Rockville, Md.) was maintained on blood agar plates and transferred daily to Trypticase soy broth (Baltimore Biological Laboratory, Cockeysville, Md.). Overnight cultures were washed twice in 0.1 M sodium sulfate and suspended in this solution to the required absorbance at 540 nm just before use. Canine MPO was prepared from pyometrial pus by the method of Agner (1), and human MPO was kindly supplied by I. Olsson, Lund, Sweden. Canine MPO was used unless otherwise indicated. Peroxidase activity was determined by the *ortho*-dianisidine assay (21) before use. One unit of enzyme is the amount which decomposes 1 μmol of H₂O₂ per min at 25°C. Estradiol 17^β, estradiol 17^α, estrone, estriol, ethinyl estradiol, estradiol-3-benzoate, testosterone, progesterone, hydroxyprogesterone, cortisone, hydrocortisone, and deoxycorticosterone were obtained from Sigma Chemical Co., St. Louis, Mo.

Determination of microbicidal activity. The components of the reaction mixture (see table legends) in a total volume of 0.5 ml were incubated in polystyrene test tubes (12 by 75 mm, Falcon 2052, Oxnard, Calif.) at 37°C for 1 h in a water-bath shaker oscillating 80 times per min. Serial 10-fold dilutions were made in 0.1 M sodium sulfate, and the viable cell count was determined by the pour plate method using Trypticase soy agar. The viable cell count was expressed as organisms per milliliter of original reaction mixture.

Statistical analyses. The geometric mean of the microbicidal data was calculated as previously described (18). Statistical differences were determined

using Student's two-tailed *t* test for independent means (not significant, $P > 0.05$).

RESULTS

Stimulation by estrogens. Table 1 demonstrates the effect of H_2O_2 alone, the MPO- H_2O_2 -chloride system, and the latter system supplemented with estradiol on the *E. coli* viable cell count over a range of H_2O_2 concentrations. Under the conditions employed, H_2O_2 alone did not have a significant inhibitory effect on the viable cell count until a concentration of 4×10^{-3} M was reached ($P < 0.001$). When H_2O_2 was supplemented with MPO and chloride, a microbicidal effect was observed at low H_2O_2 concentrations which disappeared when the H_2O_2 concentration was increased to 5×10^{-4} M. This loss of microbicidal activity was evident until a concentration of H_2O_2 was reached at which H_2O_2 alone was toxic. Peroxidases, including MPO (2), are inhibited by H_2O_2 at high concentrations, and the loss of microbicidal activity of the MPO system presumably reflects this inhibition. When estradiol was added to the MPO- H_2O_2 -chloride system, a significant increase in microbicidal activity was observed at H_2O_2 concentrations ranging from 2×10^{-4} to 1×10^{-3} M. Bactericidal activity was stimulated under these conditions by estradiol 17 β , estradiol 17 α , estrone, estriol, ethinyl estradiol, and phenol, but not by estradiol-3-benzoate, testosterone, progesterone, hydroxyprogesterone, cortisone, hydrocortisone,

and deoxycorticosterone (Table 2, system A). Table 3 demonstrates the requirement for each component of the MPO- H_2O_2 -chloride-estra-

TABLE 2. *Effect of various agents*^a

Agent	Viable cell count (organisms/ml $\times 10^{-6}$) ^b in:	
	System A	System B
None	4.32	0.001
Estradiol 17 β	0.0 ^c	4.03 ^c
Estradiol 17 α	0.00002 ^c	6.09 ^c
Estrone	0.00002 ^c	0.2 ^c
Estriol	0.0004 ^c	5.96 ^c
Ethinyl estradiol	0.0002 ^c	6.13 ^c
Phenol	0.0001 ^c	4.51 ^c
Estradiol-3-benzoate	4.16	0.001
Testosterone	5.67	0.0002
Progesterone	4.47	0.0002
Hydroxyprogesterone	5.35	0.0003
Cortisone	6.93	0.0005
Hydrocortisone	4.62	0.0009
Deoxycorticosterone	4.97	0.001

^a The reaction mixture was as described in Table 1, except that the supplements were as follows. System A: H_2O_2 , 5×10^{-4} M; MPO, 8 mU; sodium chloride, 5×10^{-2} M; steroids (or phenol), 10^{-4} M. System B: H_2O_2 , 10^{-5} M; MPO, 8 mU; sodium iodide, 5×10^{-5} M; steroids (or phenol), 10^{-4} M.

^b Mean of three to four experiments.

^c $P < 0.001$ steroid (or phenol) versus none; all others not significant.

TABLE 3. *Effect of deletions and halide substitutions*^a

Additions	Viable cell count	
	Organisms/ml ($\times 10^{-6}$) ^b	P^c
Control	6.06	
MPO + H_2O_2 + Cl^-	0.0002	<0.001
+ estradiol		
Estradiol omitted	6.21	NS ^d
NaCl omitted	6.49	NS
H_2O_2 omitted	6.44	NS
MPO omitted	6.30	NS
MPO heated	6.30	NS
MPO + H_2O_2 + I^-	4.17	NS
Estradiol added	0.01	<0.025
MPO + H_2O_2 + Br^-	5.34	NS
Estradiol added	0.0008	<0.05

^a The MPO- H_2O_2 -chloride-estradiol system was as described in Table 1, with the H_2O_2 concentration at 5×10^{-4} M. Sodium iodide (10^{-4} M) and sodium bromide (10^{-3} M) were substituted for chloride where indicated. Deletions and additions were made, and the MPO was heated at $100^\circ C$ for 15 min where indicated.

^b Mean of three to four experiments.

^c P value for the significance of the difference from the control which contained *E. coli*, buffer, ethylenediaminetetraacetic acid, and sulfate as described in Table 1.

^d NS, Not significant.

TABLE 1. *Stimulation of the MPO- H_2O_2 -chloride antimicrobial system by estradiol*^a

H_2O_2 (M)	Viable cell count (organisms/ml $\times 10^{-6}$) ^b in:		
	H_2O_2	MPO + H_2O_2 + Cl^-	MPO + H_2O_2 + Cl^- + estradiol
1×10^{-5}		0.00003	0.0001
1×10^{-4}	7.67	0.002	0.00002
2×10^{-4}	6.92	0.34	0.00003 ^c
5×10^{-4}	6.75	6.19	0.00009 ^c
1×10^{-3}	6.08	5.95	0.005 ^c
2×10^{-3}	2.56	4.26	3.35
4×10^{-3}	0.0009	0.0007	0.0003

^a The reaction mixture contained 5×10^{-2} M sodium phosphate buffer (pH 7.5), 5×10^{-5} M ethylenediaminetetraacetic acid, 4.4×10^{-2} M sodium sulfate, 6×10^5 *E. coli*, and the supplements indicated below as follows: H_2O_2 at the concentrations indicated; MPO, 8 mU; sodium chloride, 5×10^{-2} M; and estradiol, 10^{-4} M. An isotonic equivalent amount of sodium sulfate (3.3×10^{-2} M) was omitted from tubes containing sodium chloride.

^b Mean of three to six experiments.

^c Significance of the difference from the MPO- H_2O_2 - Cl^- system, $P < 0.001$; all others not significant.

diol antimicrobial system and the loss of activity on heat-inactivation of MPO. When phenol was the stimulant employed, deletion of chloride decreased but did not abolish antimicrobial activity (data not shown).

Estradiol also increased antimicrobial activity at high H_2O_2 concentration when iodide or bromide was substituted for chloride (Table 3). As for the chloride-dependent system, a bactericidal effect of the MPO- H_2O_2 -iodide (or bromide) system was readily observed in the absence of estradiol when the H_2O_2 concentration was lowered (Table 4). Comparable results were observed when MPO was replaced by a purified preparation of guinea pig eosinophil peroxidase, i.e., microbicidal activity was inhibited by high concentrations of H_2O_2 , and this inhibition was prevented by estradiol (data not shown). The canine MPO generally employed also could be replaced by human MPO.

Inhibition by estrogens. When the H_2O_2 concentration was 10^{-5} M, the MPO- H_2O_2 -halide system was bactericidal at iodide concentrations down to 10^{-5} M under the conditions used in Table 4. In contrast to the findings at high H_2O_2 concentrations, estradiol inhibited antimicrobial activity under these conditions, with a significant inhibitory effect observed at iodide concentrations ranging from 2×10^{-4} M to 2×10^{-5} M. An inhibition by estradiol also was observed when bromide was substituted for iodide; however, when chloride (10^{-1} or 10^{-2} M) was used under the conditions used in Table 4, in-

hibition by estradiol was not seen. When both chloride (5×10^{-2} M) and iodide (5×10^{-5} M) were employed, the results were comparable to those observed with iodide alone; i.e., estradiol significantly inhibited antimicrobial activity. Those steroids which stimulated antimicrobial activity at high H_2O_2 concentration (Table 2, system A) also inhibited antimicrobial activity at low H_2O_2 concentration (Table 2, system B), although complete inhibition was not observed with estrone under the conditions employed. Phenol also was inhibitory. The canine MPO generally used could be replaced by human MPO.

DISCUSSION

Phenolic estrogens were shown here to have a dual effect on the MPO- H_2O_2 -halide antimicrobial system. When the H_2O_2 concentration was increased to a level at which loss of microbicidal activity was observed, presumably due to the inactivation of MPO (2), estrogens increased microbicidal activity. In contrast, when the H_2O_2 and halide concentrations were low, the microbicidal activity evident under these conditions was inhibited by estrogens. The same estrogens which were stimulatory at high H_2O_2 concentrations were inhibitory when the H_2O_2 concentration was low, and non-estrogenic steroids (testosterone, progesterone, hydroxyprogesterone, cortisone, hydrocortisone, deoxycorticosterone) were without effect under both sets of conditions. It is of interest that ethinyl estradiol, which has a substitution in the 17 position, was active, whereas estradiol-3-benzoate, with a substitution on the phenolic hydroxyl group, was ineffective. The importance of the phenolic hydroxyl group was further emphasized by the comparable effects observed when the phenolic estrogen was replaced by phenol. Both the 17^{α} and the 17^{β} stereoisomers of estradiol were active, as were estril and estrone. In an earlier study, Jacobs et al. (5) observed that estril and estrone increased the bactericidal activity of a system consisting of the granule pellet from guinea pig PMNs, H_2O_2 , and chloride at pH 5.5. The H_2O_2 concentration used was 5×10^{-5} M. In contrast to the findings reported here, stimulation was not observed when estradiol 17^{β} was used or when the granule pellet was replaced by purified human MPO.

MPO has two heme prosthetic groups per molecule. At relatively low concentrations, H_2O_2 forms a dissociable complex with one heme and is utilized for peroxidatic reactions. Both the halide and estrogen are oxidized by MPO under these conditions, with the oxidation of the halide required for microbicidal activity. When the

TABLE 4. Inhibition of antimicrobial activity by estradiol^a

Halide	Viable cell count (organisms/ml $\times 10^{-6}$) ^b	M			
		Type	-Estradiol	+Estradiol	P ^c
NaI	5×10^{-4}		0.00001	0.001	NS
	2×10^{-4}		0.00009	0.10	<0.002
	1×10^{-4}		0.0003	1.20	<0.001
	5×10^{-5}		0.0007	4.78	<0.001
	2×10^{-5}		0.0005	4.51	<0.01
	1×10^{-5}		0.75	4.63	NS
NaBr	5×10^{-6}		4.96	4.80	NS
	1×10^{-4}		0.0002	3.50	<0.005
NaCl	5×10^{-2}		0.00001	0.00001	NS
	1×10^{-2}		0.00009	0.0006	NS
	1×10^{-3}		5.39	5.66	NS

^a The reaction mixture was as described in Table 1 except that the supplements were as follows: H_2O_2 , 10^{-5} M; MPO, 8 mU; sodium iodide and sodium bromide at the concentrations indicated; and estradiol, 10^{-4} M where indicated.

^b Mean of two to five experiments.

^c P values for the significance of the difference between the presence and absence of estradiol. NS, Not significant.

H₂O₂ and halide concentrations are low, inhibition of microbicidal activity by estradiol may be due simply to the competition between the halide and estrogen for the small amount of H₂O₂ available. It has been proposed (2) that, when the H₂O₂ concentration is increased, complexes are formed with both hemes which interact with the degradation of H₂O₂ to oxygen and water. With a further increase in the H₂O₂ concentration, the iron porphyrin structure is destroyed and the enzyme is denatured. Under the conditions used in Table 1, estradiol was protective of the microbicidal system at concentrations 1/2 to 1/20 that of H₂O₂, indicating that its effect was not due to a stoichiometric interaction with H₂O₂. It is not known whether estrogens function as oxidation-reduction catalysts under these conditions or react directly with MPO to prevent inactivation.

Estrogens are firmly bound by PMNs during phagocytosis, and covalent linkage to cellular components, due in part to oxidation by MPO and H₂O₂, has been proposed (15). It is possible, therefore, that estrogens can influence the microbicidal activity of the MPO system in the intact cell either by competing for the available H₂O₂ or by preventing a fall in activity induced by high levels of H₂O₂. Although effects of estrogens on neutrophil function have been reported (see 3), it is not known whether estrogens are present in adequate amounts or in the proper location to influence microbicidal activity *in situ*.

ACKNOWLEDGMENTS

The excellent technical assistance of Ann Waltersdorff and Joanne Fluvog and the help of Kay Tisdell in the preparation of the manuscript are greatly appreciated.

This study was supported by Public Health Service research grants HD02266 and AI07763 from the National Institutes of Health.

LITERATURE CITED

1. Agner, K. 1958. Crystalline myeloperoxidase. *Acta Chem. Scand.* **12**:89-94.
2. Agner, K. 1963. Studies on myeloperoxidase activity. I. Spectrophotometry of the MPO-H₂O₂ compound. *Acta Chem. Scand.* **17**:332-338.
3. Bodel, P., G. M. Dillard, Jr., S. S. Kaplan, and S. E. Malawista. 1972. Anti-inflammatory effects of estradiol on human blood leukocytes. *J. Lab. Clin. Med.* **80**:373-384.
4. Brökelmann, J. 1969. Peroxidase-associated binding of estradiol by the rat uterus. *J. Histochem. Cytochem.* **17**:394-407.
5. Jacobs, A. A., R. J. Selvaraj, R. R. Strauss, B. B. Paul, G. W. Mitchell, Jr., and A. J. Sbarra. 1973. The role of the phagocyte in host-parasite interactions. XXXIX. Stimulation of bactericidal activity of myeloperoxidase-containing leukocyte fractions by estrogens. *Am. J. Obstet. Gynecol.* **117**:671-678.
6. Jellinck, P. H., and R. Fletcher. 1970. Peroxidase-catalyzed conjugation of [4-¹⁴C]-estradiol with albumin and thiols. *Can. J. Biochem.* **48**:1192-1198.
7. Jellinck, P. H., and L. Irwin. 1962. The oxidation of estrone-16-C¹⁴ by peroxidase. *Can. J. Biochem. Physiol.* **40**:459-469.
8. Jellinck, P. H., and C. R. Lyttle. 1972. Estrogen-induced uterine enzymes in the control of estradiol action. *Adv. Enzyme Regul.* **11**:17-33.
9. Klebanoff, S. J. 1959. An effect of thyroxine on the oxidation of reduced pyridine nucleotides by the peroxidase system. *J. Biol. Chem.* **234**:2480-2485.
10. Klebanoff, S. J. 1960. Reduced pyridine nucleotides as activators of certain reactions catalyzed by peroxidase. *Biochim. Biophys. Acta* **44**:501-509.
11. Klebanoff, S. J. 1961. The sulfite-activated oxidation of reduced pyridine nucleotides by peroxidase. *Biochim. Biophys. Acta* **48**:93-103.
12. Klebanoff, S. J. 1962. An interrelationship between ergothioneine, certain phenolic hormones and peroxidase. *Biochim. Biophys. Acta* **56**:460-469.
13. Klebanoff, S. J. 1965. Inactivation of estrogen by rat uterine preparations. *Endocrinology* **76**:301-311.
14. Klebanoff, S. J. Antimicrobial mechanisms in neutrophilic polymorphonuclear leukocytes. *Semin. Hematol.* **12**:117-142.
15. Klebanoff, S. J. 1977. Estrogen binding by leukocytes during phagocytosis. *J. Exp. Med.* **145**:983-998.
16. Klebanoff, S. J., and R. A. Clark. 1978. The neutrophil: function and clinical disorders, p. 410-434. North-Holland Publishing Co., Amsterdam.
17. Klebanoff, S. J., and S. J. Segal. 1960. The inactivation of estradiol by peroxidase. *J. Biol. Chem.* **235**:52-55.
18. Rosen, H., and S. J. Klebanoff. 1979. Bactericidal activity of a superoxide anion generating system: a model for the polymorphonuclear leukocyte. *J. Exp. Med.* **149**:27-39.
19. Tchernitchin, A., X. Tchernitchin, A. M. Bongiovanni, and R. Chandross. 1974. Effect of hydrogen peroxide on estrogen binding by uterine eosinophils *in vitro*. *J. Steroid Biochem.* **5**:693-695.
20. Williams-Ashman, H. G., M. Cassman, and M. Klavins. 1959. Two enzymic mechanisms for hydrogen transport by phenolic oestrogens. *Nature (London)* **184**:427-429.
21. Worthington Biochemicals Corp. 1972. Worthington enzyme manual, p. 43. Worthington Biochemicals Corp., Freehold, N. J.