

NOTES

Comparison of the Virion Polypeptides of Group B Arboviruses

DANIEL SHAPIRO, DENNIS TRENT, WALTER E. BRANDT, AND PHILIP K. RUSSELL¹

*Department of Virus Diseases, Walter Reed Army Institute of Research, Washington, D.C. 20012 and
Department of Microbiology, University of Texas Medical School at San Antonio,
San Antonio, Texas 79229*

Received for publication 6 March 1972

The virion polypeptides of eight group B arboviruses were compared by polyacrylamide gel coelectrophoresis. All contained three polypeptides, V-1, V-2, and V-3. The mosquito-borne viruses had electrophoretically similar small membrane polypeptides (V-1); the tick-borne viruses had a V-1 of significantly decreased mobility.

Group B arboviruses have been found to contain three virion polypeptides by polyacrylamide gel electrophoresis (1, 3, 4, 6, 7); a fourth radioactive peak is sometimes present. We have designated the three polypeptides of Japanese encephalitis (JE) virus as V-1 (a small membrane polypeptide, molecular weight approximately 8,700), V-2 (the "core" polypeptide, molecular weight approximately 13,500) and V-3 (the large membrane glycoprotein, molecular weight approximately 53,000) (1). However, substantial differences in the molecular weights of the virion polypeptides of other group B arboviruses have been reported. We therefore examined the virion polypeptides of eight group B arboviruses by coelectrophoresis through polyacrylamide gels in order to compare them directly. Since there are major biologic and antigenic differences between subgroups within group B, we included both tick-borne and mosquito-borne viruses to determine whether structural difference may be correlated with the other parameters.

The viruses were studied in two series of coelectrophoretic experiments. Series A viruses were: (i) Suckling mouse brain passage (SMP) 34 of New Guinea "C" strain of dengue-2 (DEN-2); (ii) SMP 27 of M1/311 strain of JE; (iii) SMP 109 of Hubbard strain of St. Louis encephalitis (SLE); (iv) highly mouse-adapted French neurotropic yellow fever; (v) high mouse passage of TP-21 strain of Langat (LAN); and (vi) SMP 6 of Moscow B-4 strain of Russian spring-summer

encephalitis (RSSE) virus. These viruses were grown in LLC-MK₂ cells that were radioactively labeled with either ³H- or ¹⁴C-amino acids and then purified by rate zonal centrifugation through linear sucrose gradients by previously described methods (1, 3). The virions were dissociated with 1% sodium lauryl sulfate and 2-mercaptoethanol, mixed, and subjected to electrophoresis through 8 or 10% polyacrylamide gels either 70 or 180 mm long, as previously described (3). Series B viruses all had high mouse passage histories: (i) KB cell-adapted New Guinea B strain of DEN-2 (1); (ii) Nakayama strain of JE; (iii) Tampa Bay Human-28 strain of SLE; (iv) Ar-248 strain of West Nile (WN)₂; (v) TP-21 strain of LAN; and (vi) Byers strain of Powassan (POW). They were grown in LLC-MK₂ cells, labeled, purified, and subjected to electrophoresis through 10% polyacrylamide gels by previously described methods (6).

The results indicate that all eight viruses had fundamentally similar polypeptide compositions (Fig. 1-3). Nevertheless, two distinct subgroups could be identified, based on the electrophoretic mobility of the small membrane polypeptide V-1. Subgroup I was exemplified by the mosquito-borne viruses JE, DEN-2, SLE, and WN (Fig. 1). Each had a V-1 of similar mobility. The mobility of the V-1 of yellow fever was slightly decreased relative to that of the above four viruses (Fig. 2A), but was greater than that of the tick-borne viruses (Fig. 2B). Subgroup II was exemplified by the members of the tick-borne complex, LAN, POW, and RSSE (Fig. 3); the mobility of V-1 was decreased relative to that of subgroup I. Within this

¹ Present address: Dept. of Microbiology, Dartmouth Medical School, Hanover, N.H. 03755.




FIG. 1. Electrophoresis of subgroup I, mosquito-borne viruses. Series A: top, 40 μ liters of ^{14}C -amino acid-labeled Japanese encephalitis (JE) virions was mixed with 20 μ liters of ^3H -amino acid-labeled dengue-2 (DEN-2) virions, and the mixture was subjected to coelectrophoresis through a 70-mm, 10% acrylamide gel; bottom, 50 μ liters of ^{14}C -JE was mixed with 50 μ liters of ^3H -St. Louis encephalitis virus (SLE) and subjected to coelectrophoresis through a 180-mm, 10% acrylamide gel. Series B: top, 100 μ liters of ^{14}C -SLE was subjected to coelectrophoresis with 100 μ liters of ^3H -DEN-2; center, 100 μ liters of ^{14}C -SLE was subjected to coelectrophoresis with 100 μ liters of ^3H -JE; bottom, 100 μ liters of ^{14}C -SLE was subjected to coelectrophoresis with 75 μ liters of ^3H -West Nile virus (WN).




FIG. 2. Electrophoresis of yellow fever virus (YF). Series A (only): top, 40 μ liters of ^3H -Japanese encephalitis virus (JE) was mixed with 20 μ liters of ^{14}C -YF and subjected to coelectrophoresis through a 70-mm, 8% acrylamide gel; bottom, 20 μ liters of ^{14}C -YF was mixed with 40 μ liters of ^3H -Langat virus (LAN) and subjected to coelectrophoresis on a 70-mm, 10% acrylamide gel.

subgroup, the mobilities of V-3 and V-2 were more similar than within subgroup I.

These data indicate that the mosquito-borne group B arboviruses can be distinguished from the tick-borne group B arboviruses on the basis of the mobility of the small membrane polypeptide V-1; the tick-borne V-1 is either larger (2) or less negatively charged (5) than the mosquito-borne V-1. These observations, although strictly empirical, would raise the possibility that the small membrane polypeptide, V-1, plays a role in determining host range and may be of taxonomic significance.

We thank Timothy Van Duser and Jack McCown for their generous assistance.

Part of this work was supported by Public Health Service research grant AI-09397 from the National Institute of Allergy and Infectious Diseases.

LITERATURE CITED

1. Cardiff, R. D., W. E. Brandt, T. G. McCloud, D. Shapiro, and P. K. Russell. 1971. Immunological and biophysical separation of dengue-2 antigens. *J. Virol.* 7:15-23.
2. Shapiro, A. L., E. Viñuela, and J. V. Maizel. 1967. Molecular weight estimation of polypeptide chains by electrophoresis in SDS-polyacrylamide gels. *Biochem. Biophys. Res. Commun.* 28:815-820.
3. Shapiro, D., W. E. Brandt, R. D. Cardiff, and P. K. Russell. 1971. The proteins of Japanese encephalitis virus. *Virology* 44:108-124.
4. Stollar, V. 1969. Studies on the nature of dengue viruses. IV. The structural proteins of type 2 dengue virus. *Virology* 39:426-438.
5. Strauss, E. G., and P. Kaesberg. 1970. Acrylamide gel electrophoresis of bacteriophage Q β : electrophoresis of the intact virions and of the viral proteins. *Virology* 42:437-452.
6. Trent, D. W., and A. A. Qureshi. 1971. Structural and non-structural proteins of Saint Louis encephalitis virus. *J. Virol.* 7:379-388.
7. Westaway, E. G., and B. M. Reedman. 1969. Proteins of the Group B arbovirus Kunjin. *J. Virol.* 4:688-693.




FIG. 3. Electrophoresis of subgroup II, tick-borne, viruses. Series A: top, 50 μ liters of ^{14}C -Japanese encephalitis virus (JE) was mixed with 50 μ liters of ^3H -Langat virus (LAN) and subjected to coelectrophoresis through a 180-mm, 10% acrylamide gel; bottom, 50 μ liters of ^{14}C -Russian spring-summer encephalitis virus (RSSE) was mixed with 20 μ liters of ^3H -LAN and subjected to coelectrophoresis through a 70-mm, 8% acrylamide gel. Series B: top, 100 μ liters of ^{14}C -St. Louis encephalitis virus (SLE) was subjected to coelectrophoresis with 80 μ liters of ^3H -LAN; bottom, 100 μ liters of ^{14}C -SLE was subjected to coelectrophoresis with 100 μ liters of ^3H -Powassan virus (POW).