

ERRATA

Analysis of Immune Responses of Different Hosts to *Babesia divergens* Isolates from Different Geographic Areas and Capacity of Culture-Derived Exoantigens To Induce Efficient Cross-Protection

E. PRECIGOUT, A. GORENFLOT, A. VALENTIN, G. BISSUEL, B. CARCY, P. BRASSEUR,
Y. MOREAU, AND J. SCHREVEL

Laboratoire de Biologie Cellulaire, URA Centre National de la Recherche Scientifique 290, F-86022 Poitiers Cedex, Laboratoire de Biologie Cellulaire, Faculté de Pharmacie, F-34060 Montpellier, Laboratoire IFFA, Rhône-Mérieux, F-69342 Lyon Cedex 07, Laboratoire de Parasitologie, Hôtel-Dieu, F-76031 Rouen, and Muséum National d'Histoire Naturelle, 61, Rue Buffon, F-75007 Paris, France

Volume 59, no. 8, p. 2803, legend to Fig. 6, lines 1–3: Should read “. . . The gerbils of groups Av, Bv, Cv, and Dv were vaccinated with *B. divergens* Rouen 1987 (in vitro culture supernatant). The gerbils of groups Ac, Bc, Cc, and Dc were injected with placebo. . . .”

Transformation of *Actinobacillus actinomycetemcomitans* by Electroporation, Utilizing Constructed Shuttle Plasmids

PREM K. SREENIVASAN, DONALD J. LeBLANC, LINDA N. LEE, AND PAULA FIVES-TAYLOR

Department of Microbiology and Molecular Genetics, University of Vermont, Burlington, Vermont 05405, and Department of Microbiology, University of Texas Health Science Center at San Antonio, San Antonio, Texas 78284

Volume 59, no. 12, p. 4623, Fig. 1: The *HincII* and *XhoI* sites should be interchanged, and the orientation of the spectinomycin resistance gene should be reversed.