

## Role of the *Escherichia coli* O157:H7 O Side Chain in Adherence and Analysis of an *rfb* Locus

SIMA S. BILGE,<sup>1,2</sup> JAMES C. VARY, JR.,<sup>1,2</sup> SCOTT F. DOWELL,<sup>1,2</sup> AND PHILLIP I. TARR<sup>1,2,3\*</sup>

Departments of Pediatrics<sup>1</sup> and Microbiology,<sup>3</sup> University of Washington School of Medicine, and the Children's Hospital and Medical Center,<sup>2</sup> Seattle, Washington

Received 24 April 1996/Returned for modification 19 June 1996/Accepted 2 August 1996

Shiga-toxicogenic *Escherichia coli* strains belonging to serotype O157 are important human pathogens, but the genetic basis of expression of the O157 antigen and the role played by the lipopolysaccharide O side chain in the adherence of this organism to epithelial cells are not understood. We performed *TnphoA* mutagenesis on *E. coli* O157:H7 strain 86-24 to identify a mutant (strain F12) deficient in O-antigen expression. Nucleotide sequence analysis demonstrated that the transposon inserted within an open reading frame with significant homology to *rfbE* of *Vibrio cholerae* O1 (U. H. Stroehrer, L. E. Karageorgos, R. Morona, and P. A. Manning, Proc. Natl. Acad. Sci. USA 89:2566–2570, 1992), which is postulated to encode perosamine synthetase. This open reading frame was designated *rfbE*<sub>EcO157:H7</sub>. The guanine-plus-cytosine fraction (0.35) suggests that *rfbE*<sub>EcO157:H7</sub> may have originated in a species other than *E. coli*. *rfbE*<sub>EcO157:H7</sub> is conserved in nontoxicogenic *E. coli* O157 strains expressing a variety of other flagellar antigens but is not found in *E. coli* O55:H7 strains, which are more closely related to *E. coli* O157:H7. Strain F12 was significantly more adherent to HeLa cells in a quantitative adherence assay than was its *E. coli* O157:H7 parent, but they did not differ in other phenotypes. Restoration of the expression of the O side chain by complementation of the *TnphoA* mutation in strain F12 by a plasmid expressing intact *rfbE*<sub>EcO157:H7</sub> reduced the adherence of the hyperadherent strain F12. We conclude that *rfbE*<sub>EcO157:H7</sub> is necessary for the expression of the O157 antigen, that acquisition of *E. coli rfb* genes occurred independently in *E. coli* O157:H7 and unrelated O157 strains, and that the O side chain of *E. coli* O157:H7 lipopolysaccharide interferes with the adherence of *E. coli* O157:H7 to epithelial cells.

Shiga-toxicogenic *Escherichia coli* strains expressing lipopolysaccharide (LPS) O-antigen 157 are important human enteric pathogens. Many *E. coli* strains produce Shiga toxins, but Shiga-toxicogenic *E. coli* strains possessing this particular O antigen are the pathogens most frequently isolated from humans in North America and are the predominant cause of hemolytic uremic syndrome (34).

Sorbitol-nonfermenting *E. coli* O157:H7 and sorbitol-fermenting *E. coli* O157:NM (15) are clonally related on the basis of having similar or identical isoenzyme forms detected by multilocus enzyme electrophoresis (42, 43). *E. coli* O55:H7 is the serotype most closely related to *E. coli* O157:H7 and *E. coli* O157:NM by this analysis; 6-phosphogluconate dehydrogenase (EC 1.1.1.44), encoded by the *gnd* locus which is adjacent to the *rfb* region in *E. coli* (2), is the only differentiating electromorph among 20 enzymes studied. In contrast, *E. coli* O157 strains expressing flagellar antigens other than H7 are distantly related to *E. coli* O157:H7 (42, 43); *E. coli* O157 with non-H7 flagellar antigens display multiple polymorphisms for each of these 20 different enzymes and do not express Stx1, Stx2, or intimin.

We performed *TnphoA* mutagenesis of *E. coli* O157:H7 to identify a mutant deficient in expression of the O157 antigen for the purposes of (i) identifying *rfb* loci of this serotype, (ii) determining if these sequences are conserved in *E. coli* O157 expressing different flagellar antigens and in *E. coli* O55:H7, and (iii) assessing the role played by the O side chain of *E. coli* O157:H7 in the adherence of this organism to epithelial cells in vitro.

(These data were presented in part at the 95th General Meeting of the American Society for Microbiology [4a].)

### MATERIALS AND METHODS

**Bacterial strains used.** *E. coli* O157:H7 strain 86-24 was used as the prototype for these studies (14, 35). A spontaneously occurring nalidixic acid-resistant mutant of this organism (designated *E. coli* O157:H7 86-24<sup>nalR</sup>) was isolated from Luria-Bertani (LB) (29) agar plate containing nalidixic acid (20 mg/liter), onto which the centrifuged bacterial contents of 1 ml of overnight LB broth culture of *E. coli* O157:H7 86-24 were spread.

*E. coli* O157 strains were studied to determine the conservation of the *E. coli* O157:H7 *rfb* locus. Eighteen *E. coli* O157:H7 strains were analyzed, including 17 strains from a previous study (35) and the outbreak strain from the 1993 multi-state *E. coli* O157:H7 epidemic (3). Five strains of *E. coli* O55:H7 and four strains of *E. coli* O157:H43 were obtained from Thomas Whittam. Seven strains of *E. coli* O157 expressing H antigens 3 ( $n = 1$ ), 12 ( $n = 1$ ), 16 ( $n = 3$ ), 38 ( $n = 1$ ), and 45 ( $n = 1$ ) were kindly provided by Peter Feng. A sorbitol-fermenting, Stx2-producing *E. coli* O157:NM strain from a German patient with hemolytic uremic syndrome was contributed by Lothar Beutin. Except for this last strain, none of the non-H7 *E. coli* O157 strains studied produced Stx1 or Stx2. DNAs from non-O157 Shiga-toxicogenic *E. coli* strains isolated from humans (7) were also probed.

*E. coli* SM10 ( $\lambda$ -pir) transformed with pRT733 (36), a suicide vector into which *TnphoA* has been cloned, was used for conjugative introduction of *TnphoA* into *E. coli* O157:H7 86-24<sup>nalR</sup>, *E. coli* B171 (26), a nontoxicogenic enteropathogenic strain of serotype O111:NM, *E. coli* O26:H11, which produces Stx1 but not Stx2 (35), *E. coli* O55:H7, and *E. coli* HB101 (29) were used to study the expression of a cloned *E. coli* O157:H7 *rfb* locus in various *E. coli* strains.

***TnphoA* mutagenesis.** *E. coli* SM10 ( $\lambda$ -pir) containing pRT733 was mated with *E. coli* O157:H7 86-24<sup>nalR</sup>. Sixty separate matings produced kanamycin- and nalidixic acid-resistant transconjugants, of which approximately 0.5% expressed alkaline phosphatase (PhoA), as indicated by the presence of blue colonies when plated on agar containing phosphate (to suppress endogenous PhoA activity) and 5-bromo-4-chloro-3-indolylphosphate (X-P), a chromogenic substrate of PhoA. A total of 3,000 transconjugants (approximately 50 from each mating) were frozen in LB broth–15% glycerol at  $-70^{\circ}\text{C}$  until analysis. To determine if the O157 antigen was retained by these mutants, transconjugants were thawed, grown on LB agar containing nalidixic acid (20 mg/liter) and kanamycin (37.5 mg/liter), and tested individually for expression of this antigen by using the Oxoid (Unipath) latex particle agglutination test.

**DNA cloning and sequencing strategy.** The region of the *TnphoA* insertion in

\* Corresponding author. Mailing address: Division of Gastroenterology, Children's Hospital and Medical Center, 4800 Sand Point Way NE, Seattle, WA 98105. Phone: (206) 526-2521. Fax: (206) 528-2721. Electronic mail address: tarr@u.washington.edu.

the mutant of interest was first mapped by hybridization to identify restriction sites in the DNA flanking the transposon. Size-selected genomic DNA fragments from *E. coli* O157:H7 strain F12 were ligated into the corresponding sites in plasmid Bluescript SK<sup>+</sup> (pSK<sup>+</sup>) (Stratagene) to clone the DNA flanking the *TnphoA* insertion. A 6.6-kb fragment of DNA extending from the *Bam*HI site of *TnphoA* to the *Eco*RV site in DNA adjacent to the IS50L flank of *TnphoA* was ligated into corresponding sites in pSK<sup>+</sup>. Similarly, a 7.3-kb fragment of DNA extending from *Cl*aI sites in *TnphoA* and in DNA adjacent to the IS50R region was ligated into the *Cl*aI site of pSK<sup>+</sup>. These fragments were selected by plating transformed *E. coli* HB101 on LB agar containing kanamycin.

Initial sequence from the cloned DNA adjacent to the IS50L region of *TnphoA* was obtained by using a primer (5'-GTAAAACGACGGCCAGT3') complementary to vector sequences adjacent to the cloning site, single-stranded DNA template created with the VCS-M13 helper phage (Stratagene), and the dideoxy-chain termination procedure (30) with Sequenase (U.S. Biochemicals). All oligonucleotides were purchased from GenSet. The initial sequence from the DNA adjacent to the right flank of *TnphoA* was obtained by using Sequenase, double-strand sequencing techniques suggested by the manufacturer, and primer oligonucleotides complementary to IS50R (5'-CGGCCGCACGATGAAGAGC3') and M13 reverse primer (5'-GGAAACAGCTATGACCATG3') sequences of pSK<sup>+</sup>.

After sequence analysis of the cloned DNA originating from *E. coli* O157:H7, we performed PCR with 5'-GGGGATCCTAATCTTCTGGCATGATTGATTGGC3' and 5'-GGGAATTCCTTTACAATTCACCGCCCGCCACTCG3' (incorporating *Bam*HI and *Eco*RI sites, respectively, for cloning purposes) as primers, which were derived from the cloned DNA furthest from *TnphoA* in the regions adjacent to the left and right flanks, respectively. This amplified PCR product was cloned into the respective sites on pSK<sup>+</sup> and sequenced by double-strand sequencing techniques with the *Taq* DyeDeoxy cycle-sequencing kit (Applied Biosystems) under the conditions recommended by the manufacturer, with appropriate intervening primers and an Applied Biosystems model 373A DNA sequencing unit (Molecular Pharmacology Unit, University of Washington School of Medicine). Resulting sequences were confirmed by double-strand sequencing of the cloned DNA flanking the *TnphoA* insertion in strain F12 and compared with databases reached through the National Center for Biotechnology Information GenInfo BLAST network server (12).

The gene into which *TnphoA* inserted was amplified from parental *E. coli* O157:H7 by PCR with oligonucleotides 5'-GGGGATCCTAATCTTCTGGCATGATTGATTGGC3' and 5'-GGGAATTCCTTTACAATTCACCGCCCGCCACTCG3' (incorporating *Bam*HI and *Eco*RI sites, respectively, for ligation purposes) as primers and cloned into pSK<sup>+</sup>. The resulting construct was designated pF12.

**LPS analysis.** Bacterial LPS from equivalent numbers of organisms was extracted with hot phenol (16), separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE; 12% polyacrylamide), and visualized by silver staining (37). Electrophoretically separated LPS was transferred to Immobilon-P (Millipore), which was then preadsorbed in phosphate-buffered saline (PBS) containing 3% bovine serum albumin (Sigma), and probed with polyclonal rabbit antibody to the O157 antigen (a gift of Joy Wells) diluted 1:100 in PBS-3% BSA. Immunoreactive LPS was identified by incubation of the membrane with goat anti-rabbit antibody coupled to horseradish peroxidase (Sigma) and the enhanced chemiluminescence substrate (Amersham) followed by autoradiography.

**Outer membrane protein analysis.** Bacterial outer membrane proteins were isolated by the method of Achtman et al. (1) after overnight growth under the same conditions as used for bacteria in the adherence assays. A 20- $\mu$ g sample of protein from each strain was analyzed in separate lanes of SDS-PAGE (12% polyacrylamide) gels. The resulting separated proteins were visualized by Coomassie blue staining.

**Southern hybridization.** DNA was extracted from mutant and wild-type *E. coli* (28), digested, electrophoretically separated in 0.7% agarose gels in 0.5 $\times$  Tris-borate-EDTA, transferred to a Nytran membrane (Schleicher & Schuell), and probed. Membranes were prehybridized in 5 $\times$  SSC (1 $\times$  SSC is 0.015 M sodium citrate and 0.15 M NaCl)-1 mM EDTA-0.1% 5 $\times$  Denhardt's solution-100 mg of sonicated and heat denatured calf thymus DNA per ml before the addition of probe. All probes were labelled with the Megaprime DNA-labelling system (Amersham) and [ $\alpha$ -<sup>32</sup>P]dCTP (New England Nuclear Research Products). The probes used were a 0.7-kb *Sma*I-*Pst*I fragment from the Tn5 central region of *TnphoA* and a 1.3-kb fragment corresponding to the complete gene into which *TnphoA* inserted in the mutant of interest (produced as a PCR-derived clone [described above]).

**Phenotypic analysis of the mutant of interest.** The mutant which no longer expressed the O157 antigen was tested for phenotypes associated with the parental *E. coli* O157:H7 strain, including the production of a cytotoxin for Vero cells (11), expression of the H7 antigen (kindly performed by Jay Lewis, Washington State Department of Health), and inability to ferment sorbitol when plated on sorbitol-MacConkey agar (Prepared Media Laboratory) (41) or to produce  $\beta$ -glucuronidase from 4-methylumbelliferyl- $\beta$ -D-glucuronide (Sigma), a fluorogenic substrate of this enzyme (27).

**Adherence assay.** HeLa cells grown to confluence at 37°C in 5% CO<sub>2</sub> in minimal essential medium containing 10% heat-inactivated fetal calf serum, L-glutamine (2 mM), penicillin (100 IU/ml), and streptomycin (100  $\mu$ g/ml) were

trypsinized, diluted, added to glass chamber slides, and reincubated. The adherence assay was performed 2 days later, when the cells were approximately 80% confluent.

To assay bacterial adherence, the cells in each chamber were washed with sterile PBS and covered with 0.6 ml of incubation medium (minimal essential medium, 5% fetal calf serum, 2 mM L-glutamine, nonessential amino acids, 0.5% D-mannose). A 20- $\mu$ l portion of bacteria which had been grown overnight with agitation in LB broth at 37°C was added to the HeLa cells. Bacterial strains which had not been transformed with pSK<sup>+</sup> were incubated at 37°C in 5% CO<sub>2</sub> for 2 h. The chambers were then washed three times with sterile PBS, covered with 0.6 ml of incubation medium, and incubated again for 2 h (toxic effects from strain F12 were noted with longer incubation periods). Bacterial strains which had been transformed with pSK<sup>+</sup> were incubated with cells in the presence of ampicillin (100 mg/liter), and the incubation periods were 3 and 3 h (strain F12 transformed with pSK<sup>+</sup> was not as cytotoxic as strain F12, permitting the longer incubation periods). Nonadherent bacteria were removed by 10 washes of the chambers with PBS from a plastic wash bottle at the end of the assays.

Washed cells and adherent bacteria were fixed by the addition of 100% methanol to each chamber for 5 min, stained with Giemsa stain for 60 min, mounted, and examined for adherence. Positive and negative controls were locally adherent enteropathogenic *E. coli* B171, and nonadherent laboratory strain *E. coli* ORN172, an *E. coli* K-12 derivative from which cryptic type 1 pilus genes had been deleted (44), respectively. Five fields were examined in each chamber by an observer unaware of the identity of the organisms which had been added to each chamber, and the HeLa cells and adherent clusters (five or more bacilli per cluster) were enumerated. The values from at least eight experiments were used to determine the adherence ratio, which is expressed as the mean number of clusters ( $\pm$  standard deviation) per cell. The significance of differences between means was determined by the two-tailed *t* test.

## RESULTS

**Identification of a *TnphoA* mutant of *E. coli* O157:H7 deficient in the expression of the O157 LPS antigen.** A total of 3,000 *TnphoA* mutants of *E. coli* O157:H7 were tested individually for the loss of the O157 antigen. A single mutant, designated F12, which expressed PhoA at a low level (manifested by a faint blue color when the mutant was plated on agar containing X-P), failed to react in the O157 latex particle agglutination test. Southern blot analysis demonstrated that strain F12 had sustained a single *TnphoA* insertion in a 20-kb chromosomal *Mlu*I fragment (data not shown). Strain F12 neither fermented sorbitol after overnight incubation on sorbitol MacConkey agar nor produced  $\beta$ -glucuronidase but was motile, expressed the H7 antigen, and elaborated a cytotoxin for Vero cells; these phenotypes are identical to those of *E. coli* O157:H7 strain 86-24 and *E. coli* O157:H7 strain 86-24<sup>nalR</sup>, from which F12 was derived.

**Cloning and sequencing of the DNA adjacent to the *TnphoA* insertion in strain F12.** (1.7 and 0.6 kb) of DNA flanking the IS50L and IS50R regions, respectively, of the *TnphoA* insertion in strain F12 were sequenced. *TnphoA* had inserted into an open reading frame (ORF) beginning with an ATG codon 6 bp 3' to a Shine-Dalgarno sequence, extending to a stop codon (TAG) 1,095 nucleotides downstream (Fig. 1). This ORF encodes a putative polypeptide of 364 amino acids with a predicted molecular mass of 41,552 Da. Analysis of the deduced amino acid sequence (39) failed to demonstrate a probable cleavage site for a signal peptide. The *TnphoA* insertion site in strain F12 is noted by an arrow in Fig. 1.

The protein encoded by the ORF into which *TnphoA* inserted has extensive homology to the probable perosamine synthetase encoded by *rfbE* of *Vibrio cholerae* O1 (33) (Fig. 2). Like *V. cholerae*, the LPS of *E. coli* O157:H7 contains perosamine (22). Because perosamine, an unusual component of bacterial LPS, is found in the side chains of both organisms and because of the extensive homology between *RfbE* of *V. cholerae* O1 and the protein encoded by the ORF into which *TnphoA* inserted in strain F12, we have designated this *E. coli* O157:H7 ORF *rfbE*<sub>Eco157:H7</sub>. Of the 364 amino acids in *RfbE*<sub>Eco157:H7</sub>, 197 (54.1%) can be exactly aligned with the probable perosamine synthetase of *V. cholerae* O1: an addi-

```

GGGTTAACTGTTATGTTGACTGCTTCATTTTTATATATATTGTAATAACTTTATATGGTATAAATGTAGTTTAAAAACATATCGATAGACAGTTAA
ATATAAGAGGATGAAAATGAAATATATACCAGTTTACCAACCGTCATTGACAGGAAAAGAAAAGAATATGTAATGAATGCTGGACTCAACGTTGGATT
      M K Y I P V Y Q P S L T G K E K E Y V N E C L D S T W I
TCATCAAAGGAACTATATTCAGAAGTTGAAAATAAATTTGCGGAACAAAACCATGTGCAATATGCAACTACTGTAAGTAAATGGAACGGTTGCTCTTC
S S K G N Y I Q K F E N K F A E Q N H V Q Y A T T V S N G T V A L H
ATTTAGCTTTGTTAGCGTTAGGTATATCGGAAGGAGATGAAGTTATTGTTCCAACACTGACATATATAGCATCAGTTAATGCTATAAAAACACAGGAGC
  L A L L A L G I S E G D E V I V P T L T Y I A S V N A I K Y T G A
CACCCCCATTTTCGTTGATTGAGATAATGAACTTGGCAAATGTCTGTAGTGACATAGAACAAAAAATCACTAATAAACTAAAGCTATTATGTGTGTC
  T P I F V D S D N E T W Q M S V S D I E Q K I T N K T K A I M C V
CATTTATACGGACATCCATGTGATATGGAACAAATGTAGAAGTGGCCAAAAGTAGAAATTTGTTGTAATTTGAAGATTGCGCTGAAGCCTTTGGTTCTA
H L Y G H P C D M E Q I V E L A K S R N L F V I E D C A E A F G S K
AATATAAAGGTAATATGTGGAAACATTTGGAGATATTTCTACTTTTAGCTTTTTTGGAAATAAACTATTACTACAGGTGAAGGTGGAATGGTTGTGAC
  Y K G K Y V G T F G D I S T F S F F G N K T I T T G E G G M V V T
GAATGACAAAACACTTTATGACCGTTGTTTACATTTTAAAGGCCAAGGATTAGCTGTACATAGGCAATATTGGCATGACGTTATAGGCTACAATTATAGG
  N D K T L Y D R C L H F K G Q G L A V H R Q Y W H D V I G Y N Y R
ATGACAAATATCTGCGTCTATAGGATTAGCCAGTTAGAACAAGCTGATGATTTATATCACGAAAACGTTGAATTTGCTGATATTTATAAAAAAATA
M T N I C A A I G L A Q L E Q A D D F I S R K R E I A D I Y K K N I
TCAACAGTCTTGTAACAAGTCCACAAGGAAAGTAAAGATGTTTTTACACTTATTGGATGGTCTCAATTCTAACTAGGACCCGAGAGGAAAGAGAGGAATT
  N S L V Q V H K E S K D V F H T Y W M V S I L T R T A E E R E E L
AAGGAATCACCTTGCGATAACTCATCGAAACAAGGCCAGTTTTTACCCTGTCCACACGATGCCAATGTACTCGGAAAAATATCAAAGCACCTATA
  R N H L A D K L I E T R P V F Y P V H T M P M Y S E K Y Q K H P I
GCTGAGGATCTTGTTGGCGTGAATTAATTACCTAGTTTCCCAGCCTATCGAATGAGCAAGTTATTTATATTGTGAATCTATTAACGAATTTTATA
A E D L G W R G I N L P S F P S L S N E Q V I Y I C E S I N E F Y S
GTGATAAATAGCCTAAAATATTGTAAGGTCATTCATGAAAATTGCGTTGAATTCAGATGGATTTTACGAGTGGGGCGGTGGAATTG
  D K

```

FIG. 1. Complete nucleotide sequence of inserted DNA in pF12, with deduced amino acid sequences. The Shine-Dalgarno sequence is underlined. The arrow indicates the site of the *TnphoA* insertion in strain F12.

tional 135 (37%) of the amino acids of RfbE<sub>EcO157:H7</sub> can be conservatively matched to residues in this *V. cholerae* O1 enzyme. There is 62% homology at the nucleotide level between *rfbE*<sub>EcO157:H7</sub> and *rfbE* of *V. cholerae* O1 (results not shown).

**Complementation of the *TnphoA* insertion in *rfbE*<sub>EcO157:H7</sub> of strain F12 with *rfbE*<sub>EcO157:H7</sub>.** A DNA fragment encoding the *E. coli* O157:H7 *rfbE* locus was amplified via PCR with primers derived from sequences flanking this gene, and inserted into pSK<sup>+</sup>. The resulting construct, pF12, when introduced into strain F12, complemented the *TnphoA* mutation, as evidenced by restored reactivity in the latex particle agglutination test.

**LPS analysis of *E. coli* O157:H7 strain 86-24 and derivatives.** The LPS profiles of *E. coli* O157:H7 and derivatives are presented in Fig. 3A. *TnphoA* mutant strain F12, which does not react in the latex particle agglutination test, does not express detectable O side chain on immunoblot analysis, although the core LPS is still expressed in strain F12. LPS from only the strains reactive in the O157 latex particle agglutination test (*E. coli* O157:H7 strain 86-24, *E. coli* O157:H7 strain 86-24<sup>nalR</sup>, and strain F12 transformed with pF12) display the high-molecular-weight ladder representing O-polysaccharide side chain. When immobilized LPS was probed with antibodies to the O157 antigen, LPS from strain F12 demonstrated no reactive antigens in the core region of the gels (Fig. 3B).

LPS purified from *E. coli* O26:H11, *E. coli* O55:H7, and *E. coli* O111:NM (strain B171) transformed with pF12 failed to express O side chains recognized by antibody to the O157

```

RfbEO157 1 MKYIPVYQPSLTGKEEYVNECLDSTWISSKGNVYQKFNKFAEQNHVQY 50
RfbEEc 1 ..MIPVYEPSLDGNRKYLNDCIDSGWVSSRGKYIDRFETEFAEFLKVKH 48
51 ATTVSNGTVALHALLALGISEGDEVIVPTLYTIASVNAIKYTGATPIFV 100
49 ATTVSNGTVALHLAMSALGTTGGDEVIVPTFYVAVSNTIVQCGALPVFA 98
101 DSDNETWQMSVSDIEQKITNKTAIMCVHLYGHPCDMEQIVELAKSRNLF 150
99 EIEGESLQVSVEDVKKRINKKTKAVMAVHYGQACDIQSLRDLDCHEGLY 148
151 VIEDCAEAFGSKYKGYVGTFGDISTFSPFGNKTITTTGEGGMVVNDKTL 200
149 LIEDCAEAIQTAVNGKVGTFGDEVSTFSPFGNKTITTSGEGGMVVSNDII 198
201 YDRCLHFQGGGLAVHRQYWHDIVGYNRMNTNCAAIGLAQLEQADDFISR 250
199 IDKCLRLKNQGVVAGKRYWHDLVAYNYRMTNLCAAIGVAQLERVDKIKA 248
251 KREIADYKKNINSLVQVHVESKDVFTYWMVSIILTRTAE..REELRN 297
249 KRDI AEIYRSELAGLPMQVHVESNGTFHSHYWLTSIILDQEFVHRDGLMT 298
298 HLADKLIETRPVYFPVHTMPMYSEKYQ..HPAIEDLWGRGINLPSFSP 345
299 FLENNDESRPFYPAHTLPMYEHAEKTAFFPLSNYSYSHRGINLPSWGPL 348
346 SNEQVIYICESINEFYSDK 364
349 CDDQVKEICNCKNYFNCCI 367

```

FIG. 2. Homology between RfbE<sub>EcO157:H7</sub> and the probable perosamine synthetase of *V. cholerae* O1. Amino acids are aligned by using the Genetics Computer Group sequence analysis software package (10). Exact matches between aligned amino acids are denoted by vertical lines. Double and single dots represent amino acid pairs with a comparison value of  $\geq 0.50$  and 0.10 to 0.50, respectively (13). Gaps introduced in the sequence to optimize alignment are indicated by dots between adjacent residues.




FIG. 3. LPS analysis of *E. coli* O157:H7 and derivatives. LPS from *E. coli* O157:H7 strain 86-24<sup>naIR</sup> (lane A), *TnphoA* mutant F12 (lane B), strain F12 transformed with pSK+ (lane C), strain F12 transformed with pF12 (lane D), *E. coli* HB101 transformed with pSK+ (lane E), and *E. coli* O157:H7 strain 86-24 (lane F) were separated electrophoretically, visualized with silver stain (A) and transferred to Immobilon-P and probed with antibody to the O157 antigen (B).

antigen when analyzed by immunoblotting of electrophoresed, immobilized antigen (data not shown).

**Conservation of the *E. coli* O157:H7 *rfbE*<sub>EcO157:H7</sub> locus.** We used as a probe a fragment corresponding to *rfbE*<sub>EcO157:H7</sub> to determine the conservation of this gene in *E. coli* O157:H7 and other *E. coli* strains. Homology was detected on a 6.5-kb *EcoRI* fragment of chromosomal DNA in each of 18 *E. coli* O157:H7 strains tested (data not shown); in each of 11 *E. coli* strains of serogroup O157 expressing H antigens 3, 12, 16, 38, 43, and 45; and in a nonmotile, sorbitol-fermenting, Stx2-producing *E. coli* O157 strain (Fig. 4). One of the *E. coli* O157:H16 strains contained a region of homology to *rfbE*<sub>EcO157:H7</sub> in a slightly smaller *EcoRI* fragment (Fig. 4, lane L). Homology to *rfbE*<sub>EcO157:H7</sub> was not detected in five strains of *E. coli* O55:H7 (Fig. 4) or in five strains of Stx-producing *E. coli*, isolated from Seattle children, which expressed a variety of O and H antigens, including *E. coli* O26 (data not shown).

**Adherence properties of *E. coli* O157:H7 and O-antigen-deficient strain F12.** Table 1 demonstrates the quantitative adherence characteristics of *E. coli* O157:H7 compared with those of its O-antigen-deficient mutant. *E. coli* O157:H7 strain 86-24<sup>naIR</sup> adheres sparsely to epithelial cells; strain F12 adheres approximately sevenfold better (Fig. 5). The hyperadherence of strain F12 was significantly diminished by the introduction of pF12, which partially restores the O157 side chain antigen (Table 1; Fig. 5). *E. coli* ORN172 was completely nonadherent in these assays.

**OMP analysis of *E. coli* O157:H7 86-24<sup>naIR</sup> and F12.** To determine if the hyperadherence of strain F12 could be attributed to the increased expression of a surface protein, the OMP profiles of strain F12 and *E. coli* O157:H7 86-24<sup>naIR</sup>, with and




FIG. 4. Conservation of *rfbE*<sub>EcO157:H7</sub> in strains of *E. coli* O157 but not in strains of *E. coli* O55:H7. Shown are total DNAs from *E. coli* O157:H7 strain 86-24 (lane A), *E. coli* HB101 (lane B), *E. coli* O55:H7 strains (lanes C to G), nontoxicogenic *E. coli* O157 strains expressing H antigens 43 (lanes H to K), 16 (lanes L to N), 45 (lane O), 3 (lane P), 38 (lane Q), and 12 (lane R), and toxicogenic, sorbitol-fermenting, *E. coli* O157:NM (lane S), digested with *EcoRI*, electrophoresed, transferred to Nytran, and probed with *rfbE*<sub>EcO157:H7</sub>.

without transformation with pSK+, were compared. There were no prominent bands in the F12 strains compared with the parent strains (Fig. 6).

## DISCUSSION

*rfbE*<sub>EcO157:H7</sub> is necessary for expression of the O157 antigen of *E. coli* O157:H7. Complementation of the *TnphoA* mutation in strain F12 with *rfbE*<sub>EcO157:H7</sub> restored O157 antigenicity. Therefore, loss of expression of this antigen was not caused by a polar effect on a gene downstream of *rfbE*<sub>EcO157:H7</sub>. RfbE<sub>EcO157:H7</sub> is similar to RfbE of *V. cholerae* O1, a probable perosamine synthetase (33). The *E. coli* O157:H7 O polysaccharide consists of [→3-α-D-GalNAcP-(1→2)α-D-PerNAcP-(1→3)-α-L-Fucp-(1→4)-β-D-Glcp-(1→)]<sub>n</sub> repeats (22). Although the function of RfbE<sub>EcO157:H7</sub> is unknown, its homology to RfbE of *V. cholerae* and the presence of perosamine in the LPS of both organisms suggest that RfbE<sub>EcO157:H7</sub> is responsible for perosamine synthesis. Humans immunized with killed *V. cholerae* O1 develop antibodies which detect the *E. coli* O157 antigen (8). Perhaps the perosamine moiety contributes a cross-reactive epitope between the LPS of these organisms.

*E. coli* O157:H7 and sorbitol-fermenting, toxicogenic *E. coli* O157:NM are more closely related to *E. coli* O55:H7 than to non-H7 *E. coli* O157 strains (43), differing only in the 6-phosphogluconate dehydrogenase electromorph (43). 6-Phosphogluconate dehydrogenase polymorphism results from intra-

TABLE 1. Adherence of *E. coli* O157:H7 86-24<sup>naIR</sup> and derivatives to HeLa cells<sup>a</sup>

Strain	No. of clusters/cell (mean ± s.d.)	P <sup>b</sup>
O157:H7 86-24 <sup>naIR</sup>	0.04 ± 0.05	
F12	0.27 ± 0.30	<0.05
F12(pSK+)	0.17 ± 0.13	
F12(pF12)	0.06 ± 0.04	<0.05

<sup>a</sup> Adherence is expressed as the number of clusters of bacteria (containing five or more organisms) adherent to HeLa cells. Five fields were examined for each chamber, and the clusters and cells visualized were enumerated. Values represent examination of at least eight chambers.

<sup>b</sup> P values represent comparison of the adherence of strain F12 with that of its parent, *E. coli* O157:H7 86-24<sup>naIR</sup>, and of strain F12 transformed with pSK+ with that of strain F12 transformed with pF12.




FIG. 5. Adherence of *E. coli* O157:H7 strains 86-24<sup>nalR</sup> (A) and F12 (B) to HeLa cells.

genic and extragenic recombination in *gnd* (5, 20, 21). Whittam proposed that recombination involving the *rfb* clusters and *gnd* loci differentiated the O side chains of *E. coli* O157:H7 from *E. coli* O55:H7 (42). The absence of *rfbE*<sub>E<sub>co</sub>O157:H7</sub> from *E. coli* O55:H7 is consistent with this hypothesis. The guanine-plus-cytosine fraction of *rfbE*<sub>E<sub>co</sub>O157:H7</sub> (0.35) also suggests that this gene might not have originated in *E. coli*. Low G+C fractions have been identified in *rfb* loci of *E. coli* O7 (18), *E. coli* O101 (9), *E. coli* K-12 (31), and *Salmonella enterica* strains (24).

*stx1* and *stx2* are contained on bacteriophages (32), which were presumably acquired by *E. coli* O157:H7 via transduction from an unidentified donor species. Intimin, which mediates the ability of *E. coli* O157:H7 to induce actin aggregation, is encoded by *eaeA*<sub>E<sub>co</sub>O157:H7</sub> (15). *eaeA*<sub>E<sub>co</sub>O157:H7</sub> is in a large segment of DNA, designated *lee* (locus of enterocyte effacement) (19). *lee* (G+C content, 0.39) is integrated at the same site in the *E. coli* O157:H7 chromosome as are large fragments




FIG. 6. OMP profiles of *E. coli* O157:H7 strains 86-24<sup>nalR</sup> (lanes A and C) and F12 (lanes B and D), without (lanes A and B) and with (lanes C and D) pSK+.

which include genes encoding intimin and uropathogenic virulence factors in enteropathogenic (19) and uropathogenic (6) *E. coli* strains, respectively. Presumably, these virulence-associated segments, like *rfbE*<sub>EcO157:H7</sub>, were acquired laterally from another organism.

Preliminary analysis of DNA adjacent to *rfbE*<sub>EcO157:H7</sub> demonstrates an upstream ORF with significant homology to *orf7.4* of *Yersinia pseudotuberculosis* (reference 17 and unpublished results). *orf7.4* is postulated to encode a transmembrane protein which translocates O side chains to the bacterial periphery, and it has recently been designated *rfbX* (25). The sequenced portion of this gene in *E. coli* O157:H7, which we have tentatively designated *rfbX*<sub>EcO157:H7</sub>, also has a low G+C content (0.31); *rfbX*<sub>EcO157:H7</sub>, like *rfbE*<sub>EcO157:H7</sub>, occurs in *E. coli* strains expressing the O157 antigen but not in *E. coli* O55:H7 strains (data not shown).

Our data also suggest that the *E. coli* O157:H7 O side chain diminishes the adherence of *E. coli* O157:H7 to cultured epithelial cells. The outer membrane protein profile of strain F12 does not contain bands that are absent from strain *E. coli* O157:H7 strain 86-24<sup>nalR</sup>, so it is unlikely that a protein adhesin was hyperexpressed because of *TnphoA* insertion in *rfbE*<sub>EcO157:H7</sub>. Ail-dependent invasion of Chinese hamster ovary cells by *Y. enterocolitica* is enhanced in strains deficient in the expression of O:3 side chains. Monoclonal antibodies detect Ail in larger quantities on the surface of the hyperinvasive O:3<sup>-</sup> mutants, implying that LPS side chains of *Y. enterocolitica* mask Ail (23) and possibly other virulence determinants. *Shigella flexneri* strains expressing recombinant *inv* penetrate eukaryotic cells only after the *S. flexneri* strains are mutated so that O side chains are not expressed (40). Expression of *E. coli* and *S. typhimurium* O antigens reduces the ability of antibodies to bind to porin surface epitopes (4). Additionally, expression of the O antigen of LPS reduces the ability of monoclonal antibodies to bind to porin surface epitopes in *E. coli* and *S. typhimurium* (4) and to PhoE of various members of the family *Enterobacteriaceae* (38). These observations suggest that the O side chains of bacterial LPS physically hinder contact between outer membrane virulence factors and eukaryotic cells, possibly modulating the effects of virulence traits by bacterial pathogens. The hyperadherence of the O157<sup>-</sup> mutant in vitro raises the possibility that variable expression of LPS in vivo influences adherence of *E. coli* O157:H7 to enterocytes.

In summary, the homology of *rfbE*<sub>EcO157:H7</sub> to corresponding genes in *E. coli* O157 of differing H antigens, but not to those in the more closely related *E. coli* O55:H7, and its low G+C content support the concept that O-antigen diversity results from the horizontal acquisition of genetic material, possibly from an interspecific source, rather than from point mutations. Furthermore, our data suggest that the O157 side chain inhibits the adherence of *E. coli* O157:H7 to epithelial cells. The in vivo relevance of this observation warrants further study.

#### ACKNOWLEDGMENTS

We thank Sheila Hull, Richard Hull, Peter Reeves, and Thomas Whittam for helpful discussions; Steve Moseley for advice and encouragement; Dong Nguyen and May Yau for technical expertise; and Christine Merrikin for secretarial assistance.

This research was supported by the Crohn's Colitis Foundation of America, an American Gastroenterological Association/Blackwell Scientific Publications Research Award, and USDA grant 94-03953.

#### ADDENDUM IN PROOF

Since submission of the manuscript, another communication has reported that an O-side-chain-deficient mutant of *E. coli*

O157:H7 is hyperadherent (F. Cockerill, G. Beebakhee, R. Soni, and P. Sherman, *Infect. Immun.* **64**:3196–3200, 1996).

#### REFERENCES

- Achtman, M., A. Mercer, B. Kusecek, A. Pohl, M. Heuzenroeder, W. Aaronson, A. Sutton, and R. P. Silver. 1983. Six widespread bacterial clones among *Escherichia coli* K1 isolates. *Infect. Immun.* **39**:315–335.
- Bachmann, B. J. Linkage map of *Escherichia coli* K-12, edition 8. 1990. *Microbiol. Rev.* **54**:130–197.
- Bell, B. P., M. Goldoft, P. M. Griffin, M. A. Davis, D. C. Gordon, P. I. Tarr, C. A. Bartleson, J. H. Lewis, T. J. Barrett, J. G. Wells, et al. 1994. A multistate outbreak of *Escherichia coli* O157:H7-associated bloody diarrhea and hemolytic uremic syndrome from hamburgers. The Washington experience. *JAMA* **272**:1349–1353.
- Bentley, A. T., and P. E. Klebba. 1988. Effect of lipopolysaccharide structure on reactivity of antiporin monoclonal antibodies with the bacterial cell surface. *J. Bacteriol.* **170**:1063–1068.
- 4a. Bilge, S. S., J. C. Vary, Jr., K. Potter, S. F. Dowell, and P. I. Tarr. 1995. *Escherichia coli* O157:H7 lipopolysaccharide mutants are hyperadherent. *abstr. B-7*, p. 166. In Abstracts of the 95th General Meeting of the American Society for Microbiology 1995. American Society for Microbiology, Washington, D.C.
- Bisercic, M., J. Y. Feutrier, and P. R. Reeves. 1991. Nucleotide sequences of the *gnd* genes from nine natural isolates of *Escherichia coli*: evidence of intragenic recombination as a contributing factor in the evolution of the polymorphic *gnd* locus. *J. Bacteriol.* **173**:3894–3900.
- Blum, G., M. Ott, A. Lischewski, A. Ritter, H. Imrigh, H. Tschape, and J. Hacker. 1994. Excision of large DNA regions termed pathogenicity islands from tRNA-specific loci in the chromosome of an *Escherichia coli* wild-type pathogen. *Infect. Immun.* **62**:606–614.
- Bokete, T. N., C. M. O'Callahan, C. R. Clausen, N. M. Tang, N. Tran, S. L. Moseley, T. R. Fritsche, and P. I. Tarr. 1993. Shiga-like toxin-producing *Escherichia coli* in Seattle children: a prospective study. *Gastroenterology* **105**:1724–1731.
- Chart, H., and B. Rowe. 1993. Antibody cross-reactions with lipopolysaccharide from *E. coli* O157 after cholera vaccination. *Lancet* **341**:1282. (Letter.)
- Cheah, K. C., and P. A. Manning. 1993. Inactivation of the *Escherichia coli* B41 (O101:K99/F41) *rfb* gene encoding an 80-kDa polypeptide results in the synthesis of an antigenically altered lipopolysaccharide in *E. coli* K-12. *Gene* **123**:9–15.
- Devereux, J., P. Haerberli, and O. Smithies. 1984. A comprehensive set of sequence analysis programs for the VAX. *Nucleic Acids Res.* **12**:387–395.
- Gentry, M. K., and J. M. Dalrymple. 1980. Quantitative microtiter cytotoxicity assay for *Shigella* toxin. *J. Clin. Microbiol.* **12**:361–366.
- Gish, W., and D. J. States. 1993. Identification of protein coding regions by database similarity search. *Nat. Genet.* **3**:266–272.
- Gribskov, M., and R. R. Burgess. 1986. Sigma factors from *E. coli*, *B. subtilis*, phage SP01, and phage T4 are homologous proteins. *Nucleic Acids Res.* **14**:6745–6763.
- Griffin, P. M., S. M. Ostroff, R. V. Tauxe, K. D. Greene, J. G. Wells, J. H. Lewis, and P. A. Blake. 1988. Illnesses associated with *Escherichia coli* O157:H7 infections. A broad clinical spectrum. *Ann. Intern. Med.* **109**:705–712.
- Gunzer, F., H. Bohm, H. Russmann, M. Bitzan, S. Aleksic, and H. Karch. 1992. Molecular detection of sorbitol-fermenting *Escherichia coli* O157 in patients with hemolytic-uremic syndrome. *J. Clin. Microbiol.* **30**:1807–1810.
- Inzana, T. J. 1983. Electrophoretic heterogeneity and interstrain variation of the lipopolysaccharide of *Haemophilus influenzae*. *J. Infect. Dis.* **148**:492–499.
- Kessler, A. C., A. Haase, and P. R. Reeves. 1993. Molecular analysis of the 3,6-dideoxyhexose pathway genes of *Yersinia pseudotuberculosis* serogroup IIA. *J. Bacteriol.* **175**:1412–1422.
- Marolda, C. L., and M. A. Valvano. 1993. Identification, expression, and DNA sequence of the GDP-mannose biosynthesis genes encoded by the O7 *rfb* gene cluster of strain VW187 (*Escherichia coli* O7:K1). *J. Bacteriol.* **175**:148–158.
- McDaniel, T. K., K. G. Jarvis, M. S. Donnenberg, and J. B. Kaper. 1995. A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proc. Natl. Acad. Sci. USA* **92**:1664–1668.
- Nelson, K., and R. K. Selander. 1994. Intergeneric transfer and recombination of the 6-phosphogluconate dehydrogenase gene (*gnd*) in enteric bacteria. *Proc. Natl. Acad. Sci. USA* **91**:10227–10231.
- Ochman, H., and R. K. Selander. 1984. Standard reference strains of *Escherichia coli* from natural populations. *J. Bacteriol.* **157**:690–693.
- Perry, M. B., L. MacLean, and D. W. Griffith. 1986. Structure of the O-chain polysaccharide of the phenol-phase soluble lipopolysaccharide of *Escherichia coli* O157:H7. *Biochem. Cell Biol.* **64**:21–28.
- Pierson, D. E. 1994. Mutations affecting lipopolysaccharide enhance Ail-mediated entry of *Yersinia enterocolitica* into mammalian cells. *J. Bacteriol.* **176**:4043–4051.
- Reeves, P. 1993. Evolution of *Salmonella* O antigen variation by interspecific gene transfer on a large scale. *Trends Genet.* **9**:17–22.

25. **Reeves, P. R.** 1995. Personal communication.
26. **Riley, L. W., L. N. Junio, L. B. Libaek, and G. K. Schoolnik.** 1987. Plasmid-encoded expression of lipopolysaccharide O-antigenic polysaccharide in enteropathogenic *Escherichia coli*. *Infect. Immun.* **55**:2052–2056.
27. **Rippey, S. R., L. A. Chandler, and W. D. Watkins.** 1987. Fluorometric method for enumeration of *Escherichia coli* in molluscan shellfish. *J. Food Prot.* **50**:685–690.
28. **Samadpour, M., L. M. Grimm, B. Desai, D. Alf, J. E. Ongerth, and P. I. Tarr.** 1993. Molecular epidemiology of *Escherichia coli* O157:H7 strains by bacteriophage lambda restriction fragment length polymorphism analysis: application to a multistate foodborne outbreak and a day-care center cluster. *J. Clin. Microbiol.* **31**:3179–3183.
29. **Sambrook, J., E. F. Fritsch, and T. Maniatis.** 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
30. **Sanger, F., S. Nicklen, and A. R. Coulson.** 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
31. **Stevenson, G., B. Neal, D. Liu, M. Hobbs, N. H. Packer, M. Batley, J. W. Redmond, L. Lindquist, and P. Reeves.** 1994. Structure of the O antigen of *Escherichia coli* K-12 and the sequence of its *rfb* gene cluster. *J. Bacteriol.* **176**:4144–4156.
32. **Strockbine, N. A., L. R. Marques, J. W. Newland, H. W. Smith, R. K. Holmes, and A. D. O'Brien.** 1986. Two toxin-converting phages from *Escherichia coli* O157:H7 strain 933 encode antigenically distinct toxins with similar biologic activities. *Infect. Immun.* **53**:135–140.
33. **Stroher, U. H., L. E. Karageorgos, R. Morona, and P. A. Manning.** 1992. Serotype conversion in *Vibrio cholerae* O1. *Proc. Natl. Acad. Sci. USA* **89**:2566–2570.
34. **Tarr, P. I.** 1995. *Escherichia coli* O157:H7: clinical, diagnostic, and epidemiological aspects of human infection. *Clin. Infect. Dis.* **20**:1–8.
35. **Tarr, P. I., M. A. Neill, C. R. Clausen, J. W. Newland, R. J. Neill, and S. L. Moseley.** 1989. Genotypic variation in pathogenic *Escherichia coli* O157:H7 isolated from patients in Washington, 1984–1987. *J. Infect. Dis.* **159**:344–347.
36. **Taylor, R. K., C. Manoil, and J. J. Mekalanos.** 1989. Broad-host-range vectors for delivery of *TnphoA*: use in genetic analysis of secreted virulence determinants of *Vibrio cholerae*. *J. Bacteriol.* **171**:1870–1878.
37. **Tsai, C. M., and C. E. Frasch.** 1982. A sensitive silver stain for detecting lipopolysaccharides in polyacrylamide gels. *Anal. Biochem.* **119**:115–119.
38. **van der Ley, P., O. Kuipers, J. Tommassen, and B. Lugtenberg.** 1986. O-antigenic chains of lipopolysaccharide prevent binding of antibody molecules to an outer membrane pore protein in *Enterobacteriaceae*. *Microb. Pathog.* **1**:43–49.
39. **Von Heijne, G.** 1986. A new method for predicting signal sequence cleavage sites. *Nucleic Acids Res.* **14**:4683–4690.
40. **Voorhis, D. L., S. Dillon, S. B. Formal, and R. R. Isberg.** 1991. An O antigen can interfere with the function of the *Yersinia pseudotuberculosis* invasin protein. *Mol. Microbiol.* **5**:317–325.
41. **Wells, J. G., B. R. Davis, I. K. Wachsmuth, L. W. Riley, R. S. Remis, R. Sokolow, and G. K. Morris.** 1983. Laboratory investigation of hemorrhagic colitis outbreaks associated with a rare *Escherichia coli* serotype. *J. Clin. Microbiol.* **18**:512–520.
42. **Whittam, T. S.** 1995. Genetic population structure and pathogenicity in enteric bacteria, p. 217–245. *In* S. Baumberg, J. P. W. Young, S. R. Saunders, and E. M. H. Wellington (ed.), *Population genetics of bacteria*. Cambridge University Press, Cambridge.
43. **Whittam, T. S., M. L. Wolfe, I. K. Wachsmuth, F. Orskov, I. Orskov, and R. A. Wilson.** 1993. Clonal relationships among *Escherichia coli* strains that cause hemorrhagic colitis and infantile diarrhea. *Infect. Immun.* **61**:1619–1629.
44. **Woodall, L. D., P. W. Russell, S. L. Harris, and P. E. Orndorff.** 1993. Rapid, synchronous, and stable induction of type 1 piliation in *Escherichia coli* by using a chromosomal *lacUV5* promoter. *J. Bacteriol.* **175**:2770–2778.

---

*Editor:* P. E. Orndorff