

# The Predicted Amino Acid Sequence of the *Salmonella typhimurium* Virulence Gene *mviA*<sup>+</sup> Strongly Indicates that MviA Is a Regulator Protein of a Previously Unknown *S. typhimurium* Response Regulator Family

WILLIAM H. BENJAMIN, JR.,<sup>1,2\*</sup> XINPING WU,<sup>2</sup> AND W. EDWARD SWORDS<sup>2</sup>

*The Departments of Pathology<sup>1</sup> and Microbiology,<sup>2</sup> The University of Alabama at Birmingham, Birmingham, Alabama 35294*

Received 20 October 1995/Returned for modification 3 January 1996/Accepted 18 March 1996

**The *Salmonella typhimurium* virulence gene *mviA*<sup>+</sup> has a predicted amino acid sequence with homology to the N-terminal 112-amino-acid sequence of response regulator proteins. A previously described mutant allele (*mviA*), which restores virulence to avirulent LT2 strains, was shown to contain a point mutation which would be predicted to cause a single amino acid change, V-102→G (W. H. Benjamin, Jr., J. Yother, P. Hall, and D. E. Briles, *J. Exp. Med.* 174:1073–1083, 1991). A comparison of the nucleotide sequence of *mviA*<sup>+</sup> with that of the *Escherichia coli* and *Salmonella typhi* genes revealed a high degree of conservation.**

We have shown that differences in the *mviA* gene in *Salmonella typhimurium* strains of the LT2 background directly affect the ability of the bacterium to grow in *Ity*<sup>s</sup> (susceptible genotype) but not *Ity*<sup>r</sup> mice (4–6). Thus, an understanding of the mechanism of action of *mviA*<sup>+</sup> may shed light on the resistance mechanism of *Ity* as well as on salmonella virulence. *mviA*<sup>+</sup> was initially characterized by its ability to confer avirulence upon LT2 strains of *S. typhimurium* (6). A comparison of the deduced amino acid sequence of *mviA*<sup>+</sup> with that of known proteins reveals a high degree of homology with a well-characterized family of bacterial response regulatory proteins and suggests that MviA is the second component of a previously unknown two-component response regulatory system. Two-component response regulator systems are commonly used by prokaryotes in environment-dependent control of gene expression (1, 9, 21). On the basis of sequence homology, it is estimated that approximately 40 two-component response regulator systems exist in *Escherichia coli*. However, functional data to support this assignment are available for only a portion of the response regulator-like genes that have been sequenced (1, 18). Several two-component regulatory systems (*phoP/phoQ*, *envZ/ompR*, *bvgS/bvgA*, and *virG/virA*) are known bacterial virulence genes (3, 11–13, 15, 17, 20).

The intact *mviA*<sup>+</sup> allele from an avirulent *S. typhimurium* LT2 strain was cloned and sequenced. The *mviA*<sup>+</sup> sequence was found on a 1.92-kb *Bst*XI-*Cla*I fragment of pWB4184 (6), which includes a 1.57-kb *Dra*I fragment that constitutes the minimal subclone causing avirulence (reference 6 and data not shown). DNA sequencing of double-stranded plasmids was performed with the Sequenase kit (U.S. Biochemicals). On the basis of initial sequence data obtained from the cloned *mviA*<sup>+</sup> gene in plasmid pWB4184, appropriate primers were constructed, and the entire *mviA*<sup>+</sup> gene was amplified by PCR from chromosomal DNA prepared from cultures of *S. typhimurium* by previously described methods (14). PCR products were made single stranded by exonuclease digestion with a presequencing kit (U.S. Biochemicals). The primers used include the following pUC19 universal and reverse primers: *mviAa*, 5' ATGCAGCAAGATTTGCTCTC 3' (nucleotides

[nt] 53 to 72); *mviAb*, 5' ACCAATGGCTGCGTCATGTT 3' (nt 506 to 487); *mviAc*, 5' TTTAACGGCTTATTGCAGGA 3' (nt 1156 to 1175); *mviAd*, 5' CATTCCGCAGACAACATCAA 3' (nt 1503 to 1489); *mviAe*, 5' AGCAAATTCTGACAGGC GCA 3' (nt 422 to 431); *mviAf*, 5' GGGGGTGAAGATGTC TTGCT 3' (nt 789 to 809); and *mviAg*, 5' TCAACCCGCG AGTTAAACAT 3' (nt 895 to 875). A single open reading frame was identified which is predicted to encode a protein 337 amino acids in length with a molecular mass of 37,292 Da and with a pI of 5.5. No obvious transmembrane or leader sequences were noted. A typical promoter [–35 (TATAACC) N<sub>17</sub> (TTTAAT)] was identified 89 bp upstream of the predicted start codon of the open reading frame (16).

GenBank searches yielded no significant homology at the nucleotide level, except for the apparent *mviA*<sup>+</sup> homolog from *E. coli*, which was previously designated *orf37* (7, 8). Because of the high degree of homology and an almost identical chromosomal map location, we believe that *orf37* is a homolog of *mviA*<sup>+</sup>; hereafter, we refer to it as *E. coli mviA*<sup>+</sup>. The 110 best matches to *mviA*<sup>+</sup>, obtained from a TfastA search with the 112 N-terminal amino acids predicted from the *mviA*<sup>+</sup> sequence were analyzed, and 84 unique sequences were identified. Of these, 77 were identified as having the response regulator motif, on the basis of the invariant amino acids corresponding to MviA residues D-15, D-58, T- or S-83, and K-108 (21, 22). The deduced sequence was 90% identical to the *E. coli* MviA predicted protein product (Fig. 1). In contrast to *E. coli* MviA, the additional 76 known and apparent response regulators had 20 to 37% amino acid sequence homology with MviA over at least 98 amino acids. In each case, the region of homology covered most of the N-terminal 112 amino acids of MviA (Fig. 1). The hydrophobic regions described by Stock et al. (21, 22) were also highly conserved between MviA and the response regulator protein sequences. By using the seven *mviA* oligonucleotide primers, *mviA*<sup>+</sup> homologs were amplified from *E. coli* (strain LE392) and two strains of *Salmonella typhi* (the virulent strain Q5 and the avirulent vaccine strain Ty21a). The PCR products were sequenced as described above, and the nucleotide sequences obtained were compared with that of *mviA*<sup>+</sup> from *S. typhimurium*. Q5 was found to have a single nucleotide difference, in codon 98 (TTG → TTA); this is a silent mutation, because it has no effect on the predicted product. Ty21a was

\* Corresponding author. Phone: (205) 934-1923. Fax: (205) 934-0605. Electronic mail address: Benjamin@wp.path.uab.edu.



fore be useful in the rational construction of attenuated strains for use as live vaccines.

**Nucleotide sequence accession number.** All sequences obtained in this study have been submitted to GenBank under accession number U53173.

This work was supported by a grant from the National Institutes of Health (RO1 AI15946) to William H. Benjamin. W. Edward Swords was supported in part by training grant T32 AI107041 from the NIH. Genetics Computer Group sequence analysis was made available through the computer resources of the University of Alabama at Birmingham Center for AIDS Research (P30 AI27767).

We thank David E. Briles for assistance and Rebecca C. Tart and Melissa Caimano for critical reading of the manuscript.

#### REFERENCES

- Albright, L. M., E. Huala, and F. M. Ausubel. 1989. Prokaryotic signal transduction mediated by sensor and regulator protein pairs. *Annu. Rev. Genet.* **23**:311–336.
- Alexeyev, M. F. 1995. Three kanamycin resistance gene cassettes with different polylinkers. *BioTechniques* **18**:52–55.
- Aricó, B., V. Scarlato, D. M. Monack, S. Falkow, and R. Rappuoli. 1991. Structural and genetic analysis of the *bvg* locus in *Bordetella* species. *Mol. Microbiol.* **5**:2481–2491.
- Benjamin, W. H., Jr., C. L. Turnbough, Jr., J. D. Goguen, B. S. Posey, and D. E. Briles. 1986. Genetic mapping of novel virulence determinants of *Salmonella typhimurium* to the region between *trpD* and *supD*. *Microb. Pathog.* **1**:115–124.
- Benjamin, W. H., Jr., C. L. Turnbough, Jr., B. S. Posey, and D. E. Briles. 1986. *Salmonella typhimurium* virulence genes necessary to exploit the *Ity<sup>sis</sup>* genotype of the mouse. *Infect. Immun.* **51**:872–878.
- Benjamin, W. H., Jr., J. Yother, P. Hall, and D. E. Briles. 1991. The *Salmonella typhimurium* locus *mviA* regulates virulence in *Ity<sup>3</sup>* but not *Ity<sup>1</sup>* mice: functional *mviA* results in avirulence; mutant (nonfunctional) *mviA* results in virulence. *J. Exp. Med.* **174**:1073–1083.
- Bösl, M., and H. Kersten. 1994. Organization and functions of genes in the upstream region of *tyrT* of *Escherichia coli*: phenotypes of mutants with partial deletions of a new gene (*tgs*). *J. Bacteriol.* **176**:221–231.
- Bösl, M. R. 1993. Genetic map of the *tyrT* region of *Escherichia coli* from 27.1 to 27.7 minutes based exclusively on sequence data. *J. Bacteriol.* **175**:7751–7753.
- Cozzone, A. 1993. ATP-dependent protein kinases in bacteria. *J. Cell. Biochem.* **51**:7–13.
- Dahl, M. K., T. Msadek, F. Kunst, and G. Rapoport. 1991. Mutational analysis of the *Bacillus subtilis* DegU regulator and its phosphorylation by the DegS protein kinase. *J. Bacteriol.* **173**:2539–2547.
- Dorman, C. J., S. Chatfield, C. F. Higgins, C. Hayward, and G. Dougan. 1989. Characterization of porin and *ompR* mutants of a virulent strain of *Salmonella typhimurium*: *ompR* mutants are attenuated in vivo. *Infect. Immun.* **57**:2136–2140.
- Dziejman, M., and J. J. Mekalanos. 1995. Two-component signal transduction and its role in the expression of bacterial virulence factors, p. 305–317. In J. A. Hoch and T. J. Silhavy (ed.), *Two-component signal transduction*. ASM Press, Washington, D.C.
- Fields, P. I., E. A. Groisman, and F. Heffron. 1989. A *Salmonella* locus that controls resistance to microbiocidal proteins from phagocytic cells. *Science* **243**:1059–1062.
- Gay, N. J. 1984. Construction and characterization of an *Escherichia coli* strain with a *uncI* mutation. *J. Bacteriol.* **158**:820–825.
- Groisman, E. A., E. Chiao, C. J. Lipps, and F. Heffron. 1989. *Salmonella typhimurium* *phoP* virulence gene is a transcriptional regulator. *Proc. Natl. Acad. Sci. USA* **86**:7077–7081.
- Harr, R., M. Haggström, and P. Gustafsson. 1983. Search algorithm for pattern match analysis of nucleic acid sequences. *Nucleic Acids Res.* **11**:2943–2957.
- Mekalanos, J. J. 1992. Environmental signals controlling expression of virulence determinants in bacteria. *J. Bacteriol.* **174**:1–7.
- Parkinson, J. S., and E. C. Kofoid. 1992. Communication modules in bacterial signaling proteins. *Annu. Rev. Genet.* **26**:71–112.
- Pazour, G. J., C. N. Ta, and A. Das. 1992. Constitutive mutations of *Agrobacterium tumefaciens* transcriptional activator *virG*. *J. Bacteriol.* **174**:4169–4174.
- Stibitz, S., A. A. Weiss, and S. Falkow. 1988. Genetic analysis of a region of the *Bordetella pertussis* chromosome encoding filamentous hemagglutinin and the pleiotropic regulatory locus *vir*. *J. Bacteriol.* **170**:2904–2913.
- Stock, J. B., A. J. Ninfa, and A. M. Stock. 1989. Protein phosphorylation and regulation of adaptive responses in bacteria. *Microbiol. Rev.* **53**:450–490.
- Stock, J. B., A. M. Stock, and J. M. Mottonen. 1990. Signal transduction in bacteria. *Nature (London)* **344**:395–400.
- Zubay, G. 1973. In vitro synthesis of protein in microbial systems. *Annu. Rev. Genet.* **7**:267–287.

Editor: A. O'Brien