

## Intrapulmonary *Hartmannella vermiformis*: a Potential Niche for *Legionella pneumophila* Replication in a Murine Model of Legionellosis

JOAN BRIELAND,<sup>1\*</sup> MARK McCLAIN,<sup>2</sup> MAUREEN LEGENDRE,<sup>3</sup> AND CARY ENGLEBERG<sup>2,4</sup>

Unit for Laboratory Animal Medicine<sup>1</sup> and Departments of Microbiology and Immunology,<sup>2</sup> Pathology,<sup>3</sup> and Medicine,<sup>4</sup> University of Michigan Medical School, Ann Arbor, Michigan 48109-0614

Received 18 April 1997/Returned for modification 28 May 1997/Accepted 29 August 1997

The potential role of inhaled protozoa as a niche for intrapulmonary replication of *Legionella pneumophila* was investigated in vivo with mutant strains of *L. pneumophila* which have reduced virulence for the amoeba *Hartmannella vermiformis*. *L. pneumophila* AA488 and AA502 were derived from wild-type strain AA100 after transposon mutagenesis. These mutants have reduced virulence for *H. vermiformis* but are fully virulent for mononuclear phagocytic cells. A/J mice, which are susceptible to replicative *L. pneumophila* lung infections, were inoculated intratracheally with *L. pneumophila* AA100, AA488, or AA502 ( $10^6$  bacteria per mouse) or were coinoculated with one of the *L. pneumophila* strains ( $10^6$  bacteria per mouse) and uninfected *H. vermiformis* ( $10^6$  amoebae per mouse). The effect of coinoculation with *H. vermiformis* on intrapulmonary growth of each *L. pneumophila* strain was subsequently assessed. In agreement with our previous studies, coinoculation with *H. vermiformis* significantly enhanced intrapulmonary growth of the parent *L. pneumophila* strain (AA100). In contrast, intrapulmonary growth of *L. pneumophila* AA488 or AA502 was not significantly enhanced by coinoculation of mice with *H. vermiformis*. These studies demonstrate that *L. pneumophila* virulence for amoebae is required for maximal intrapulmonary growth of the bacteria in mice coinoculated with *H. vermiformis* and support the hypothesis that inhaled amoebae may potentiate intrapulmonary growth of *L. pneumophila* by providing a niche for bacterial replication.

*Legionella pneumophila* is a bacterial parasite of many species of freshwater protozoa and an occasional intracellular pathogen of humans (2, 3, 21, 24, 26, 27, 29, 32, 36, 42). Infection of humans with *L. pneumophila* usually occurs after bacteria from natural aquatic sources contaminate potable and nonpotable water systems. The portal of entry is the lung, via inhalation of aerosols or microaspiration of contaminated water (10, 14–16). Following inhalation, the bacteria invade and replicate within host mononuclear phagocytic cells, primarily within alveolar macrophages. The intracellular infection of alveolar macrophages that is characteristic of Legionnaires' disease is remarkably similar in its ultrastructural features to *L. pneumophila* infection of protozoal organisms (1, 23, 24, 26, 32, 36, 40, 41).

Since the discovery of the bacterium-protozoan interaction, there has been much speculation but limited research directed towards elucidating the relevance of protozoa to human *L. pneumophila* infection (36). Protozoa play a key role in the persistence of *L. pneumophila* in natural aquatic water systems, since growth of *L. pneumophila* in the environment in the absence of these organisms has rarely been documented (22, 37, 43). Protozoa may also serve as a vehicle to protect *L. pneumophila* through the process of water treatment and/or serve as a reservoir for growth of the bacteria in potable water systems (4). Amoebae which are permissive for growth of *L. pneumophila* are frequently found in water implicated as the source of the bacteria in outbreaks of Legionnaire's disease (4, 10, 11, 19, 20). It is not known whether the presence of amoebae is a prerequisite for an outbreak, since contamination of potable

water systems with protozoa is exceedingly common (23, 36, 42).

Using an animal model of replicative *L. pneumophila* lung infection in A/J mice inoculated intratracheally with virulent bacteria, we have recently demonstrated that protozoa may have a more direct effect on the pathogenesis of replicative *L. pneumophila* lung infection. Specifically, we have shown that intrapulmonary growth of *L. pneumophila* in A/J mice is potentiated by coinoculation with *Hartmannella vermiformis*, an amoebal species which is permissive for growth of the bacteria and a natural reservoir of *L. pneumophila* in the environment (7). These results support the hypothesis that coincident inhalation of protozoa may enhance the severity of Legionnaire's disease. However, the mechanism(s) by which intrapulmonary *H. vermiformis* organisms potentiate *L. pneumophila* replication in vivo has not been thoroughly investigated.

In the present study, the potential role of intrapulmonary *H. vermiformis* as a niche for *L. pneumophila* replication was assessed with mutant strains of *L. pneumophila* which have reduced virulence for *H. vermiformis* and a previously described murine model of replicative *L. pneumophila* lung infection in A/J mice (8). We demonstrate that bacterial virulence for *H. vermiformis* is required for maximal intrapulmonary growth of *L. pneumophila* in A/J mice coinoculated with the amoebae. These results support the hypothesis that intrapulmonary *H. vermiformis* organisms likely potentiate growth of *L. pneumophila* in vivo by their ability to provide a niche for *L. pneumophila* replication.

A virulent strain of *L. pneumophila* (AA100) originally isolated from an outbreak of Legionnaires' disease was used in all assays (8). The bacteria were maintained and passaged on buffered charcoal-yeast extract (BCYE) agar (18). The amoeba *H. vermiformis* CDC-19 (ATCC 50237), originally isolated from a water sample obtained during an investigation of nos-

\* Corresponding author. Mailing address: 018 Animal Research Facility, University of Michigan Medical School, Ann Arbor, MI 48109-0614. Phone: (313) 764-0277. Fax: (313) 936-3235.

ocomial legionellosis, was maintained axenically as confluent monolayers of trophozoites in T-75 flasks in ATCC culture medium 1034 supplemented with 10% fetal bovine serum as previously described (19).

To identify *L. pneumophila* mutants with reduced virulence for *H. vermiformis*, phosphatase-positive ( $\text{Pho}^+$ ) mutant strains of *L. pneumophila* were generated by transposon insertion mutagenesis of *L. pneumophila* AA100, with the previously described alkaline phosphatase fusion-generating transposon mTn10*phoA* (28). The results of these experiments identified approximately 4,000 mutagenized legionellae, yielding 206  $\text{Pho}^+$  *L. pneumophila* mutant bacterial strains. The ability of each  $\text{Pho}^+$  mutant *L. pneumophila* strain to grow in U937 cells (a human monocytic cell line which is permissive for *L. pneumophila* growth [12, 33]) and *H. vermiformis* was subsequently assessed. Briefly, differentiated U937 cells were inoculated with the parent bacterial strain (AA100) or with a  $\text{Pho}^+$  strain of *L. pneumophila* at a multiplicity of infection (MOI) of 20:1 for 2 h at 37°C. *L. pneumophila*-infected U937 cell monolayers were then washed and incubated for an additional 2 h in media containing gentamicin (50 µg/ml) to kill extracellular bacteria. The cells were subsequently washed and either immediately hypotonically lysed or incubated for an additional 24 to 72 h (37°C, 5%  $\text{CO}_2$ ). Following incubation, the culture supernatant was removed, the cells were hypotonically lysed, and the corresponding cell supernatant and lysate were pooled. An aliquot of each sample was serially diluted in water and cultured on BCYE. The number of *L. pneumophila* CFU per sample, indicative of growth of  $\text{Pho}^+$  *L. pneumophila* mutants in U937 cells, was determined and compared to that of the parent *L. pneumophila* strain (AA100). To assess the virulence of each *L. pneumophila* strain for *H. vermiformis*, axenically maintained *H. vermiformis* monolayers were inoculated with each mutant *L. pneumophila* strain at an MOI of 1:1. Immediately after inoculation, an aliquot of the culture supernatant was removed and serially diluted, and *L. pneumophila* CFU in the inocula were determined. *L. pneumophila*-infected *H. vermiformis* monolayers were then incubated for 24 to 72 h, at which time an aliquot of the culture supernatant was removed and cultured on BCYE. The number of *L. pneumophila* CFU in culture supernatant, indicative of growth of  $\text{Pho}^+$  *L. pneumophila* mutants in *H. vermiformis*, was determined and compared to that of the parent *L. pneumophila* strain (AA100). As shown in Fig. 1, these screening methods identified three  $\text{Pho}^+$  *L. pneumophila* strains, designated AA488, AA501, and AA502, which are as virulent as *L. pneumophila* AA100 for cultured mononuclear phagocytic cells but have significantly reduced virulence for *H. vermiformis*.

Virulence of *L. pneumophila* AA488, AA501, and AA502 in vivo was subsequently assessed with a previously described murine model of legionellosis in intratracheally inoculated A/J mice (8). For preparation of in vivo inocula, each *L. pneumophila* strain was grown on BCYE agar for 48 h and resuspended in phosphate-buffered saline ( $4 \times 10^7$  bacteria per ml). Female 6- to 8-week-old A/J mice (Jackson Laboratories, Bar Harbor, Maine), which are susceptible to replicative *L. pneumophila* lung infections (8), were subsequently inoculated intratracheally with the parent bacterial strain (AA100) or with one of the three mutant *L. pneumophila* strains ( $10^6$  bacteria per mouse) (8). At 48 h postinoculation, the mice were humanely euthanized, the lungs were excised and homogenized, and *L. pneumophila* CFU per lung were determined by culture of lung homogenate on bacteriological media (8, 17). (The mice were euthanized at this time point because the results of our previous studies have demonstrated that intrapulmonary growth of *L. pneumophila* AA100 in intratracheally inoculated A/J mice




FIG. 1. Kinetics of growth of  $\text{Pho}^+$  *L. pneumophila* strains for mammalian cells and amoebae. (A) Growth of  $\text{Pho}^+$  *L. pneumophila* strains in differentiated U937 cells was monitored as previously described and compared to that of the parent bacterial strain (AA100). (B) Growth of  $\text{Pho}^+$  *L. pneumophila* strains in the amoeba *H. vermiformis* was determined as previously described and compared to that of the parent bacterial strain (AA100). Data for both panels A and B represent the mean log increase in CFU from duplicate wells of a minimum of two separate experiments. \*, significant difference in growth compared with that in *L. pneumophila* AA100 (Student's *t* test [ $P < 0.05$ ]).

is maximal at 48 h postinoculation [8].) As shown in Fig. 2, while there was no significant difference between intrapulmonary growth of either *L. pneumophila* AA488 or AA502 and the parent bacterial strain (AA100), significantly fewer *L. pneumophila* CFU were recovered from lung homogenates of mice inoculated intratracheally with *L. pneumophila* AA501. These results demonstrate that *L. pneumophila* AA501 has reduced virulence in vivo. Consequently, this strain was eliminated from further analysis.

In subsequent experiments, the growth kinetics of *L. pneumophila* AA488, AA502, and AA100 in the lung were compared. A/J mice were inoculated intratracheally with either *L. pneumophila* AA488, AA502, or AA100 ( $10^6$  bacteria per mouse). At 24 to 72 h postinoculation, the mice were humanely euthanized and *L. pneumophila* CFU in lung homogenates were determined by culture of lung homogenate on bacteriological media (8, 17). As shown in Fig. 3, the levels of intrapulmonary growth of *L. pneumophila* AA488 and AA502 were similar to that of *L. pneumophila* AA100. These results dem-




FIG. 2. Intrapulmonary growth of  $\text{Pho}^+$  *L. pneumophila* strains in A/J mice at 48 h postinoculation. A/J mice were inoculated intratracheally with a  $\text{Pho}^+$  *L. pneumophila* strain (AA488, AA501, or AA502) which is virulent for U937 cells but has reduced virulence for *H. vermiformis* in vitro ( $10^6$  bacteria per mouse). At 48 h postinoculation, the mice were euthanized, and the lungs were excised and homogenized. *L. pneumophila* CFU per lung were subsequently determined by culture of lung homogenates on bacteriological media. Results represent the mean  $\pm$  standard error of 8 to 22 mice per treatment group. \*, significant difference in growth compared with that in mice inoculated with *L. pneumophila* AA100 (Mann-Whitney test [ $P < 0.05$ ]).

onstrate that *L. pneumophila* AA488 and AA502 are fully virulent in vivo.

The effect of coinoculation with *H. vermiformis* on the intrapulmonary growth of each *L. pneumophila* strain was subsequently assessed. A/J mice were inoculated intratracheally with the parent strain of *L. pneumophila*, AA100 (which is virulent for both mononuclear phagocytic cells and *H. vermiformis* [ $10^6$  bacteria per mouse]), or with  $\text{Pho}^+$  *L. pneumophila* AA488 or AA502 (strains which are fully virulent for mononuclear phagocytic cells in vivo but have reduced virulence for *H. vermiformis* [ $10^6$  bacteria per mouse]), or they were coin-




FIG. 3. Kinetics of intrapulmonary growth of  $\text{Pho}^+$  *L. pneumophila* strains in A/J mice. A/J mice were inoculated intratracheally with *L. pneumophila* AA488, AA502, or AA100 ( $10^6$  bacteria per mouse). At 24 to 72 h postinoculation, the mice were euthanized, and the lungs were excised and homogenized. *L. pneumophila* CFU per lung were subsequently determined by culture of lung homogenates on bacteriological media. Results represent the mean  $\pm$  standard error of 3 to 22 mice per treatment group.

TABLE 1. Effect of *H. vermiformis* on intrapulmonary growth of *L. pneumophila* strains at 48 h postinoculation

Inoculum	<i>L. pneumophila</i> CFU/lung 48 h postinoculation <sup>a</sup>
AA100	$(9.68 \pm 4.63) \times 10^7$
AA100+ amoebae	$(2.10 \pm 0.58) \times 10^{9b}$
AA488	$(6.51 \pm 3.33) \times 10^7$
AA488+ amoebae	$(1.97 \pm 1.0) \times 10^8$
AA502	$(2.01 \pm 1.21) \times 10^8$
AA502+ amoebae	$(1.32 \pm 0.73) \times 10^9$

<sup>a</sup> Results represent mean  $\pm$  standard error of 7 to 22 mice per treatment group.

<sup>b</sup> Significant increase in intrapulmonary growth of *L. pneumophila* in mice coinoculated with *H. vermiformis* compared with that in mice inoculated with the same strain of bacteria alone (Mann-Whitney test [ $P < 0.05$ ]).

oculated with these bacterial strains ( $10^6$  bacteria per mouse) and *H. vermiformis* ( $10^6$  amoebae per mouse) as previously described (7, 8, 38). At 48 h postinoculation, the mice were humanely euthanized and the lungs were excised and homogenized. *L. pneumophila* CFU per lung were determined by culture of lung homogenates on bacteriological media. Bacterial growth was assessed at 48 h postinoculation, because we have previously demonstrated that growth of *L. pneumophila* in mice coinoculated with *H. vermiformis* and strain AA100 is maximal at this time point (7). In agreement with the results of our previous studies (7), coinoculation of A/J mice with *H. vermiformis* significantly enhanced intrapulmonary growth of the parent strain of *L. pneumophila* (AA100) (Table 1). In contrast, coinoculation with *H. vermiformis* did not result in a significant increase in intrapulmonary growth of *L. pneumophila* AA488 or AA502 compared to that in mice inoculated with the same strain of bacteria alone. These results demonstrate that *L. pneumophila* virulence for *H. vermiformis* is required for maximal enhancement in intrapulmonary growth of the bacteria in mice coinoculated with the amoebae and support the hypothesis that intrapulmonary *H. vermiformis* may enhance growth of *L. pneumophila* in the lung by providing a niche for bacterial replication.

However, our results also demonstrate that there were small increases (i.e., three- and sixfold, respectively) in the growth of *L. pneumophila* in mice coinoculated with *H. vermiformis* and *L. pneumophila* AA488 or AA502 at 48 h postinfection compared to that in mice inoculated with the same strain of bacteria. While this increase in bacterial growth is not statistically significant, these results suggest that intrapulmonary *H. vermiformis* may potentiate the growth of *L. pneumophila*, at least in part, by a mechanism which is independent of *L. pneumophila* virulence for *H. vermiformis*. We have previously demonstrated that intrapulmonary *H. vermiformis* alters host immune responses during replicative *L. pneumophila* lung infections, resulting in exaggerated production of cytokines, including gamma interferon and tumor necrosis factor alpha (7). Previous studies by our laboratory and others have demonstrated that mononuclear phagocytic cell activation, mediated in large part by gamma interferon and tumor necrosis factor alpha, is required for resolution of replicative *L. pneumophila* infection in permissive mononuclear phagocytic cells (5, 6, 8, 9, 25, 31). However, exaggerated production of proinflammatory cytokines, including tumor necrosis factor alpha, has been shown to be detrimental to resolution of other inflammatory lung diseases, including adult respiratory distress syndrome (30, 39). The potential relationship between *H. vermiformis*-induced altered host immune responses and enhanced intrapulmonary

growth of *L. pneumophila* in mice coinoculated with the amoebae remains to be thoroughly investigated.

Alternatively, it is possible that growth of *L. pneumophila* strains (which are virulent for amoebae) in intrapulmonary *H. vermiformis* may enhance bacterial virulence for mammalian cells, thereby facilitating growth of the bacteria in vivo. Previous in vitro studies have demonstrated that *L. pneumophila*, following growth within *Acanthamoeba castellanii*, another freshwater protozoan, has enhanced invasiveness for cultured mammalian cells (13). The effect of growth of *L. pneumophila* in intrapulmonary *H. vermiformis* on virulence of the bacteria in vivo remains to be investigated.

In summary, we have isolated mutant strains of *L. pneumophila* which have reduced virulence for *H. vermiformis* but are fully virulent for mammalian cells in vivo. These mutant *L. pneumophila* strains suggest that different bacterial gene products are likely required for *L. pneumophila* infection of mononuclear phagocytic cells and freshwater protozoa. We have previously shown that coinoculation of A/J mice with the amoeba *H. vermiformis* potentiates intrapulmonary growth of *L. pneumophila*, thereby demonstrating that coincident inhalation of protozoa may enhance the severity of replicative *L. pneumophila* lung infection (7). The results of our current study demonstrate that *L. pneumophila* virulence for *H. vermiformis* is required for maximal intrapulmonary growth of the bacteria. These results support the hypothesis that intrapulmonary amoebae may facilitate growth of *L. pneumophila* in the lung, at least in part, by providing a niche for bacterial replication in vivo. However, this mechanism is not mutually exclusive, because *H. vermiformis*-induced changes in host immune responses and/or increased bacterial virulence, resulting from intra-amoebic growth of the bacteria, may also contribute to enhanced intrapulmonary growth of *L. pneumophila*.

While our results support a role of coinhaled protozoa in the pathogenesis of experimentally induced legionellosis, the potential role of inhaled amoebae in the pathogenesis of Legionnaires' disease in humans remains unknown. It is likely that people may inhale *H. vermiformis*, because amoebae can be aerosolized (35) and asymptomatic persons frequently have titers of antibody to amoebal species, including *H. vermiformis*, indicative of human exposure (34). However, it is likely impossible to validate a role of inhaled *H. vermiformis* in the pathogenesis of legionellosis in humans by identification of amoebae in the lung. We have previously demonstrated that *H. vermiformis* is identifiable in the murine lung only during the initial phase of replicative *L. pneumophila* lung infection (i.e., at <72 h postinoculation [7]). Diagnostic tests of human patients suspected of having Legionnaires' disease are conducted considerably later in the course of the infection, potentially after inhaled amoebae have been cleared from the lung. Nevertheless, the results of our present studies indicate that further investigation into the potential role of coinhaled protozoa in the pathogenesis of Legionnaires' disease in humans is warranted. It is anticipated that the results of these studies may have significant implications with regard to the design of strategies to prevent, control, and/or treat human legionellosis.

This work was supported by Public Health Service grants NIH R29-HL-49136 and RR00200.

#### REFERENCES

1. Abu Kwaik, Y. 1996. The phagosome containing *Legionella pneumophila* within the protozoan *Hartmannella vermiformis* is surrounded by the rough endoplasmic reticulum. *Appl. Environ. Microbiol.* **62**:2002-2028.
2. Abu Kwaik, Y., B. S. Fields, and N. C. Engleberg. 1994. Protein expression by the protozoan *Hartmannella vermiformis* upon contact with its bacterial parasite *Legionella pneumophila*. *Infect. Immun.* **62**:1860-1866.
3. Anand, C., A. R. Skinner, A. Malic, and J. B. Kurtz. 1983. Interaction of

- Legionella pneumophila* and a free living amoebae (*Acanthamoeba palestinensis*). *J. Hyg. Camb.* **91**:167-178.
4. Barbaree, J. M., B. S. Fields, J. C. Feeley, G. W. Gorman, and W. T. Martin. 1986. Isolation of protozoa from water associated with a legionellosis outbreak and demonstration of intracellular multiplication of *Legionella pneumophila*. *Appl. Environ. Microbiol.* **51**:422-424.
5. Bhardwaj, N., T. W. Nash, and M. A. Horwitz. 1986. Interferon gamma activated human monocytes inhibit intracellular multiplication of *Legionella pneumophila*. *J. Immunol.* **137**:2662-2669.
6. Blanchard, D. K., J. Y. Djeu, T. W. Klein, H. Friedman, W. E. Stewart II, and W. E. Stewart. 1988. Protective effects of tumor necrosis factor in experimental *Legionella pneumophila* infections in mice via activation of PMN function. *J. Leukocyte Biol.* **43**:429-435.
7. Brieland, J., M. McClain, L. Heath, C. Chrisp, G. Huffnagle, M. LeGendre, M. Hurley, J. Fantone, and C. Engleberg. 1996. Coinoculation with *Hartmannella vermiformis* enhances replicative *Legionella pneumophila* lung infection in a murine model of Legionnaires' disease. *Infect. Immun.* **64**:2449-2456.
8. Brieland, J. K., P. Freeman, R. Kunkel, C. Chrisp, M. Hurley, J. Fantone, and C. Engleberg. 1994. Replicative *Legionella pneumophila* lung infection in intratracheally inoculated A/J mice. A murine model of human Legionnaires' disease. *Am. J. Pathol.* **145**:1537-1546.
9. Brieland, J. K., D. G. Remick, P. T. Freeman, M. C. Hurley, J. C. Fantone, and N. C. Engleberg. 1995. In vivo regulation of replicative *Legionella pneumophila* lung infection by endogenous tumor necrosis factor alpha and nitric oxide. *Infect. Immun.* **63**:3253-3258.
10. Brieman, R. F. 1993. Modes of transmission in epidemic and non-epidemic *Legionella* infection: directions for further study, p. 30-35. In J. M. Barbaree, R. F. Brieman, and A. P. Dufour (ed.), *Legionella*: current status and emerging perspectives. American Society for Microbiology, Washington, D.C.
11. Brieman, R. F., B. S. Fields, G. N. Sanden, L. Volmer, A. Meier, and J. S. Spika. 1990. Association of shower use with Legionnaires' disease: possible role of amoebae. *JAMA* **263**:2924-2926.
12. Cianciotto, N. P., B. I. Eisenstein, C. H. Mody, G. B. Toews, and N. C. Engleberg. 1989. A *Legionella pneumophila* gene encoding a species-specific surface protein potentiates initiation of intracellular infection. *Infect. Immun.* **57**:1255-1262.
13. Cirillo, J. D., S. Falkow, and L. S. Tompkins. 1994. Growth of *Legionella pneumophila* in *Acanthamoeba castellanii* enhances invasion. *Infect. Immun.* **62**:3254-3261.
14. Davis, G. S., W. C. Winn, D. W. Gump, J. E. Craighead, and H. N. Beaty. 1982. Legionnaires' pneumonia after aerosol exposure in guinea pigs and rats. *Am. Rev. Respir. Dis.* **126**:1050-1057.
15. Davis, G. S., W. C. Winn, and H. N. Beaty. 1981. Legionnaires' disease. Infections caused by *Legionella pneumophila* and legionella-like organisms. *Clin. Chest Med.* **2**:145-166.
16. Dondoro, T. J., Jr., R. C. Rendtorff, G. F. Mallison, R. M. Weeks, J. S. Levy, E. W. Wong, and W. Schaffner. 1980. An outbreak of Legionnaires' disease associated with a contaminated air-conditioning cooling tower. *N. Engl. J. Med.* **302**:365-370.
17. Edelstein, P. H. 1981. Improved semiselective medium for isolation of *Legionella pneumophila* from contaminated clinical and environmental specimens. *J. Clin. Microbiol.* **14**:298-303.
18. Feeley, J. C., R. J. Gibson, G. W. Gorman, N. C. Langford, J. K. Rasheed, C. D. Mackel, and W. B. Baine. 1979. Charcoal yeast extract agar: primary isolation medium for *Legionella pneumophila*. *J. Clin. Microbiol.* **10**:437-441.
19. Fields, B. S., T. A. Nerad, T. K. Sawyer, C. H. King, J. M. Barbaree, W. T. Martin, W. E. Morrill, and G. N. Sanden. 1990. Characterization of an axenic strain of *Hartmannella vermiformis* obtained from an investigation of nosocomial legionellosis. *J. Protozool.* **37**:581-583.
20. Fields, B. S., G. N. Sanden, and J. M. Barbaree. 1989. Intracellular multiplication of *Legionella pneumophila* in amoebae isolated from hospital hot water tanks. *Curr. Microbiol.* **18**:131-139.
21. Fields, B. S., E. B. Shotts, Jr., J. C. Feeley, G. W. Gorman, and W. T. Martin. 1984. Proliferation of *Legionella pneumophila* as an intracellular parasite of the ciliated protozoan *Tetrahymena pyriformis*. *Appl. Environ. Microbiol.* **47**:467-471.
22. Fields, B. S., S. R. Fields, J. N. Loy, E. H. White, S. L. Steffens, and E. B. Shotts. 1993. Attachment and entry of *Legionella pneumophila* in *Hartmannella vermiformis*. *J. Infect. Dis.* **167**:1146-1150.
23. Henke, M., and K. M. Seidel. 1986. Association between *Legionella pneumophila* and amoebae in water. *Isr. J. Med. Sci.* **22**:690-695.
24. Holden, E. P., H. H. Winkler, D. O. Wood, and E. D. Leinbach. 1984. Intracellular growth of *Legionella pneumophila* within *Acanthamoeba castellanii* Neff. *Infect. Immun.* **45**:18-24.
25. Horwitz, M. A., and S. C. Silverstein. 1981. Activated human monocytes inhibit the intracellular multiplication of Legionnaires' disease bacteria. *J. Exp. Med.* **154**:1618-1635.
26. Horwitz, M. A. 1983. The Legionnaires' disease bacterium (*Legionella pneumophila*) inhibits phagosome-lysosome fusion in human monocytes. *J. Exp. Med.* **158**:2108-2126.
27. Horwitz, M. A., and S. C. Silverstein. 1980. Legionnaires' disease bacterium

- (*Legionella pneumophila*) multiplies intracellularly in human monocytes. *J. Clin. Invest.* **66**:441–450.
28. McClain, M. S., and N. C. Engleberg. 1996. Construction of an alkaline phosphatase fusion-generating transposon, mTn10*phoA*. *Gene* **170**:147–148.
  29. McDade, J. E., C. C. Shepard, D. W. Fraser, T. R. Tsai, M. A. Redus, and W. R. Dowdle. 1977. Legionnaires' disease: isolation of a bacterium and demonstration of its role in other respiratory disease. *N. Engl. J. Med.* **297**:1197–1203.
  30. Meduri, G. U., G. Kohler, S. Headley, E. Tolley, F. Stentz, and A. Postlethwaite. 1995. Inflammatory cytokines in the BAL in patients with ARDS. Persistent elevation over time predicts poor outcome. *Chest* **108**:1303–1314.
  31. Nash, T. W., D. M. Libby, and M. A. Horwitz. 1988. IFN-gamma activated human alveolar macrophages inhibit the intracellular multiplication of *Legionella pneumophila*. *J. Immunol.* **140**:3978–3981.
  32. Newsome, A. L., R. L. Baker, R. D. Miller, and R. R. Arnold. 1985. Interactions between *Naegleria fowleri* and *Legionella pneumophila*. *Infect. Immun.* **50**:449–452.
  33. Pearlman, E., A. Jiwa, N. C. Engleberg, and B. I. Eisenstein. 1988. Growth of *Legionella pneumophila* in a human macrophage-like (U937) cell line. *Microb. Pathog.* **5**:87–95.
  34. Powell, E., A. L. Newsome, and S. D. Allen. 1994. Free-living amoeba antigens recognized by naturally occurring human antibodies, abstr. C-374, p. 512. *In Abstracts of the 94th General Meeting of the American Society for Microbiology* 1993. American Society for Microbiology, Washington, D.C.
  35. Rodriguez-Zaragoza, S., P. Bonilla, E. Ramirez, and R. Ortiz. 1992. Seasonal variation of free-living amoebae isolated from the atmosphere of San Luis Potosi, S.L.P., Mexico, abstr. Q-163, p. 362. *In Abstracts of the 92nd General Meeting of the American Society for Microbiology* 1992. American Society for Microbiology, Washington, D.C.
  36. Rowbotham, T. J. 1980. Preliminary report on the pathogenicity of *Legionella pneumophila* for fresh water and soil amoebae. *J. Clin. Pathol.* **33**:1179–1183.
  37. Rowbotham, T. J. 1986. Current views on the relationships between amoebae, legionellae and man. *Isr. J. Med. Sci.* **22**:678–689.
  38. Stein-Streilein, J., and J. Guffee. 1986. *In vivo* treatment of mice and hamsters with antibodies to asialo GM1 increases morbidity and mortality to pulmonary influenza infection. *J. Immunol.* **136**:1436–1441.
  39. St. John, R. C., and P. M. Dorinsky. 1993. Clinical implications of basic research. Immunologic therapy for ARDS, septic shock and multiple organ failure. *Chest* **103**:932–943.
  40. Swanson, M. S., and R. R. Isberg. 1995. Association of *Legionella pneumophila* with the macrophage endoplasmic reticulum. *Infect. Immun.* **63**:3609–3620.
  41. Swanson, M. S., and R. R. Isberg. 1996. Identification of *Legionella pneumophila* mutants that have aberrant intracellular fates. *Infect. Immun.* **64**:2585–2594.
  42. Tyndall, R. L., and E. L. Domingue. 1982. Cocultivation of *Legionella pneumophila* and free-living amoebae. *Appl. Environ. Microbiol.* **44**:954–959.
  43. Wadowsky, R. M., L. J. Butler, M. K. Cook, S. M. Verma, M. A. Paul, B. S. Fields, G. Keleti, J. L. Sykora, and R. B. Yee. 1988. Growth-supporting activity for *Legionella pneumophila* in tap water cultures and implication of Hartmannellid amoebae as growth factors. *Appl. Environ. Microbiol.* **54**:2677–2682.

---

Editor: J. M. Mansfield