

Molecular Variation among Type IV Pilin (*bfpA*) Genes from Diverse Enteropathogenic *Escherichia coli* Strains

T. ERIC BLANK,¹ HAILANG ZHONG,¹ ALISON L. BELL,² THOMAS S. WHITTAM,²
AND MICHAEL S. DONNENBERG^{1*}

Division of Infectious Diseases, Department of Medicine, University of Maryland School of Medicine, Baltimore, Maryland 21201,¹ and Institute of Molecular Evolutionary Genetics and Department of Biology, Pennsylvania State University, University Park, Pennsylvania 16802²

Received 12 June 2000/Returned for modification 7 September 2000/Accepted 13 September 2000

Typical enteropathogenic *Escherichia coli* (EPEC) strains produce bundle-forming pili (BFP), type IVB fimbriae that have been implicated in EPEC virulence, antigenicity, autoaggregation, and localized adherence to epithelial cells (LA). BFP are polymers of bundlin, a pilin protein that is encoded by the *bfpA* gene found on a large EPEC plasmid. Striking sequence variation has previously been observed among type IV pilin genes of other gram-negative bacterial pathogens (e.g., *Pseudomonas* and *Neisseria* spp.). In contrast, the established sequences of *bfpA* genes from two distantly related prototype EPEC strains vary by only a single base pair. To determine whether bundlin sequences vary more extensively, we used PCR to amplify the *bfpA* genes from 19 EPEC strains chosen for their various serotypes and sites and years of isolation. Eight different *bfpA* alleles were identified by sequencing of the PCR products. These alleles can be classified into two major groups. The α group contains three alleles derived from strains carrying O55, O86, O111, O119, O127, or O128 somatic antigens. The β group contains five alleles derived from strains carrying O55, O110, O128ab, O142, or nontypeable antigens. Sequence comparisons show that bundlin has highly conserved and variable regions, with most of the variation occurring in the C-terminal two-thirds of the protein. The results of multilocus enzyme electrophoresis support the hypothesis that *bfpA* sequences have spread horizontally across distantly related clonal lineages. Strains with divergent bundlin sequences express bundlin protein, produce BFP, and carry out autoaggregation and LA. However, four strains lack most or all of these phenotypes despite having an intact *bfpA* gene. These results have important implications for our understanding of bundlin structure, transmission of the *bfp* gene cluster among EPEC strains, and the role of bundlin variation in the evasion of host immune system responses.

Enteropathogenic *Escherichia coli* (EPEC) is one of several pathovars of *E. coli* capable of causing diarrhea (45). While human EPEC infections, which are manifest primarily in infants, were once commonplace in industrialized nations (56), they are now identified primarily in developing countries (10). EPEC strains possess distinct virulence factors not found in most *E. coli* strains (45). Typical EPEC strains produce bundle-forming pili (BFP), long, flexible, rope-like structures composed of intertwining fibers (19). Based on protein sequence analysis and morphology, BFP belong to the type IV group of fimbriae or pili found on a variety of gram-negative bacteria, many of which are human, animal, or plant pathogens (67, 70, 77). BFP have recently been shown to elicit an antibody response in natural infections (37, 38, 53) and a modest response in experimentally infected adults (13). BFP are required for two phenotypes of EPEC that can be studied in vitro and probably play a role in colonization in vivo. The first is autoaggregation, which is the ability of EPEC to reversibly form multicellular clusters in liquid culture (5, 76). The second is localized adherence (LA), which is the capacity of EPEC to form defined multicellular clusters (microcolonies) upon epithelial cells (7, 59) and upon human intestinal tissue (26). Both of these phenomena seem to be manifestations of the ability of BFP to intertwine, creating a fibrous network connecting indi-

vidual bacteria. BFP also appear to adhere directly to epithelial cells (19, 31), although no BFP receptor on epithelial cells has been identified. During the course of tissue culture infection, BFP bundles undergo a transformation to a longer and thicker form that is correlated with dispersal of bacteria from microcolonies (31). In a recent clinical study, volunteers fed mutant EPEC strains lacking BFP developed diarrhea at a much lower frequency and of less severity than did volunteers receiving equivalent doses of an isogenic strain producing BFP (5). Thus, BFP are dynamic fimbriae that contribute to EPEC pathogenesis.

BFP biogenesis is specified by a cluster of 14 *bfp* genes (65, 66) located on large (~50- to 70-MDa) EPEC adherence factor (EAF) plasmids (3, 47, 48). The first gene of this cluster is *bfpA*, which encodes bundlin, a pilin protein that is the only known structural component of BFP. Like other type IV pilins, bundlin is synthesized with a leader peptide removed by a prepilin peptidase, has a hydrophobic region at its mature N terminus, and has two cysteine residues forming a disulfide bond near the C terminus. Bundlin is membrane associated (2, 76). It seems likely that the ultimate function of most of the *bfp* gene products is to remove bundlin subunits from the bacterial membrane and assemble them into pili (2).

Little is known about the structure of bundlin monomers or how they assemble into BFP. However, some information on these topics can be inferred from studies of other type IV pilin. The crystal structures of type IV pilin monomers from *Neisseria gonorrhoeae* and *Pseudomonas aeruginosa* have been solved (15, 25, 52). These pilins are ladle shaped with a long N-terminal α -helical handle and a globular head. Models of the

* Corresponding author. Mailing address: Division of Infectious Diseases, Department of Medicine, University of Maryland School of Medicine, 10 South Pine St., Medical School Teaching Facility 9-00, Baltimore, MD 21201. Phone: (410) 706-7560. Fax: (410) 706-8700. E-mail: mdonnenb@umaryland.edu.

TABLE 1. EPEC strains used in this study

Strain	Serotype	Isolation:			Reference(s)	Source
		Yr	Location	Circumstances		
10	O119:H+	1986	Santiago, Chile	Health center-based study	36, 46	CVD ^a
009-271082	O111:NM	1982	Lima, Peru	Infant cohort study	44	CVD
010-311082	O128:H+	1982	Lima, Peru	Infant cohort study	44	CVD
012-050982	O142:H+	1982	Lima, Peru	Infant cohort study	44	CVD
0659-79	O119:H6	1979	Florida	Infant diarrhea	18, 20, 35, 47	CDC ^b
2309-77	O111ab:H2	1977	California	Infant stool	47	CDC
Beta	O55:H6	1947	Aberdeen, Scotland	Infant diarrhea outbreak	17, 57	R. M. Robins-Browne
C771	O142:H6	Pre-1960	Djakarta, Indonesia	Infant diarrhea	51, 57	R. M. Robins-Browne
CA89-4221	NT ^c :H+	1989	Montreal, Quebec, Canada	Canine diarrhea	4, 14	J. Harel
DIF043256	O111:H2	1986	Morelos, Mexico	Acute infant diarrhea	9	A. Cravioto
E56/54	O128ab:H2	1953	England	Infant diarrhea outbreak	57, 69	R. M. Robins-Browne
E851/71	O142:H6	1971	Glasgow, Scotland	Infant diarrhea outbreak	7, 29, 34, 35, 47	B. Rowe
E990	O86:NM	1950	England	Infant diarrhea outbreak	57, 68	R. M. Robins-Browne
E2348/69	O127:H6	1969	Taunton, England	Infant diarrhea outbreak	7, 11, 34, 35, 47	CVD
HSP19/4	O86:H+	1995	São Paulo, Brazil	Acute infant diarrhea		I. C. A. Scaletsky
MA551/1	O55:NM	1998	São Luiz, Brazil	Acute infant diarrhea		I. C. A. Scaletsky
RN191/1	O111:H+	1996	Natal, Brazil	Acute infant diarrhea		I. C. A. Scaletsky
RN410/1	O119:H+	1998	Natal, Brazil	Pediatric control		I. C. A. Scaletsky
Stoke W	O111ab:NM	1947	Aberdeen, Scotland	Pediatric control	17, 57	R. M. Robins-Browne
Z188-93	O110:H6	1993	Germany	Avian diarrhea	60	J. E. Lohr

^a CVD, Center for Vaccine Development, University of Maryland School of Medicine, Baltimore.

^b CDC, Centers for Disease Control and Prevention, Atlanta, Ga.

^c NT, nontypeable.

way in which pilin monomers pack together into the pilus fiber have been proposed for the gonococcal and *Pseudomonas* pili (16, 25, 52), as well as the toxin-coregulated pili (TCP) of *Vibrio cholerae* (R. Chattopadhyaya and A. C. Ghose, unpublished data; Protein Data Bank accession number 1QQZ).

Extensive sequence variability has been observed in the type IV pilins of *Dichelobacter nodosus*, *Eikenella corrodens*, *Moraxella bovis*, *P. aeruginosa* (reviewed in reference 70), *N. gonorrhoeae* (52), *N. meningitidis* (1), and *Vibrio cholerae* (27, 43, 50, 55). In some of these species, pilin variability is clearly linked to changes in antigenicity. The host immune response may have played a major role in selecting for pilin variants with altered sequences. In the *N. gonorrhoeae* genome, extra copies of partial pilin genes are also present. These normally silent copies can be exchanged with the active copy via recombination, leading to antigenic variation that can aid the gonococcus in evading the host immune response (reviewed in references 41 and 62). A comparison of many gonococcal pilin gene products shows that extensive sequence variation occurs in amino acid residues of the "head" region but hardly any occurs in residues of the "handle" region (52). A hypervariable region of gonococcal pilin occurs between the disulfide-bonded cysteine residues. This region, as well as a disaccharide moiety and the extreme C terminus of the pilin, is predicted to be exposed on the outside of the pilus fiber (16, 52). In *P. aeruginosa* pilins, the disulfide region is also variable in sequence and serves as a binding domain for cell surface receptors (reviewed in reference 23).

In contrast to the considerable variation noted in other type IV pilins, the bundlins encoded by the two known *bfpA* alleles contain only a single amino acid difference (11, 64). Bundlin proteins from diverse EPEC strains might be expected to exhibit more significant sequence variation than has previously been noted. The current study was undertaken to assess the level of sequence diversity in bundlin proteins among EPEC strains and to gain insight into structural and functional constraints on different regions of the molecule.

MATERIALS AND METHODS

EPEC strains. The origins of the EPEC strains examined in this study are listed in Table 1. To analyze the motility of each strain, a loopful of culture was stabbed into a motility agar plate (1% Bacto-Tryptone, 0.5% NaCl, 0.35% agar) and examined after overnight incubation at 37°C. Based on this analysis, strains lacking H serotype information were designated H+ (motile) or NM (nonmotile) (Table 1).

Isolation and sequencing of *bfpA* genes. Partial and complete *bfpA* genes were amplified by PCR using DeepVent DNA polymerase in accordance with the manufacturer's (New England Biolabs) instructions. EPEC colonies were picked from Luria agar plates and boiled for 10 min in ThermoPol buffer (New England Biolabs) containing 2 to 6 mM MgSO₄, DeepVent DNA polymerase (2 U), oligonucleotide primers (0.35 to 1.2 μM), and deoxynucleoside triphosphates (400 μM each) were added to bring the total reaction volume to 100 μl before beginning the PCR. Primer pairs used for PCR were Donne-28 and -29, -362 and -363, -423 and -363, -6 and -382, and -423 and -382 (Table 2). The precise reaction conditions varied with the strain and primers used. The PCR products were examined by agarose gel electrophoresis. If the PCR mixture contained a single DNA species of the expected size, it was purified directly using the Wizard PCR Preps DNA Purification System (Promega). If additional DNA species were present, the desired PCR product was isolated from a preparative agarose gel. Automated DNA sequencing of PCR products was carried out by the staff of The Biopolymer Laboratory (University Of Maryland School of Medicine) using

TABLE 2. Oligonucleotide primers used in this study

Primer	Sequence ^a	Annealing site ^a
Coding strand		
Donne-6	5'- <u>TCTTGGTGCTTGCGTGC</u> -3'	19-208
Donne-28	5'-CGCGGATCCATGGTTTCTAAATCATGAAT-3'	274-294
Donne-362	5'-AGGTCTGTCTTTGATTGAA-3'	309-327
Donne-423	5'-GATTATTCCGTGACCTATT-3'	223-241
Noncoding strand		
Donne-26	5'-GTAAGCGTCAGATAGTAACCAA-3'	638-617
Donne-29	5'-GCGAAGCTTTTACTTTCATAAAATATGTAAC-3'	855-835
Donne-363	5'-CCTGAGTAAACAGGATAC-3'	953-935
Donne-382	5'-TCCTTCGGGTGATTGTGTA-3'	1300-1282
Donne-416	5'-GTTGCGCTCATTACTTCT-3'	443-426

^a The underlined portion of each primer corresponds to the annealing site indicated within the *bfp* cluster nucleotide sequence (EMBL accession number Z68186).

two or more of the primers listed in Table 2. For some strains, the entire double-stranded sequence of the *bfpA* gene could be obtained using only primers Donne-423 and Donne-363.

Sequence analysis. DNASIS (Hitachi Software) was used to align *bfpA* nucleotide sequences from each strain and determine a consensus sequence. Multiple sequence alignment was performed on the *bfpA* gene sequences and the deduced bundlin amino acid sequences using DNASIS and CLUSTAL W (71). Gene trees were constructed with the computer program MEGA (Molecular Evolutionary Genetics Analysis, version 1.0; S. Kumar, K. Tamura, and M. Nei, Institute of Molecular Evolutionary Genetics, The Pennsylvania State University, 1993 [http://www.bio.psu.edu/People/Faculty/Nei/Lab/megaform.txt]). The proportions of polymorphic synonymous (p_S) and nonsynonymous (p_N) sites were calculated by the method of Nei and Gojibori (49). To examine variation in the functional constraints of different parts of the molecule, these statistics were tabulated in a sliding-window analysis of 30 codons along the gene by the program PSWIN. The theoretical three-dimensional structure of bundlin was analyzed and colored using RASMOL (58) (http://www.umass.edu/microbio/rasmol/).

MLEE. Every strain was characterized by its profile of electromorphs for 20 enzymes by multilocus enzyme electrophoresis (MLEE) (63, 78, 79). Genetic relationships between strains were determined based on a matrix of genetic distances between all pairs constructed by comparison of the allelic arrays. A dendrogram was constructed with MEGA using the neighbor-joining algorithm.

Immunoblotting. Overnight cultures of EPEC in LB were diluted 1:100 into 20 ml of Dulbecco's modified Eagle medium–nutrient mixture F-12 (DMEM–F-12) containing 15 mM HEPES buffer and lacking phenol red (Gibco-BRL Life Technologies catalog no. 11039-021). After 6 h of growth at 37°C with shaking, the bacteria were pelleted by centrifugation at $2,500 \times g$ for 5 min and then resuspended in 1.2 ml of cell lysis buffer (20 mM Tris-HCl [pH 8.0], 500 mM NaCl, 0.1 mM EDTA, 0.1% Triton X-100). The optical density at 600 nm of a portion of these samples was measured on a spectrophotometer. Aliquots (75 μ l or less) of these samples were mixed with 25 μ l of 4 \times sodium dodecyl sulfate–polyacrylamide gel electrophoresis loading buffer (2) and then adjusted to a total volume of 100 μ l with additional cell lysis buffer such that the resulting gel samples were derived from roughly equivalent densities of bacteria. The gel samples were boiled for 10 min, and then 5 μ l of each was loaded per lane of a sodium dodecyl sulfate–15% polyacrylamide gel. After electrophoretic separation, the protein in the gel was electrotransferred to a polyvinylidene difluoride membrane. The membrane was blocked with phosphate-buffered saline containing 0.1% Tween 20 plus 5% nonfat dry milk and then incubated sequentially with a polyclonal antibody (1:5,000 dilution) directed against six-His-tagged prebundlin (81), an anti-rabbit horseradish peroxidase conjugate (1:30,000 dilution), and ECL Western blotting detection reagents (Amersham Pharmacia Biotech).

Transmission electron microscopy. To prepare samples for electron microscopy, 1-ml aliquots were removed from EPEC cultures grown in DMEM–F-12 for 5 h and the bacteria were pelleted by brief centrifugation. Most of the medium was decanted, and the bacterial pellet was gently resuspended in the remainder. Aliquots (10 μ l) were spotted onto electron microscopy grids, which were dried for 10 min and then washed, stained, and examined for BFP as described previously (66).

Autoaggregation assay. EPEC strains were cultured overnight at 37°C in LB. The resulting stationary-phase cultures were diluted 1:250 (making appropriate adjustments for optical density at 600 nm) into 20 ml of DMEM–F-12. These cultures were shaken at 250 rpm for 5 h in 50-ml conical polypropylene tubes at 37°C. Autoaggregation was initially gauged by visually inspecting the cultures for bacterial aggregates and sedimentation. For some strains, it was necessary to use light microscopy in order to detect aggregates. Culture aliquots (5 μ l) were examined in hanging-drop slides at $\times 40$ magnification.

LA assay. LA to HeLa epithelial cells was assayed as described previously (12) using the eight-well chamber slide modification.

Nucleotide sequence accession numbers. The nucleotide sequences of the *bfpA* genes isolated from all of the strains listed in Table 1 (except prototype strain E2348/69) have been deposited in GenBank under accession numbers AF304468 through AF304486.

RESULTS

Identification of novel *bfpA* alleles. The goal of this study was to assess the amount of sequence variation among *bfpA* genes found in EPEC strains. To this end, we initially analyzed 13 strains from our laboratory collection (a subset of those listed in Table 1) representing a variety of serotypes and well-documented outbreaks (or sporadic cases) of human diarrhea. These strains were originally isolated from diverse geographical locations between 1947 and 1986. They belong to classical EPEC O serogroups O55, O86, O111, O119, O127, O128, and O142. Three of them (C771, E990, and Stoke W) are type strains for the preparation of O antisera. Canine EPEC strain CA89-4221 and avian EPEC strain Z188-93, which are both

probe positive for the *bfpA* gene, were also examined. PCR was used to amplify the *bfpA* gene from each EPEC strain. The DNA sequences of both strands of the PCR products were determined, and the resulting *bfpA* coding sequences were compared to each other and to the known *bfpA* sequences from prototypic strains E2348/69 and B171 (GenBank accession numbers Z12295 and U27184). Eight different *bfpA* alleles were identified among the 15 strains. Six of these alleles have not been previously described. Each *bfpA* allele specifies a prebundlin protein carrying a variant amino acid at one or more positions (Fig. 1).

Genetic relationships among *bfpA* alleles and correspondence to O serogroups. Sequence comparisons of the *bfpA* alleles show that they fall into two distinct clonal groups, which we term *bfpA* alpha (α) and *bfpA* beta (β) (Fig. 2 and Table 3). The three α alleles show polymorphism at only 7 (1.2%) of 579 nucleotides. The α prebundlins differ at only 6 (3.1%) of 193 amino acid residues. The previously reported $\alpha 1$ allele is found in prototype O127 strain E2348/69 (11), as well as in the single O86 and O55 strains initially examined in the current study. The previously reported $\alpha 2$ allele is carried by prototype O111 strain B171 (64) and each of the other O111 strains examined. The $\alpha 1$ allele varies from $\alpha 2$ at a single nucleotide, resulting in an amino acid substitution. The $\alpha 3$ allele of *bfpA* is found in both of the O119 strains but in only one of the two O128 strains we initially examined. This allele varies from $\alpha 1$ and $\alpha 2$ at six nucleotides located in the 3' half of the gene. Five of these six nucleotide changes specify altered amino acids relative to $\alpha 2$ prebundlin. Three of the sequence changes are identical to those found throughout the β group, suggesting a closer relationship to a common α/β ancestor.

Among the human-derived strains, β alleles were identified in each of the three strains of the O142 serogroup and in one of the two strains of the O128 serogroup. Unique β alleles were also found in the avian and canine EPEC strains. The five *bfpA* β alleles are more diverse in sequence than the α alleles, with the exception of $\beta 4$ and $\beta 5$, which differ from one another by only one nucleotide. At the nucleotide level, β alleles are polymorphic at 51 (8.7%) of 585 nucleotides. The β prebundlins vary at 22 (11.3%) of 195 amino acids. Relative to α alleles, β alleles of *bfpA* carry either a net insertion ($\beta 1/\beta 2/\beta 3$) or a net deletion ($\beta 4/\beta 5$) of one codon directly after codon 110 plus a one-codon insertion between codons 164 and 165 (Fig. 1 and 2). When both α and β sequences are considered, *bfpA* exhibits polymorphism in 94 (16.1%) of 585 nucleotides and prebundlin varies at 36 (18.5%) of 195 amino acids.

Evidence of selection for sequence diversification near the 3' end of the *bfpA* gene. To determine how the level of selective constraint varies along the length of the *bfpA* gene, the proportions of nonsynonymous and synonymous codon changes (p_N and p_S , respectively) were calculated within a 30-codon sliding window (Fig. 3). Nonsynonymous codon changes result in amino acid substitution; synonymous codon changes do not. The difference, $p_N - p_S$, is a measure of the degree of selective constraint. The zero-difference line indicates selectively neutral variation, where p_S and p_N are equal. The more positive the value, the less the contribution of synonymous substitutions and the greater the contribution of replacement substitutions. A positive difference in a particular region suggests that some selective pressure is driving diversification in that region, whereas a negative difference indicates selection against mutations resulting in amino acid variation. The results shown in Fig. 3 suggest that most bundlin is under selective constraint. However, a single region of *bfpA*, located between codons 143 and 174, exhibits a positive value of $p_N - p_S$. This region corresponds to amino acids 130 through 161 of mature bund-




FIG. 1. Variable amino acids are clustered in prebundlin. An alignment of the prebundlin amino acid sequences encoded by the eight *bfpA* alleles described in this study is shown. Note that amino acids 1 to 13 comprise the cleaved leader peptide. The α 1 (ALPHA1.AMI) prebundlin prototype sequence is listed on the top line. In the remaining seven sequences (ALPHA2.AMI through ALPHA3.AMI and BETA1.AMI through BETA5.AMI), invariant amino acids are indicated by dots. Variant amino acids are boxed and indicated by single-letter abbreviations.

lin, which are located between the disulfide-bonded cysteines. An attractive possibility is that this region of bundlin is located on the surface of the BFP filament and that the host immune system has selected for antigenic diversity at this site.

Genetic relationships between EPEC strains. MLEE was carried out to define the overall phylogenetic relationships between selected EPEC strains. A dendrogram resulting from this analysis (Fig. 4) shows that all of the α 2 alleles are found in a closely related cluster of O111 strains. Among the remaining EPEC strains, however, there is a curious lack of correspondence between the deduced clonal pattern of the strains and the pattern of *bfpA* alleles they possess. Alleles of both the α and β types are found among strains in either of the two major clonal EPEC groups, EPEC 1 and EPEC 2 (54, 78). In the EPEC 1 group, the α 1, α 3, β 1, β 3, and β 5 alleles are interspersed among each other in a way that obviously does not correlate with the overall strain lineage. These results strongly suggest that the *bfpA* gene has been recently transferred between EPEC strains through multiple horizontal transfer events.

Examination of *bfpA* alleles in recent EPEC isolates. To determine whether *bfpA* α alleles have remained stable over recent decades, we examined a set of four EAF⁺ EPEC strains of very recent Brazilian origin (Table 1). The *bfpA* alleles derived from these strains were isolated by PCR and sequenced. As expected from their serogroups, the O86, O111, and O119 strains (HSP19/4, RN191/1, and RN410/1) possessed *bfpA* genes of the α 1, α 2, and α 3 varieties, respectively. These results indicate that *bfpA* α sequences have been retained unchanged in current EPEC strains. The O55 strain (MA551/

1), however, did not carry the α 1 allele found in the previously examined O55 EPEC (strain β). Rather, it carried a β 5 allele identical to that carried by O142 strains C771 and E851/71. This unexpected finding provides a further suggestion of the horizontal spread of a particular *bfpA* allele.

Expression and function of BFP variants. Selected EPEC strains were examined to determine whether they produced detectable bundlin and BFP and whether they exhibited the BFP-dependent phenotypes of autoaggregation and LA. Some of these characteristics have been previously reported for a subset of these strains (see references in Table 1). Prototype EPEC strain E2348/69 served as a positive control for all of our studies. To identify bundlin, a polyclonal antiserum was used to probe whole-cell extracts from each of the EPEC strains (Fig. 5 and Table 3). Bundlin protein was readily detected in extracts from 10 out of 12 strains tested carrying α alleles of *bfpA* and 5 of 7 strains carrying β alleles. Bundlin was not identified in strain 009-271082 or 2309-77 carrying the α 2 allele or in strain E56/54 or C771 carrying the β 4 or β 5 allele. Particular β bundlins migrated more slowly (β 1 and β 2) or more quickly (β 5) than did α bundlins (Fig. 5 and data not shown). The relative migration correlates well with the lengths (180 amino acids for α 1 and β 5 and 182 amino acids for β 1 and β 2) and molecular masses (19,986 Da for β 5, 20,269 Da for α 1, 20,306 Da for β 2, and 20,328 Da for β 1) predicted for these bundlin types. Such a migration anomaly has been noted for bundlin from other strains in previous studies (18, 20, 38).

Ten of the strains were examined by electron microscopy for the presence of BFP fimbriae. Fimbrial bundles were readily detected in six strains (Table 3). No BFP was seen in samples




FIG. 2. Eight *bfpA* alleles cluster into two major groups. To the left is a phylogenetic tree for the *bfpA* gene from 17 EPEC strains constructed by the neighbor-joining algorithm based on the gamma distance with $\alpha = 2$. Branch lengths in terms of nucleotide substitutions per 100 sites are given above the major branches. Allele designations are listed for each strain. To the right is a graph of the locations of polymorphic nucleotide sites, which are marked as vertical lines that indicate nucleotides differing from those of the $\alpha 2$ allele at each position. Gaps indicate codons absent from specific alleles.

of the remaining four strains, which had also failed to express detectable bundlin. Noticeably more BFP was produced by $\alpha 1$ strains E990 and E2348/69 than the remaining strains. All of the strains were tested for LA and autoaggregation (Table 3). The $\alpha 2$ allele strains that failed to produce both bundlin and BFP (009-271082 and 2309-77) exhibited neither autoaggregation nor LA. The β allele strains E56/54 and C771, which failed to produce both bundlin and BFP, also did not exhibit autoaggregation. However, they did exhibit LA, as has been previously noted (57). All of the remaining strains exhibited both phenotypes. In the LA assay, no conspicuous differences were noted in the size, pattern, or relative numbers of adherent clusters produced by strains carrying different *bfpA* alleles. While some strains consistently produced aggregates in the autoaggregation assay that could be readily seen by the naked eye, the aggregates of other strains (e.g., E851/71) could be detected only by microscopy. To summarize, most strains expressed BFP that had normal structure and function regardless of the *bfpA* allele they carried. Four strains, however, were defective for BFP expression and function in most or all of the assays. In each of these cases, other strains carrying an identical or closely related *bfpA* allele produced functional BFP, suggesting that the allele sequence itself is not the cause of the defect.

DISCUSSION

In this study, we have identified eight alleles of the *bfpA* gene by surveying 19 EPEC strains of diverse origins. Previous studies using prototype EPEC strains E2348/69 and B171 had identified two of these alleles, $\alpha 1$ and $\alpha 2$, that differ by only a single nucleotide (11, 64). The eight *bfpA* alleles fall into two distinct

TABLE 3. Expression of BFP-related phenotypes by EPEC strains

<i>bfpA</i> allele	Strain	Serotype	Bundlin ^a	BFP ^b	LA ^c	Autoaggregation ^d
$\alpha 1$	Beta	O55:H6	+	ND ^e	+	+
$\alpha 1$	E990	O86:NM	+	+	+	+
$\alpha 1$	E2348/69	O127:H6	+	+	+	+
$\alpha 1$	HSP19/4	O86:H+	+	ND	+	+
$\alpha 2$	009-271082	O111:NM	-	-	-	-
$\alpha 2$	2309-77	O111ab:H2	-	-	-	-
$\alpha 2$	DIF043256	O111:H2	+	ND	+	+
$\alpha 2$	Stoke W	O111ab:NM	+	ND	+	+
$\alpha 2$	RN191/1	O111:H+	+	ND	+	+
$\alpha 3$	10	O119:H+	+	ND	+	+
$\alpha 3$	0659-79	O119:H6	+	ND	+	+
$\alpha 3$	010-311082	O128:H+	+	+	+	+
$\alpha 3$	RN410/1	O119:H+	+	ND	+	+
$\beta 1$	012-050982	O142:H+	+	+	+	+
$\beta 2$	Z188-93	O110:H6	+	ND	+	+
$\beta 3$	CA89-4221	NT ^f :H+	+	+	+	+
$\beta 4$	E56/54	O128ab:H2	-	-	+	-
$\beta 5$	C771	O142:H6	-	-	+	-
$\beta 5$	E851/71	O142:H6	+	+	+	+
$\beta 5$	MA551/1	O55:NM	+	ND	+	+

^a Antigenic detection of bundlin protein by immunoblotting. +, presence of bundlin; -, absence of bundlin.

^b Visual detection of BFP by electron microscopy. +, pili present; -, pili absent.

^c LA of EPEC to cultured epithelial cells. +, presence of LA; -, absence of LA.

^d Autoaggregation of EPEC in tissue culture medium. +, presence of autoaggregation; -, absence of autoaggregation.

^e ND, not determined.

^f NT, nontypeable.




FIG. 3. Evidence for diversifying selection near the 3' end of the *bfpA* gene. A sliding-window plot of the proportions of synonymous and nonsynonymous sites (p_s and p_n , respectively) that have mutated among eight *bfpA* alleles is shown. The difference ($p_n - p_s$) is a measure of the level of selective constraint on various parts of the molecule. Note that $p_n - p_s > 0$ for codons 143 to 174 (indicated by the arrow), a pattern consistent with the effect of diversifying selection.

groups based upon sequence analysis. By analogy to the intimins, we call these groups α and β . The α group, isolated from 12 of the EPEC strains examined here plus prototype strains B171 and E2348/69, contains three different alleles




FIG. 4. The alleles of *bfpA* identified in EPEC strains do not correlate precisely with the overall clonal lineage. Shown is a dendrogram of genetic relationships of EPEC strains from this study and representative DEC strains (79). The genetic distance was estimated in terms of electrophoretically detectable codon differences per enzyme locus for 20 enzymes. Previously identified DEC clusters (EPEC 1 and 2 and EHEC 1 and 2) are indicated (54, 78). Strain serotypes are in parentheses. Alleles of *bfpA* identified in specific EPEC strains are displayed in large, shaded type.







FIG. 5. Immunoblot of bundlin protein from EPEC whole-cell extracts. The arrow indicates the position of bundlin. The extracts are from the following strains: lane 1, E2348/69; lane 2, 0659-79; lane 3, E851/71; lane 4, 2309-77; lane 5, 10; lane 6, 012-050982; lane 7, 009-271082; lane 8, 010-311082; lane 9, DIF043256; lane 10, β ; lane 11, Stoke W; lane 12, E56/54; lane 13, C771; lane 14, E990. The remaining strains exhibited detectable bundlin but are not shown here.

showing little sequence variation. The β group, represented by seven strains, exhibits more significant sequence variation, both internally and when compared to α alleles. We note that additional alleles of *bfpA* have recently been deposited in sequence databases. Bortolini and colleagues have described a nonfunctional *bfpA* allele (GenBank accession number AF119170) that is prematurely truncated by an insertion sequence and is found in certain clonally related O128ab:H2 and O119:H2 EPEC strains (6). We did not identify this allele in our study; however, the residual 5' *bfpA* sequence of this allele is identical to that of the β_3 allele, with the exception of a single nucleotide. Incomplete *bfpA* sequences can be found in GenBank records with accession numbers AF233895 through AF233899 (S. Subpasu, M. Yamasaki, J. Yatsuyanagi, O.-A. Ratchatrachenchai, and K. Ito, unpublished data). One of these sequences appears to be from an α_1 allele, but the remaining four differ from our α_3 , β_1 , and β_4/β_5 alleles at 1 to 10 nucleotide positions. These sequences support our division of *bfpA* alleles into two groups and attest that there is further sequence diversity to be identified in the *bfpA* gene. The existence of multiple *bfpA* alleles has important implications for the study of BFP structure and function, for our understanding of the evolution of EPEC as a pathogen, and for EPEC diagnosis and vaccine development as detailed below.

Implications for the structure of bundlin and BFP. To better understand the structural significance of amino acid polymorphism in bundlin, we carried out a graphical analysis of variability in the sequence of the protein (Fig. 6A). We also mapped the variable residues onto a theoretical three-dimensional (3D) structure model of α_1 bundlin (Fig. 6B and C) that has been recently deposited in the Protein Data Bank (<http://www.rcsb.org/pdb>) under accession number 1QT2 (R. Chatopadhyaya and A. C. Ghose, unpublished data). The variable amino acids of bundlin are generally clustered, within both the linear sequence and the theoretical 3D structure. Variable residues which group together in the 3D structure are sometimes derived from different segments of the linear bundlin sequence (Fig. 6). Most of the highly variable amino acids are located in the C-terminal third of bundlin. Many of the variable residues are located on the pilin surface in the theoretical 3D model. Variable residues 58, 61, 63 and 64, 79, 81, and 92 (numbered in accordance with the 180-residue mature α_1 bundlin prototype) occur along one side of the pilin head (shown in green in Fig. 6B and C). Variable residues 33, 95, 97 and 98, 100 and 101, 136 to 142, and 163 (shown in red) are located in a cleft on the lower convex face of the pilin head. Residues 110, 149 to 151, 154 and 155, 165, and 167 to 169 (shown in blue) are found on one side of the upper convex face of the pilin head. Residues 145 and 146 and residue 174



(shown in violet) are located on the flattened reverse face of the pilin. One possibility is that these variable positions define regions of molecular interactions either between pilin monomers or between BFP filaments. A more appealing possibility is that these variable positions define regions on the surface of BFP that are directly exposed to the host immune system. Interestingly, other regions of bundlin show little or no variation. These include the cleaved leader peptide, the α -helical handle of the pilin, and five beta strands that probably serve to create the structure of the pilin head. Such regions may lack variation due to constraints inherent in critical structural domains.

Like other type IV pilins, bundlin contains two disulfide-bonded cysteine residues near its C terminus. These residues are completely conserved in all bundlin variants, consistent with the previous finding that replacement of either cysteine with serine greatly reduces bundlin stability (81). Replacement of the entire disulfide region of bundlin with the corresponding region of the *V. cholerae* TcpA pilin also leads to instability of the hybrid pilin (40). Studies of multiple type IV pili have shown the importance of variation in the disulfide region. The hypervariable disulfide region of the gonococcal pilin participates in antigenic variation, while that of the *P. aeruginosa* pilin is involved in host cell interaction. A detailed mutational analysis of the TcpA pilin indicates that an intact disulfide region is critical for both proper structure and function of TCP (30). Based on these findings and on the 3D model of Chattopadhyaya and Ghose, it was suggested that this region is surface exposed on TCP and defines surfaces of interaction both between pilin monomers and between individual pilus fibers within a bundle. Interestingly, there is minimal variation in a subregion of ~20 to 25 residues surrounding the more N-terminal cysteine of both BfpA (Fig. 6A) and TcpA (43), corresponding in part to the structural domain defined by Kirn and colleagues (30). Furthermore, there is evidence of constraint against nonsynonymous substitutions in this subregion (Fig. 3), indicating selection against diversity. In contrast, much of the remaining disulfide region is highly variable in both pilins. In fact, our analysis (Fig. 3) suggests that it is under positive selection for diversification. Thus, the disulfide region of type IV pilins may be caught in a struggle to maintain structural integrity, as well as pilin-pilin and filament-filament interactions, while at the same time presenting a changing antigenic face to the host immune system.

Implications for the role of BFP in the evolution of EPEC as a pathogen. EPEC strains are defined as diarrheagenic *E. coli* (DEC) strains producing attaching and effacing histopathology but lacking Shiga-like toxins (28). Typical EPEC strains also carry the EAF plasmid and adhere in a BFP-dependent localized fashion to cultured epithelial cells. The importance of the EAF plasmid and plasmid-encoded BFP in pathogenicity has been demonstrated in volunteer studies (5, 35). Further support is provided by some case-control studies, where EAF⁺, but not EAF⁻, strains have been significantly associated with diarrhea (reviewed in reference 28). As bundlin is a virulence factor and a potentially important antigen of EPEC, it is useful to understand whether *bfpA* alleles are capable of being mo-

bilized between different clonal groups of EPEC strains and how this might take place.

The majority of EPEC strains form two clonal groups, EPEC 1 and EPEC 2 (54, 78), with EPEC 1 exhibiting greater strain diversity. The low G+C content of the 11.5-kb *bfp* gene cluster on the EAF plasmid suggests that it has been derived from a non-*E. coli* bacterium (72). The 14 *bfp* cluster genes from EPEC 1 strain E2348/69 (66) and EPEC 2 strain B171 (65) are more similar to each other in nucleotide sequence than would be expected from the distance between their clonal frames, based on MLEE analysis. Therefore, Whittam and McGraw (78) hypothesized that the EAF plasmid carrying the *bfp* cluster was acquired independently by EPEC 1 and then by EPEC 2 after the divergence of these two groups. The results of the current study indicate that the evolutionary picture is more complex. There has been a radiation of bundlin alleles into at least two major types, α and β , both of which appear to have an ancient origin. In addition, *bfpA* alleles appear to have been exchanged multiple times among divergent EPEC strains, presumably as a result of conjugation (see below). Domesticated animals are a potential reservoir of β alleles, as EPEC strains isolated from a pet dog and a pet bird were both found to carry unique β alleles in this study. The complicated intermingling of both α and β alleles of *bfpA* in EPEC 1 (Fig. 4) suggests that, in addition to mutation, *bfpA* sequences have diverged by multiple recombination, horizontal transfer, and positive-selection events. At least two transfers of *bfpA* into the EPEC 2 group have also taken place.

Since the *bfpA* gene is located on the EAF plasmid, the most likely mechanism for its movement between EPEC strains is plasmid transfer by conjugation. The available evidence seems to indicate that EAF plasmids from some EPEC strains are capable of being mobilized for conjugation while others are not (24, 32, 39). The EAF plasmid of O111 strain B171, which has been completely sequenced, appears to lack plasmid transfer genes (72). It does, however, contain many remnants of insertion elements, suggesting an alternative method of *bfp* cluster transfer via a transposon. The lack of transfer genes on the O111 EAF plasmid may explain why the α 2 allele of *bfpA* was identified only among a closely related cluster of O111 strains.

An unexpected finding of this study was the identification of four strains that contained an intact *bfpA* gene yet did not produce detectable bundlin or BFP or exhibit autoaggregation. Two of the strains (E56/54 and C771) still exhibited LA, suggesting that they either produce sufficient amounts of BFP under the conditions of the adherence assay or that they produce an alternative adherence factor. In such strains, *bfpA* may be regulated differently than in prototype strains. Alternatively, expression of bundlin may have been abrogated during the course of the original infection or subsequently during multiple passages through various laboratories. It is possible that BFP is not strictly required for adherence to cells and infection of the host by some EPEC strains (6, 26).

Implications for EPEC diagnosis. The presence of multiple *bfpA* alleles in nature requires a re-evaluation of the methods used to identify typical EPEC strains. Molecular diagnosis of *bfpA*⁺ EPEC has previously been carried out with hybridiza-

FIG. 6. Variable amino acids may be clustered on the surface of bundlin. (A) Graphical analysis of bundlin variation. The bars represent amino acid residues that vary among the eight bundlin types identified in this study. The heights of the bars are proportional to the numbers of alternate residues per site. The positions of conserved cysteine residues are indicated below the x axis. (B and C) Theoretical 3D structure of bundlin (Chattopadhyaya and Ghose, unpublished). Colored amino acid residues are those that vary between bundlin types. These residues have been assigned to four color groups (green, red, blue, and violet) in accordance with their relative locations in the molecule. These colors correspond to those displayed in the graph (A). The disulfide-bonded cysteines are yellow. Panel B shows the convex face of bundlin that is likely to be facing outward on the surface of the BFP filament. Panel C shows the flattened face on the opposite side of the molecule that is likely to be facing toward the center of the filament, based on existing models of gonococcal and *V. cholerae* pili.

tion probes consisting of restriction digest fragments, PCR products, or labeled oligonucleotides (4, 18, 42, 45, 64). PCR amplification has also been described for the identification of *bfpA* from *E. coli* (22, 74, 75, 80). The current results indicate that oligonucleotides for *bfpA* identification should be designed such that the annealing regions represent sequences conserved in all known *bfpA* alleles. Use of a poorly conserved oligonucleotide sequence might result in the incorrect designation of bonafide *bfpA*⁺ strains as BFP negative. Results from the functional studies we conducted, along with the findings of others (38), show the limitations of using probes or PCR for diagnosis—some strains carrying an intact *bfpA* gene may not actually produce readily detectable levels of bundlin or BFP. Therefore, accurate designation of an *E. coli* strain as BFP positive ultimately requires one or more tests of BFP expression or function. Our results have shown for the first time that diverse EPEC strains exhibit autoaggregation, a phenotype that is BFP dependent. While the autoaggregation assay is not yet in common use, it appears to be an acceptable and facile substitute for LA in those laboratories lacking tissue culture facilities but having a simple microscope.

Antigenic variation of bundlin and implications for vaccine development. BFP has been proposed as a potential component of an EPEC vaccine (33, 61). This concept is supported by the detection of antibundlin antibodies in children with natural EPEC infections (38) and in healthy children and mothers living in areas where EPEC is endemic (37, 53). Further support is provided by the finding that antisera raised against BFP blocks LA (19). Studies of EPEC infections in adult volunteers have demonstrated a correlation between the level of antibundlin antibodies and the propensity for loss of the plasmid encoding BFP (13). Since the plasmid-cured strain is less pathogenic than the plasmid-containing strain (35), this association establishes a link between antibundlin antibodies and virulence. The results of the current study have a direct bearing on considerations for vaccine formulations. Strains of serogroups O55, O111, and O119 have the highest prevalence in areas where EPEC is endemic (8, 21, 36, 73). Our results, which demonstrate that O55 strains, at least, may possess either α or β *bfpA* alleles, suggest that successful vaccines may need to include both classes of antigen. Further studies are needed to test the hypotheses that immune responses discriminate between these classes of bundlin proteins and that the immune response is a significant factor driving the evolution of *bfpA* allelic variants of EPEC.

In summary, our analysis of 19 EPEC strains indicates that there is heretofore unappreciated allelic diversity in the *bfpA* gene. Although some strains with intact *bfpA* alleles failed to produce BFP, most of the strains produced BFP having readily detectable expression and function. The *bfpA* gene contains both variable and invariable regions, whose locations are likely to correlate with their role in bundlin and BFP filament structure. In particular, the most highly variable stretch of bundlin is located in a region of the molecule that is predicted to be surface exposed and under diversifying selection from a host immune response. This study additionally establishes evidence for the horizontal transfer of *bfpA* across diverse EPEC lineages, which likely contributed to the evolution of this important pathogen.

ACKNOWLEDGMENTS

We are grateful to each of the researchers who previously provided EPEC strains that were used in this study, including Alejandro Cravioto, José Harel, J. E. Lohr, Roy Robins-Browne, Bernard Rowe, Isabel Scaletsky, and Nancy Strockbine. We thank Jorge Girón for critical review of the manuscript.

This investigation was supported by a Public Health Service grant (AI37606) to M.S.D. and a National Research Service Award postdoctoral training fellowship (AI10191) to T.E.B.

REFERENCES

- Aho, E. L., J. W. Botten, R. J. Hall, M. K. Larson, and J. K. Ness. 1997. Characterization of a class II pilin expression locus from *Neisseria meningitidis*: evidence for increased diversity among pilin genes in pathogenic *Neisseria* species. *Infect. Immun.* **65**:2613–2620.
- Anantha, R. P., K. D. Stone, and M. S. Donnenberg. 2000. Effects of *bfp* mutations on biogenesis of functional enteropathogenic *Escherichia coli* type IV pili. *J. Bacteriol.* **182**:2498–2506.
- Baldini, M. M., J. B. Kaper, M. M. Levine, D. C. Candy, and H. W. Moon. 1983. Plasmid-mediated adhesion in enteropathogenic *Escherichia coli*. *J. Pediatr. Gastroenterol. Nutr.* **2**:534–538.
- Beaudry, M., C. Zhu, J. M. Fairbrother, and J. Harel. 1996. Genotypic and phenotypic characterization of *Escherichia coli* isolates from dogs manifesting attaching and effacing lesions. *J. Clin. Microbiol.* **34**:144–148.
- Bieber, D., S. W. Ramer, C. Y. Wu, W. J. Murray, T. Tobe, R. Fernandez, and G. K. Schoolnik. 1998. Type IV pili, transient bacterial aggregates, and virulence of enteropathogenic *Escherichia coli*. *Science* **280**:2114–2118.
- Bortolini, M. R., L. R. Trabulsi, R. Keller, G. Frankel, and R. Sperandio. 1999. Lack of expression of bundle-forming pili in some clinical isolates of enteropathogenic *Escherichia coli* (EPEC) is due to a conserved large deletion in the *bfp* operon. *FEMS Microbiol. Lett.* **179**:169–174.
- Cravioto, A., R. J. Gross, S. M. Scotland, and B. Rowe. 1979. An adhesive factor found in strains of *Escherichia coli* belonging to the traditional infantile enteropathogenic serotypes. *Curr. Microbiol.* **3**:95–99.
- Cravioto, A., R. E. Reyes, F. Trujillo, F. Uribe, A. Navarro, J. M. De La Roca, J. M. Hernández, G. Pérez, and V. Vázquez. 1990. Risk of diarrhea during the first year of life associated with initial and subsequent colonization by specific enteropathogens. *Am. J. Epidemiol.* **131**:886–904.
- Cravioto, A., A. Tello, A. Navarro, J. Ruiz, H. Villafán, F. Uribe, and C. Eslava. 1991. Association of *Escherichia coli* HEp-2 adherence patterns with type and duration of diarrhoea. *Lancet* **337**:262–264.
- Donnenberg, M. S. 1995. Enteropathogenic *Escherichia coli*, p. 709–726. In M. J. Blaser, P. D. Smith, J. I. Ravdin, H. B. Greenberg, and R. L. Guerrant (ed.), *Infections of the gastrointestinal tract*. Raven Press, Ltd., New York, N.Y.
- Donnenberg, M. S., J. A. Girón, J. P. Nataro, and J. B. Kaper. 1992. A plasmid-encoded type IV fimbrial gene of enteropathogenic *Escherichia coli* associated with localized adherence. *Mol. Microbiol.* **6**:3427–3437.
- Donnenberg, M. S., and J. P. Nataro. 1995. Methods for studying adhesion of diarrheagenic *Escherichia coli*. *Methods Enzymol.* **253**:324–336.
- Donnenberg, M. S., C. O. Tacket, G. Losonsky, G. Frankel, J. P. Nataro, G. Dougan, and M. M. Levine. 1998. Effect of prior experimental human enteropathogenic *Escherichia coli* infection on illness following homologous and heterologous rechallenge. *Infect. Immun.* **66**:52–58.
- Drolet, R., J. M. Fairbrother, J. Harel, and P. Hélie. 1994. Attaching and effacing and enterotoxigenic *Escherichia coli* associated with enteric colibacillosis in the dog. *Can. J. Vet. Res.* **58**:87–92.
- Forest, K. T., S. A. Dunham, M. Koomey, and J. A. Tainer. 1999. Crystallographic structure reveals phosphorylated pilin from *Neisseria*: phosphoserine sites modify type IV pilus surface chemistry and fibre morphology. *Mol. Microbiol.* **31**:743–752.
- Forest, K. T., and J. A. Tainer. 1997. Type-4 pilus-structure: outside to inside and top to bottom—a minireview. *Gene* **192**:165–169.
- Giles, C., G. Sangster, and J. Smith. 1949. Epidemic gastroenteritis of infants in Aberdeen during 1947. *Arch. Dis. Child.* **24**:45–53.
- Girón, J. A., M. S. Donnenberg, W. C. Martin, K. G. Jarvis, and J. B. Kaper. 1993. Distribution of the bundle-forming pilus structural gene (*bfpA*) among enteropathogenic *Escherichia coli*. *J. Infect. Dis.* **168**:1037–1041.
- Girón, J. A., A. S. Y. Ho, and G. K. Schoolnik. 1991. An inducible bundle-forming pilus of enteropathogenic *Escherichia coli*. *Science* **254**:710–713.
- Girón, J. A., F. Qadri, T. Azim, K. J. Jarvis, J. B. Kaper, and M. J. Albert. 1995. Monoclonal antibodies specific for the bundle-forming pilus of enteropathogenic *Escherichia coli*. *Infect. Immun.* **63**:4949–4952.
- Gomes, T. A. T., M. A. M. Vieira, I. K. Wachsmuth, P. A. Blake, and L. R. Trabulsi. 1989. Serotype-specific prevalence of *Escherichia coli* strains with EPEC adherence factor genes in infants with and without diarrhea in São Paulo, Brazil. *J. Infect. Dis.* **160**:131–135.
- Gunzburg, S. T., N. G. Tornieporth, and L. W. Riley. 1995. Identification of enteropathogenic *Escherichia coli* by PCR-based detection of the bundle-forming pilus gene. *J. Clin. Microbiol.* **33**:1375–1377.
- Hahn, H. P. 1997. The type-4 pilus is the major virulence-associated adhesin of *Pseudomonas aeruginosa*—a review. *Gene* **192**:99–108.
- Hales, B. A., C. A. Hart, R. M. Batt, and J. R. Saunders. 1992. The large plasmids found in enterohemorrhagic and enteropathogenic *Escherichia coli* constitute a related series of transfer-defective Inc F-IIA replicons. *Plasmid* **28**:183–193.
- Hazes, B., P. A. Sastry, K. Hayakawa, R. J. Read, and R. T. Irvin. 2000. Crystal structure of *Pseudomonas aeruginosa* PAK pilin suggests a main-

- chain-dominated mode of receptor binding. *J. Mol. Biol.* **299**:1005–1017.
26. Hicks, S., G. Frankel, J. B. Kaper, G. Dougan, and A. D. Phillips. 1998. Role of intimin and bundle-forming pili in enteropathogenic *Escherichia coli* adhesion to pediatric intestinal tissue in vitro. *Infect. Immun.* **66**:1570–1578.
 27. Iredell, J. R., and P. A. Manning. 1994. Biotype-specific *tcpA* genes in *Vibrio cholerae*. *FEMS Microbiol. Lett.* **121**:47–54.
 28. Kaper, J. B. 1996. Defining EPEC. *Rev. Microbiol. São Paulo* **27**(Suppl. 1):130–133.
 29. Kennedy, D. H., G. H. Walker, R. J. Fallon, J. F. Boyd, R. J. Gross, and B. Rowe. 1973. An outbreak of infantile gastroenteritis due to *E. coli* 0142. *J. Clin. Pathol.* **26**:731–737.
 30. Kirn, T. J., M. J. Lafferty, C. M. Sandoe, and R. K. Taylor. 2000. Delineation of pilin domains required for bacterial association into microcolonies and intestinal colonization by *Vibrio cholerae*. *Mol. Microbiol.* **35**:896–910.
 31. Knutton, S., R. K. Shaw, R. P. Anantha, M. S. Donnenberg, and A. A. Zorngani. 1999. The type IV bundle-forming pilus of enteropathogenic *Escherichia coli* undergoes dramatic alterations in structure associated with bacterial adherence, aggregation and dispersal. *Mol. Microbiol.* **33**:499–509.
 32. Laporta, M. Z., M. L. M. Silva, I. C. A. Scaletsky, and L. R. Trabulsi. 1986. Plasmids coding for drug resistance and localized adherence to HeLa cells in enteropathogenic *Escherichia coli* O55:H– and O55:H6. *Infect. Immun.* **51**:715–717.
 33. Levine, M. M. 1996. Vaccines against enteropathogenic *Escherichia coli*. *Rev. Microbiol. São Paulo* **27**(Suppl. 1):126–129.
 34. Levine, M. M., E. J. Bergquist, D. R. Nalin, D. H. Waterman, R. B. Hornick, C. R. Young, S. Sotman, and B. Rowe. 1978. *Escherichia coli* strains that cause diarrhoea but do not produce heat-labile or heat-stable enterotoxins and are non-invasive. *Lancet* **i**:1119–1122.
 35. Levine, M. M., J. P. Nataro, H. Karch, M. M. Baldini, J. B. Kaper, R. E. Black, M. L. Clements, and A. D. O'Brien. 1985. The diarrheal response of humans to some classic serotypes of enteropathogenic *Escherichia coli* is dependent on a plasmid encoding an enteroadhesiveness factor. *J. Infect. Dis.* **152**:550–559.
 36. Levine, M. M., V. Prado, R. Robins-Browne, H. Lior, J. B. Kaper, S. L. Moseley, K. Gicquelais, J. P. Nataro, P. Vial, and B. Tall. 1988. Use of DNA probes and HEp-2 cell adherence assay to detect diarrheagenic *Escherichia coli*. *J. Infect. Dis.* **158**:224–228.
 37. Loureiro, I., G. Frankel, J. Adu-Bobie, G. Dougan, L. R. Trabulsi, and M. M. Carneiro-Sampaio. 1998. Human colostrum contains IgA antibodies reactive to enteropathogenic *Escherichia coli* virulence-associated proteins: intimin, BfpA, EspA, and EspB. *J. Pediatr. Gastroenterol. Nutr.* **27**:166–171.
 38. Martinez, M. B., C. R. Taddei, A. Ruiz-Tagle, L. R. Trabulsi, and J. A. Girón. 1999. Antibody response of children with enteropathogenic *Escherichia coli* infection to the bundle-forming pilus and locus of enterocyte effacement-encoded virulence determinants. *J. Infect. Dis.* **179**:269–274.
 39. McConnell, M. M., H. Chart, S. M. Scotland, H. R. Smith, G. A. Willshaw, and B. Rowe. 1989. Properties of adherence factor plasmids of enteropathogenic *Escherichia coli* and the effect of host strain on expression of adherence to HEp-2 cells. *J. Gen. Microbiol.* **135**:1123–1134.
 40. McNamara, B. P., and M. S. Donnenberg. 2000. Evidence for specificity in type 4 pilus biogenesis by enteropathogenic *Escherichia coli*. *Microbiology* **146**:719–729.
 41. Meyer, T. F., C. P. Gibbs, and R. Haas. 1990. Variation and control of protein expression in *Neisseria*. *Annu. Rev. Microbiol.* **44**:451–477.
 42. Nagayama, K., Z. Bi, T. Oguchi, Y. Takarada, S. Shibata, and T. Honda. 1996. Use of an alkaline phosphatase-conjugated oligonucleotide probe for the gene encoding the bundle-forming pilus of enteropathogenic *Escherichia coli*. *J. Clin. Microbiol.* **34**:2819–2821.
 43. Nandi, B., R. K. Nandy, A. C. Vicente, and A. C. Ghose. 2000. Molecular characterization of a new variant of toxin-coregulated pilus protein (TcpA) in a toxigenic non-O1/non-O139 strain of *Vibrio cholerae*. *Infect. Immun.* **68**:948–952.
 44. Nataro, J. P., M. M. Baldini, J. B. Kaper, R. E. Black, N. Bravo, and M. M. Levine. 1985. Detection of an adherence factor of enteropathogenic *Escherichia coli* with a DNA probe. *J. Infect. Dis.* **152**:560–565.
 45. Nataro, J. P., and J. B. Kaper. 1998. Diarrheagenic *Escherichia coli*. *Clin. Microbiol. Rev.* **11**:142–201.
 46. Nataro, J. P., J. B. Kaper, R. Robins-Browne, V. Prado, P. Vial, and M. M. Levine. 1987. Patterns of adherence of diarrheagenic *Escherichia coli* to HEp-2 cells. *Pediatr. Infect. Dis. J.* **6**:829–831.
 47. Nataro, J. P., K. O. Maher, P. Mackie, and J. B. Kaper. 1987. Characterization of plasmids encoding the adherence factor of enteropathogenic *Escherichia coli*. *Infect. Immun.* **55**:2370–2377.
 48. Nataro, J. P., I. C. A. Scaletsky, J. B. Kaper, M. M. Levine, and L. R. Trabulsi. 1985. Plasmid-mediated factors conferring diffuse and localized adherence of enteropathogenic *Escherichia coli*. *Infect. Immun.* **48**:378–383.
 49. Nei, M., and T. Gojobori. 1986. Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Mol. Biol. Evol.* **3**:418–426.
 50. Novais, R. C., A. Coelho, C. A. Salles, and A. C. Vicente. 1999. Toxin-co-regulated pilus cluster in non-O1, non-toxicogenic *Vibrio cholerae*: evidence of a third allele of pilin gene. *FEMS Microbiol. Lett.* **171**:49–55.
 51. Ørskov, F., I. Ørskov, T. A. Rees, and K. Sahab. 1960. Two new *E. coli* O-antigens: O141 and O142 and two new coli K-antigens: K85 and K86. *Acta Pathol. Microbiol. Scand.* **48**:48–50.
 52. Parge, H. E., K. T. Forest, M. J. Hickey, D. A. Christensen, E. D. Getzoff, and J. A. Tainer. 1995. Structure of the fibre-forming protein pilin at 2.6 Å resolution. *Nature* **378**:32–38.
 53. Parissi-Crivelli, A., J. M. Parissi-Crivelli, and J. A. Girón. 2000. Recognition of enteropathogenic *Escherichia coli* virulence determinants by human colostrum and serum antibodies. *J. Clin. Microbiol.* **38**:2696–2700.
 54. Reid, S. D., C. J. Herbelin, A. C. Bumbaugh, R. K. Selander, and T. S. Whittam. 2000. Parallel evolution of virulence in pathogenic *Escherichia coli*. *Nature* **406**:64–67.
 55. Rhine, J. A., and R. K. Taylor. 1994. TcpA pilin sequences and colonization requirements for O1 and O139 *Vibrio cholerae*. *Mol. Microbiol.* **13**:1013–1020.
 56. Robins-Browne, R. M. 1987. Traditional enteropathogenic *Escherichia coli* of infantile diarrhea. *Rev. Infect. Dis.* **9**:28–53.
 57. Robins-Browne, R. M., W. C. Yam, L. E. O'Gorman, and K. A. Bettelheim. 1993. Examination of archetypal strains of enteropathogenic *Escherichia coli* for properties associated with bacterial virulence. *J. Med. Microbiol.* **38**:222–226.
 58. Sayle, R. A., and E. J. Milner-White. 1995. RASMOL: biomolecular graphics for all. *Trends Biochem. Sci.* **20**:374.
 59. Scaletsky, I. C. A., M. L. M. Silva, and L. R. Trabulsi. 1984. Distinctive patterns of adherence of enteropathogenic *Escherichia coli* to HeLa cells. *Infect. Immun.* **45**:534–536.
 60. Schremmer, C., J. E. Lohr, U. Wasthuber, J. Kösters, K. Ravelshofer, H. Steinrück, and L. H. Wieler. 1999. Enteropathogenic *Escherichia coli* in *Psittaciformes*. *Avian Pathol.* **28**:349–354.
 61. Schrieffer, A., J. R. Maltez, N. Silva, M. Y. Stoeckle, M. Barral-Netto, and L. W. Riley. 1999. Expression of a pilin subunit BfpA of the bundle-forming pilus of enteropathogenic *Escherichia coli* in an *aroA* live *Salmonella* vaccine strain. *Vaccine* **17**:770–778.
 62. Seifert, H. S. 1996. Questions about gonococcal pilus phase and antigenic variation. *Mol. Microbiol.* **21**:433–440.
 63. Selander, R. K., D. A. Caugant, H. Ochman, J. M. Musser, M. N. Gilmour, and T. S. Whittam. 1986. Methods of multilocus enzyme electrophoresis for bacterial population genetics and systematics. *Appl. Environ. Microbiol.* **51**:873–884.
 64. Sohel, I., J. L. Puente, W. J. Murray, J. Vuopio-Varkila, and G. K. Schoolnik. 1993. Cloning and characterization of the bundle-forming pilin gene of enteropathogenic *Escherichia coli* and its distribution in *Salmonella* serotypes. *Mol. Microbiol.* **7**:563–575.
 65. Sohel, I., J. L. Puente, S. W. Ramer, D. Bieber, C.-Y. Wu, and G. K. Schoolnik. 1996. Enteropathogenic *Escherichia coli*: identification of a gene cluster coding for bundle-forming pilus morphogenesis. *J. Bacteriol.* **178**:2613–2628.
 66. Stone, K. D., H.-Z. Zhang, L. K. Carlson, and M. S. Donnenberg. 1996. A cluster of fourteen genes from enteropathogenic *Escherichia coli* is sufficient for biogenesis of a type IV pilus. *Mol. Microbiol.* **20**:325–337.
 67. Strom, M. S., and S. Lory. 1993. Structure-function and biogenesis of the type IV pili. *Annu. Rev. Microbiol.* **47**:565–596.
 68. Taylor, J., and R. E. Charter. 1952. The isolation of serological types of *Bact. coli* in two residential nurseries and their relation to infantile gastroenteritis. *J. Pathol. Bacteriol.* **64**:715–728.
 69. Taylor, J., and R. E. Charter. 1955. *Escherichia coli* O.128 causing gastroenteritis of infants. *J. Clin. Pathol.* **8**:276–281.
 70. Tennent, J. M., and J. S. Mattick. 1994. Type 4 fimbriae, p. 127–146. *In* P. Klemm (ed.), *Fimbriae: adhesion, genetics, biogenesis, and vaccines*. CRC Press, Inc., Boca Raton, Fla.
 71. Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**:4673–4680.
 72. Tobe, T., T. Hayashi, C. G. Han, G. K. Schoolnik, E. Ohtsubo, and C. Sasakawa. 1999. Complete DNA sequence and structural analysis of the enteropathogenic *Escherichia coli* adherence factor plasmid. *Infect. Immun.* **67**:5455–5462.
 73. Toledo, M. R., M. Alvariza, J. Murahovschi, S. R. Ramos, and L. R. Trabulsi. 1983. Enteropathogenic *Escherichia coli* serotypes and endemic diarrhea in infants. *Infect. Immun.* **39**:586–589.
 74. Tornieporth, N. G., J. John, K. Salgado, P. de Jesus, E. Latham, M. C. Melo, S. T. Gunzburg, and L. W. Riley. 1995. Differentiation of pathogenic *Escherichia coli* strains in Brazilian children by PCR. *J. Clin. Microbiol.* **33**:1371–1374.
 75. Tsukamoto, T. 1996. PCR methods for detection of enteropathogenic *Escherichia coli* (localized adherence) and enteroaggregative *Escherichia coli*. *Kansenshogaku Zasshi* **70**:569–573.
 76. Vuopio-Varkila, J., and G. K. Schoolnik. 1991. Localized adherence by

- enteropathogenic *Escherichia coli* is an inducible phenotype associated with the expression of new outer membrane proteins. *J. Exp. Med.* **174**:1167–1177.
77. **Wall, D., and D. Kaiser.** 1999. Type IV pili and cell motility. *Mol. Microbiol.* **32**:1–10.
78. **Whittam, T. S., and E. A. McGraw.** 1996. Clonal analysis of EPEC serogroups. *Rev. Microbiol. São Paulo* **27**(Suppl. 1):7–16.
79. **Whittam, T. S., M. L. Wolfe, I. K. Wachsmuth, F. Ørskov, I. Ørskov, and R. A. Wilson.** 1993. Clonal relationships among *Escherichia coli* strains that cause hemorrhagic colitis and infantile diarrhea. *Infect. Immun.* **61**:1619–1629.
80. **Wieler, L. H., E. Vieler, C. Erpenstein, T. Schlapp, H. Steinruck, R. Bauerfeind, A. Byomi, and G. Baljer.** 1996. Shiga toxin-producing *Escherichia coli* strains from bovines: association of adhesion with carriage of *eae* and other genes. *J. Clin. Microbiol.* **34**:2980–2984.
81. **Zhang, H.-Z., and M. S. Sonnenberg.** 1996. DsbA is required for stability of the type IV pilin of enteropathogenic *Escherichia coli*. *Mol. Microbiol.* **21**:787–797.

Editor: A. D. O'Brien