

Two Novel Superantigens Found in Both Group A and Group C *Streptococcus*

Thomas Proft, Phillip D. Webb, Vanessa Handley, and John D. Fraser*

Department of Molecular Medicine and Pathology, Faculty of Medical and Health Sciences,
University of Auckland, Auckland, New Zealand

Received 16 July 2002/Returned for modification 16 October 2002/Accepted 21 November 2002

Two novel streptococcal superantigen genes (*speL_{Se}* and *speM_{Se}*) were identified from the *Streptococcus equi* genome database at the Sanger Center. Genotyping of 8 *S. equi* isolates and 40 *Streptococcus pyogenes* isolates resulted in the detection of the orthologous genes *speL* and *speM* in a restricted number of *S. pyogenes* isolates (15 and 5%, respectively). Surprisingly, the novel superantigen genes could not be found in any of the analyzed *S. equi* isolates. The results suggest that both genes are located on a mobile element that enables gene transfer between individual isolates and between streptococci from different Lancefield groups. *S. equi* pyrogenic exotoxin L (SPE-L_{Se})/streptococcal pyrogenic exotoxin L (SPE-L) and SPE-M_{Se}/SPE-M are most closely related to SMEZ, SPE-C, SPE-G, and SPE-J, but build a separate branch within this group. Recombinant SPE-L (rSPE-L) and rSPE-M were highly mitogenic for human peripheral blood lymphocytes, with half-maximum responses at 1 and 10 pg/ml, respectively. The results from competitive binding experiments suggest that both proteins bind major histocompatibility complex class II at the β -chain, but not at the α -chain. The most common targets for both toxins were human V β 1.1 expressing T cells. Seroconversion against SPE-L and SPE-M was observed in healthy blood donors, suggesting that the toxins are expressed *in vivo*. Interestingly, the *speL* gene is highly associated with *S. pyogenes* M89, a serotype that is linked to acute rheumatic fever in New Zealand.

Bacterial superantigens (SAGs) are small highly mitogenic proteins that are produced as monomeric precursor molecules with typical bacterial signal peptides (8). The mature secreted toxins simultaneously bind to major histocompatibility complex (MHC) class II antigens and T-cell-receptor (TcR) molecules, which leads to the stimulation of large numbers of T cells (16, 19, 27). Unlike conventional antigens, SAGs bind MHC class II outside the peptide binding groove and bind mainly to the CDR2 and HV4 regions on the TcR V β chain (12, 15, 17, 27, 38). Each SAG targets a specific set of TcR V β chains leaving a characteristic “fingerprint” in the stimulated T cells. The oligoclonal T-cell stimulation results in high systemic levels of the proinflammatory cytokines tumor necrosis factor alpha and interleukin-1 β (IL-1 β) and T-cell mediators, such as IL-2 and gamma interferon, that are released from antigen presenting cells and T cells (13, 27). This can cause hypotension, fever, and shock (9, 24, 37).

SAGs are produced mainly by the gram-positive bacteria *Streptococcus pyogenes* and *Staphylococcus aureus* (2, 27). In addition, some SAGs have also been found in other bacteria, e.g., the *Mycoplasma arthritidis* mitogen (10) and the *Yersinia pseudotuberculosis* mitogen (23).

S. pyogenes, a Lancefield group A streptococcus, is a human pathogen that causes a wide range of diseases, including acute tonsillitis, scarlet fever, necrotizing fasciitis, bacteremia, streptococcal toxic shock syndrome, and rheumatic fever (6, 11, 37).

The streptococcal SAGs, which include streptococcal pyrogenic exotoxin A (SPE-A), -C, and -G to -J; SSA; and several allelic variants of streptococcal mitogenic exotoxin Z (SMEZ) are believed to play an important role as virulence factors in at least some of these diseases (7, 9, 24). Enterotoxins from *S. aureus*, which include staphylococcal enterotoxin A (SEA) through SEM and toxic shock syndrome toxin (TSST), cause food poisoning (except TSST) and toxic shock syndrome (13, 27, 23a).

Recently, two pyrogenic exotoxins from *Streptococcus equi* (SePE-H and SePE-I) were identified (4). These toxins are nearly identical to SPE-H and SPE-I, differing in just 3 amino acids (aa) and 1 aa, respectively. *S. equi* is a Lancefield group C streptococcus that causes strangles, a contagious inflammatory disease of the upper respiratory tract and associated lymph nodes of equids (18).

The structurally related staphylococcal and streptococcal SAGs share primary sequence identities from 20 to > 90% and have molecular masses of 24 to 28 kDa. They can be classified into four subfamilies (i) group A, which comprises SPE-C, SPE-G, SPE-J, and SMEZ; (ii) group B, which contains SEB, SEC1-3, SEG, SPE-A, and SSA; (iii) group C, with SEA, SED, SEE, SEH, and SEJ; (iv) group D, which contains SEI, SPE-I, and SePE-I; and group E, with SPE-H and SePE-H. TSST builds a separate branch within this family tree.

Several *sag* genes are located on mobile DNA elements, such as bacteriophage, integrated in the bacterial genome (14). Horizontal transfer of these mobile DNA elements is a proposed mechanism that might explain the close relation between SAGs from different bacterial species. This could also explain why individual bacterial isolates carry only certain combinations of *sag* genes. For example, *speH* and *speI* were found

* Corresponding author. Mailing address: Department of Molecular Medicine and Pathology, Faculty of Medical and Health Sciences, University of Auckland, Private Bag 92019, Auckland, New Zealand. Phone: 64-9-373-7599, ext. 6036. Fax: 64-9-373-7492. E-mail: jd.fraser@auckland.ac.nz.

in *S. pyogenes* isolates at frequencies of only 24 and 16%, respectively, and both genes are located on prophage SF370.2 (14, 32). Thus far, no streptococcal or staphylococcal isolate has been found that carries a complete set of known *sag* genes.

Crystal structures of SEA, SEB, SEC2, SED, TSST, SPE-A, SPE-C, SPE-H, and SMEZ-2 show a conserved folding pattern comprising an NH₂-terminal β -barrel globular domain and a COOH-terminal globular domain based on a β -grasp motif (3, 31).

Some SAGs, such as TSST and SEB, bind exclusively to the invariant MHC class II α -chain via a generic binding site located within the β -barrel domain (21, 36). Other SAGs, such as SPE-C, SPE-G, SPE-H, and SPE-J, contain a zinc binding motif within the COOH-terminal domain that is used to bind the polymorphic MHC class II β -chain via a tetrameric zinc coordination complex that involves the highly conserved His 81 residue on the HLA-DR1 β -chain (25, 26).

Recently, we identified four novel *sag* genes by mining the *S. pyogenes* M1 genomic database at the University of Oklahoma. The genes for *speG*, *speH*, *speJ*, and *speI* were cloned and expressed in *Escherichia coli*, and the purified recombinant proteins showed the typical SAG features (32, 33). The *S. pyogenes* genome project was completed in 2000, and only one more *sag* gene was found; it was named *speK*. However, *speK* is a pseudogene with an incomplete open reading frame (ORF) (14).

We have now extended our search for novel SAGs by mining other streptococcal genome databases. We here describe the identification of two novel *sag* genes from *S. equi* (*speL_{se}* and *speM_{se}*) and their respective orthologues in *S. pyogenes* (*speL* and *speM*). (Two genes in this gene family were designated *sepe-h* and *sepe-i* in a previous publication [4].) One intriguing feature of the *speL* gene is the strong association to the M89 serotype, which is linked to acute rheumatic fever in New Zealand.

MATERIALS AND METHODS

Bioinformatic tools. The novel *S. equi* SAGs were identified by searching the microbial genome databases at the National Center for Biotechnology Information (http://www.ncbi.nlm.nih.gov/Microb_blast/unfinishedgenome.html) with highly conserved $\beta 5$ and $\alpha 4$ regions of streptococcal and staphylococcal SAGs, using a TBLASTN search program.

The ORFs were defined by translating the DNA sequences around the matching regions and aligning the protein sequences to known SAGs using ClustalW at <http://clustalw.genome.ad.jp> (39) and ClustalX. The leader sequences were predicted using the SignalP V1.1 server at <http://www.cbs.dtu.dk/services/SignalP>. (29). The SAG family tree was generated using ClustalW and the TreeView Computer program (30).

Genotyping of *S. equi* and *S. pyogenes* isolates. The New Zealand *S. pyogenes* isolates were obtained from the Institute of Environmental Sciences and Research in Porirua, New Zealand. Isolate 33117 was obtained from Pasteur Institute, Lille, France. The *S. equi* isolates were obtained from the Department of Microbiology, Otago University, Dunedin, New Zealand.

The streptococcal isolates were grown in brain heart infusion medium (Difco) at 37°C, and genomic DNA was prepared as described previously (34). Genotyping was carried out by PCR with 50 ng of purified streptococcal genomic DNA using the specific primer pairs *sepe-l/spe-l* (5'-GCGGATCCGATACGTACAA TACAAATG-3'; 5'-GCGAATCAATAGCATTGACC-3'), *sepe-m/spe-m* (5'-GCGGATCCGAGGGGACTATTAATATTAAG-3'; 5'-GCGAATTCGGT TCCTTGACTACTAAC-3').

A primer pair specific to a DNA region encoding the 23S rRNA (33) was used as a positive control.

Generation of recombinant SPE-L (rSPE-L) and rSPE-M. The *spe-l* and *spe-m* genes were amplified by PCR from 50 ng of genomic DNA of isolates 10846 and

FP 4223, respectively, using the same primers as for genotyping. The PCR products encoding the mature proteins without the predicted leader sequences were enriched using the Wizard PCR DNA purification system (Promega) and cloned into T-tailed pBluescript SK II vectors (Stratagene), and this was followed by transformation into *E. coli* XL1 blue.

The cloned *sag* genes were analyzed by the dideoxy chain termination method using a Licor automated DNA sequencer (model 4200). Three subclones from every cloning experiment were analyzed to ensure that no *Taq* polymerase-related mutations were introduced. The DNA sequences have been annotated in the EMBL/GenBank/DBJ databases.

The *speL* and *speM* genes were cloned from pBluescript into the *Bam*HI-*Eco*RI cloning site of pET32-3c expression vector. This vector is a modified version of pET32a (Novagene) that expresses the highly specific protease 3C cleavage site (EVLVQ|GP [vertical bar indicates cleavage site]) from a picornavirus (40) just upstream of the inserted DNA. rSPE-L and rSPE-M were expressed in *E. coli* AD494(DE3) (Novagene) as thioredoxin (Trx) fusion proteins. Cultures were grown at 30°C and induced for 3 to 4 h after adding 0.2 mM isopropyl- β -D-thiogalactopyranoside (IPTG).

The Trx fusion proteins were purified on Ni²⁺ IDA Sepharose (Sigma), and the mature proteins were cleaved off from Trx on the column using protease 3C (provided by Keith Hudson, University of Oxford, Oxford, United Kingdom) overnight at room temperature. The eluates containing the recombinant toxins were collected and analyzed on sodium dodecyl sulfate–12.5% polyacrylamide gels according to the procedure of Laemmli.

Toxin proliferation assay. Human peripheral blood lymphocytes (PBLs) were purified from blood of a healthy donor by Histopaque Ficoll (Sigma) fractionation. The PBLs were incubated in 96-well round-bottom microtiter plates at 10⁵ cells per well with RPMI-10 (RPMI with 10% fetal calf serum) containing various dilutions of recombinant toxins. The dilution series was performed in 1:5 steps from a starting toxin concentration of 10 ng/ml. After 3 days 0.1 μ Ci of [³H]thymidine was added to each well and cells were incubated for another 24 h. Cells were harvested, and the radiation in cells was counted on a scintillation counter.

TcR V β analysis. V β enrichment analysis was performed by anchored multiprimer amplification. Human PBLs were incubated with recombinant toxin (20 pg/ml) at 10⁶ cells/ml for 3 days. A twofold volume expansion of the culture followed, with medium containing IL-2 at 20 ng/ml. After another 24 h, stimulated and resting cells were harvested and RNA was prepared using Trizol reagent (Life Technologies). A 500-bp β -chain DNA probe was obtained by anchored multiprimer PCR as described previously (20), radiolabeled, and hybridized to individual V β s and a C β DNA region dot blotted on a nylon membrane. The membrane was analyzed on a Molecular Dynamics Storm PhosphorImager using ImageQuant software. Individual V β s were expressed as a percentage of all the V β s determined by hybridization to the C β probe.

Radiolabeling and LG-2 binding experiments. Recombinant toxin was radiolabeled by the chloramine T method as previously described (25). Labeled toxin was separated from free iodine by size exclusion chromatography using Sephadex G-25 (Pharmacia). LG2 cells were used for cell binding experiments, as previously described (25). Briefly, cells were harvested, resuspended in RPMI-10, and mixed at 10⁶ cells/ml with ¹²⁵I-tracer toxin (1 ng) and 0.0001 to 10 μ g of unlabeled toxin and incubated at 37°C for 1 h. After washing with ice cold RPMI the pelleted cells were analyzed in a gamma counter.

For competitive binding studies, 1 ng of ¹²⁵I-tracer toxin (rSPE-L or rSPE-M) was incubated with 0.0001 to 10 μ g of unlabeled toxin (rSPE-L, rSPE-M, rSPE-C, rSEB, and rTSST) for 1 h.

Seroconversion experiments. Five microliters of serum from human healthy donors were mixed with 10 μ l of phosphate-buffered saline and 10 μ l of ¹²⁵I-labeled rSPE-L or rSPE-M and incubated for 1 h at 37°C. As controls, bovine serum albumin or rabbit antiserum against SPE-C or SMEZ was used instead of human sera. After adding 50 μ l of protein A-staphylococcus cells (Toxin Technology) the samples were incubated for another 30 min at 4°C before the cells were spun down, washed in 1 ml of phosphate-buffered saline, and counted in a Cobra Gamma counter.

Nucleotide sequence accession numbers. The accession numbers for the *speL* and *speM* genes are AF514282 and AF514283, respectively.

RESULTS

Identification and sequence analysis of SAGs. A number of complete and incomplete microbial genome projects are accessible at the National Center for Biotechnology Information Web page, which can be mined for DNA sequences of interest

using standard BLAST search tools (http://www.ncbi.nlm.nih.gov/Microb_blast/unfinishedgenome.html). The genome databases of *Streptococcus gordonii*, *Streptococcus pneumoniae*, *S. equi*, and *Streptococcus mutans* were searched with a SA_g peptide sequence corresponding to the highly conserved α 4 region using the TBLASTN program.

Four significant matches and predicted ORFs with homology to streptococcal and staphylococcal SA_gs were found by aligning translated DNA sequences with complete protein sequences of known SA_gs. All ORFs were identified from the incomplete *S. equi* genome database at the Sanger Center, which contains a collection of more than 200 DNA sequence contiguous sequences derived from a shotgun plasmid library of the complete *S. equi* genome.

Two of these ORFs correlate with the previously described SePE-I and SePE-H (4). The third ORF identified showed 99% nucleotide identity to the hypothetical exotoxin gene *speL* found on the prophage Φ NIH1.1. sequence (NC003157) (22). This was labeled *speL_{Se}* according to the *S. equi* nomenclature (*S. equi* pyrogenic exotoxin L). The fourth ORF showed no significant sequence homology to any known SA_g, but contained the highly conserved SA_g α 4 peptide sequence and was named *speM_{Se}*. The *speL_{Se}* and *speM_{Se}* genes are adjacent at the 5' end of a 4,470-bp contiguous sequence separated by a 266-bp intergenic region.

Genotyping of *S. equi* and *S. pyogenes* isolates. Genomic DNA from eight *S. equi* isolates were screened by PCR using specific primer pairs for the *speL_{Se}* and *speM_{Se}* genes. In addition, a primer pair specific to a streptococcal DNA region encoding the 23S rRNA was used as a positive control. Neither *speL_{Se}* nor *speM_{Se}* could be detected in any of the *S. equi* isolates, suggesting that they are located on mobile DNA elements and thus restricted in their distribution to certain *S. equi* isolates (data not shown).

Screening of *S. pyogenes* isolates (including 29 M-types and three MNT) identified the *speL_{Se}* gene in 6 of 40 cases (15%) and the *speM_{Se}* gene in 2 of 40 cases (5%) (Table 1). The *speL_{Se}* gene was restricted to only 5 M-types (M28, M41, M56, M59, and M89), and the *speM_{Se}* gene was found in only 2 M-types (M80 and M92).

Interestingly, the M28 isolate was obtained from a patient suffering from streptococcal toxic shock syndrome, and M89 strains are strongly associated with acute rheumatic fever in New Zealand (28).

New Zealand *S. pyogenes* M89 isolates were genotyped for the *spe-l* gene, and 8 of 11 (73%) were positive, indicating a strong association between this M-type and the *speL* gene (Table 1).

Cloning and DNA analysis of *speL* and *speM* from *S. pyogenes*. The *speL_{Se}* gene was cloned from isolate 10846 (M89), and DNA sequence analysis revealed a strong sequence homology (99% nucleotide identity) to the *S. equi speL_{Se}*. The gene was named *speL* according to the *S. pyogenes* nomenclature for the streptococcal pyrogenic exotoxin family. The *speL_{Se}* and *speL* genes differ by only 7 bp, and the deduced protein sequences differ by 4 aa, of which two differences are conservative exchanges (Y225H and E229D) and two are moderate (I112S and N233D). Furthermore, the *speL* gene is 100% identical to the hypothetical exotoxin gene *speL* found on prophage Φ NIH1.1. sequence (NC003157).

The *speM_{Se}* was cloned from isolate FP4223 (serotype M80), and DNA sequence comparison showed strong nucleotide homology to *speM_{Se}* from *S. equi* (98.1% nucleotide identity) and was therefore labeled *spe-m*. The *speM_{Se}* and *speM* genes differ by 13 bp, and the deduced protein sequences differ by 8 aa, of which 5 are conservative exchanges (V41I, I51M, H55Y, Y74H, and E137D). The three nonconservative exchanges are E37V, G175D, and N191K.

The calculated molecular masses are 27.4 kDa (SPE-L_{Se} and SPE-L) and 26.2 kDa (SPE-M_{Se} and SPE-M), and the calculated isoelectric points are 6.69 (SPE-L_{Se}), 6.53 (SPE-L), 6.7 (SPE-M_{Se}), and 7.1 (SPE-M). The minor differences between SPE-L_{Se} and SPE-L and between SPE-M_{Se} and SPE-M suggest that the proteins are orthologous with identical or very similar functions in the two streptococcal species.

A revised SA_g family tree, based on primary amino acid sequence homology, shows that SPE-L_{Se} and SPE-L as well as SPE-M_{Se} and SPE-M belong to the same clade as SMEZ, SPE-C, SPE-G, and SPE-J (group A) but build a separate branch within that group (Fig. 1). SPE-L_{Se}-SPE-L and SPE-M_{Se}-SPE-M are most closely related to each other (42% amino acid identity and 51% similarity) followed by SPE-C (32% amino acid identity and 40% similarity).

All novel SA_gs contain a zinc binding motif near the C terminus that is common among most of the streptococcal SA_gs, including all members of group A. Zinc is essential for binding to the polymorphic MHC class II β -chain via a divalent zinc ion (26, 35). In SPE-C, the zinc-binding residues are H167, H201, and D203, and in SMEZ-2 the zinc ion is bound by residues H162, H202, and D204 (3). The multiple protein sequence alignment (Fig. 2) shows that the most likely candidates for zinc binding are residues H191, H221, and D223 for SPE-L_{Se} and SPE-L and H186, H217, and D218 for SPE-M_{Se} and SPE-M.

One important difference between the novel SA_gs and other streptococcal SA_gs is a gap in amino acid sequence between residues 207 and 208 (in SPE-L_{Se} and SPE-L) and residues 202 and 203 (in SPE-M_{Se} and SPE-M) (Fig. 2). In the protein structures of SPE-C, SPE-H, and SMEZ-2, the amino acids in this region form part of the β 10- α 5 loop and the structure determining the α 5 helix (3, 35). Structure-determining regions, unlike surface loops, are generally strongly conserved in SA_gs, suggesting that there is potential for a major difference in the SPE-L/SPE-M protein structure in this region compared to other SA_gs.

Expression of rSPE-L and rSPE-M. To produce recombinant proteins of SPE-L and SPE-M, the individual subcloned genes (coding for the mature toxins without leader sequence) were transferred to expression vector pET32-3c and transformed into *E. coli* AD494(DE3). Fusion proteins of Trx-SPE-L and Trx-SPE-M were completely soluble and gave yields of approximately 20 mg per liter. The recombinant toxins were cleaved off from the Trx using protease 3C, while still bound on the affinity column. The recombinant proteins differ from their native counterparts by four additional N-terminal residues (GPGS) introduced by the *Bam*HI restriction site and the protease 3C recognition site. However, a potential influence on the function of the recombinant proteins seems rather unlikely, as the crystal structures of the closely related proteins SMEZ-2, SPE-C, and SPE-H show that the N terminus is

TABLE 1. Genotyping of *S. pyogenes* isolates from New Zealand^a

<i>S. pyogenes</i> isolate	M/emm	Site	Disease or condition	PCR result with primer for:										
				23S	<i>speA</i>	<i>speC</i>	<i>speG^b</i>	<i>speH^b</i>	<i>speI</i>	<i>speJ</i>	<i>ssa</i>	<i>speL</i>	<i>speM</i>	<i>smez^b</i>
FP 1943	M53	TS	ST	+	-	+	+	-	-	+	-	-	-	+
FP 2658	M59	TS	ST	+	-	-	+	-	-	+	-	+	-	+
FP 4223	M80	TS	ST	+	-	-	+	-	-	+	-	-	+	+
FP 5417	M41	TS	ST	+	+	-	+	-	-	+	-	+	-	+
FP 5971	M57	TS	ST	+	+	+	+	+	-	+	-	-	-	+
1/5045	M4	TS	ST	+	+	+	+	-	-	+	+	-	-	+
82/20	M4	SK	Ulcer	+	-	-	+	-	-	+	+	-	-	+
85/167	M12	TS	ST	+	-	-	+	+	+	+	-	-	-	+
85/314	M89	WS	Wound	+	-	-	+	-	-	+	-	+	-	+
85/437	M81	WS	Infections eczema	+	-	+	+	-	-	+	+	-	-	+
85/723	M22	Ear	Otitis	+	-	+	+	-	-	+	+	-	-	+
86/435	M4	TS	RF	+	-	+	+	-	-	+	+	-	-	+
87/19	M12	TS	ST	+	+	+	+	+	+	+	-	-	-	+
89/26	M1	TS	AGN	+	+	-	+	-	-	+	-	-	-	+
90/424	M4	TS	ST	+	+	+	+	-	-	+	+	-	-	+
94/229	M49	HVS	Endometritis	+	+	-	+	+	+	+	-	-	-	- ^c
94/712	M89	WS	Cellulitis	+	-	+	+	-	-	+	-	-	-	+
95/31(2)	M89	WS	Abscess	+	-	-	+	-	-	+	-	-	-	+
96/1	M4	HVS	Endometritis	+	-	+	+	-	-	+	+	-	-	+
9779	emm56	TS	ST	+	-	-	+	-	-	+	-	+	-	+
9893	M82	TS	ST	+	-	+	+	+	+	+	-	-	-	+
10019	emm44	TS	ST	+	-	+	+	+	+	+	-	-	-	+
10303	emm59	TS	ST	+	-	-	+	-	-	+	-	+	-	+
10438	ST3018	TS	ST	+	+	-	+	-	-	+	-	-	-	+
10463	emm49	TS	ST	+	+	-	+	-	-	+	+	-	-	+
10649	ST2267	TS	ST	+	-	-	+	-	-	+	-	-	-	+
10763	M88	TS	ST	+	-	+	+	-	-	+	+	-	-	+
10791	MNT	TS	ST	+	-	+	+	+	+	+	+	-	-	+
10989	M87	TS	ST	+	-	-	+	-	-	+	-	-	-	+
11070	emm65	TS	ST	+	-	-	+	+	-	+	-	-	-	- ^c
11152	M85	TS	ST	+	-	-	+	+	-	+	-	-	-	- ^c
11222	M92	TS	ST	+	+	+	+	+	-	+	-	-	+	+
11227	emm14	TS	ST	+	-	-	+	-	-	+	-	-	-	+
11276	MNT	TS	ST	+	+	-	+	-	-	+	+	-	-	+
11574	ST809	TS	ST	+	-	-	+	-	-	+	-	-	-	+
11646	ST4547	TS	ST	+	-	-	+	-	-	+	+	-	-	+
11686	M91	TS	ST	+	-	-	+	-	-	+	+	-	-	+
11789	MNT	TS	ST	+	+	+	+	-	-	+	-	-	-	+
ATCC	M1	Wound	ND	+	-	+	+	+	+	+	-	-	-	+
33117	T28R28	ND	STSS	+	-	-	+	-	-	+	-	+	-	+
% With positive PCR result				100	31	42	100	26	16	100	32	15	5	92
Strain M89														
85/314	M89	WS	Wound	+	-	-	+	-	-	+	-	+	-	+
95/361	M89	PS	Abscess	+	-	-	+	+	-	+	-	+	-	+
82/675	M89	WS	Wound	+	-	+	+	-	-	+	-	+	-	+
95/31	M89	WS	Abscess	+	-	-	+	-	-	+	-	-	-	+
94/712	M89	WS	Cellulitis	+	-	+	+	-	-	+	-	+	-	+
10846	M89	TS	ST	+	+	-	+	-	-	+	-	+	-	+
89/54	M89	TS	ST	+	+	-	+	-	-	+	-	+	-	+
94/11	M89	PS	Abscess	+	-	-	+	-	-	+	-	+	-	+
95/127	M89	BC	Cellulitis	+	-	-	+	-	-	+	-	+	-	+
84/781	M89	TS	ST	+	+	-	+	-	-	+	-	-	-	+
96/364	M89	BC	Burns	+	-	-	+	-	-	+	-	-	-	+
% With positive PCR result				100	27	18	100	9	0	100	0	73	0	100

^a Fifty-one *S. pyogenes* isolates (38 from New Zealand, 1 from France, 1 ATCC reference strain, and 11 *S. pyogenes* M89 isolates collected in New Zealand) were genotyped. The results are based on PCR analysis using purified genomic DNA and specific primers for each *sag* gene. The primers for *speL* and *speM* were designed using the DNA sequences of the orthologous *speL_{spe}* and *speM_{spe}* genes from *S. equi*. Site and disease abbreviations are as follows: TS, throat site; WS, wound site; SK, skin; PS, pus site; HVS, high vaginal site; BC, blood culture; ST, sore throat; RF, rheumatic fever; AGN, acute glomerulonephritis; STSS, streptococcal toxic shock; ND, not determined.

^b Reported previously (34).

^c Positive by Southern blotting.




FIG. 1. Family tree of streptococcal and staphylococcal SAGs. The tree was created using ClustalW (39) and the TreeView computer program (30). The novel SAGs belong to group A, together with SPE-C, SPE-G, SPE-J, and SMEZ, but build a separate branch within this group.

facing away from both the TcR and the MHC class II binding site (3, 35).

Eluates containing the purified toxins were collected and analyzed for purity on a sodium dodecyl sulfate–12% polyacrylamide gel.

To ensure the native conformation of the purified recombinant toxins, a standard [^3H]thymidine incorporation assay was performed to test for their potency to stimulate PBLs. Both toxins were active on human T cells (Fig. 3). rSPE-C was included as a reference protein.

The half-maximal responses for rSPE-L and rSPE-M were 1 and 10 pg/ml, respectively. No activity was detected at less than 0.1 pg/ml. SPE-L was as potent as rSPE-C that also had a half-maximal value of 1 pg/ml. rSPE-M was less active than rSPE-C but still more potent than the previously described SPE-H (50 pg/ml) (33).

Significant seroconversion against SPE-L and SPE-M was detected in sera from healthy human blood donors. Out of twenty subjects, seven were identified with seroconversion against SPE-L, and six serum samples had significant antibody titers against SPE-M (data not shown). These results suggest that both toxins are actually expressed by the bacteria.

V β specificity of recombinant toxins. The human TcR V β specificity of the recombinant toxins was determined by multiprimer anchored PCR and dot blot analysis using a panel of 21 human V β DNA regions. The V β enrichment after stimulation with toxin was compared to the V β profile of PBLs stimulated with the unspecific T-cell mitogen concanavalin A (ConA).

Both toxins showed very similar results, with V β 1.1 TcR being the primary target. T cells carrying the V β 1.1 TcR were stimulated >80 times and nearly 70 times more with rSPE-L or rSPE-M, respectively, than with ConA (Table 2).

Interestingly, the total of the V β s stimulated with rSPE-L or rSPE-M was only 78 and 82%, respectively, suggesting that one or more additional V β s were stimulated which are not represented in the panel. To further investigate this possibility, the V β cDNAs from rSPE-L- and rSPE-M-stimulated T cells were cloned into a pBluescript vector and the DNA sequences of 10 randomly selected clones were analyzed. Five out of five cDNAs from rSPE-L-stimulated T cells and four of five cDNAs from rSPE-M stimulated T cells were identified as V β 1.1. One rSPE-M-expanded cDNA was identified as V β 5.8. However, V β 5.8 was not significantly enriched by either toxin (around 4.5%) compared to ConA-stimulated cells (3.3%) (Table 2).

MHC class II binding. As previously mentioned, the protein sequences of SPE-L and SPE-M contain the conserved zinc binding motif that is required for binding to the polymorphic MHC class II β -chain.

In an attempt to determine the orientation of the toxins on MHC class II, competition binding experiments were performed. The recombinant toxins were radiolabeled and tested with excess of unlabeled toxin for binding to LG-2 cells. rSEB, rTSST, and rSPE-C were used as reference proteins. SPE-C binds exclusively to the MHC class II β -chain using the zinc binding motif (25, 26), while SEB and TSST both exclusively bind to the MHC class II α -chain in a different, zinc-independent binding mode (1, 21, 36). The results are shown in Fig. 4. Binding of radiolabeled rSPE-L or rSPE-M was inhibited with excess amounts of unlabeled rSPE-C, rSPE-L, and rSPE-M. In contrast, excess amounts of unlabeled rSEB or rTSST did not influence binding of radiolabeled rSPE-L and rSPE-M. These results suggest that both, SPE-L and SPE-M, bind MHC class II exclusively to the β -chain in a zinc-dependent binding mode. This is a consistent feature among all group A members of the SAG family.

DISCUSSION

Two novel *sag* genes, *speL_{se}* and *speM_{se}*, have been identified from the *S. equi* genome database at the Sanger Center. Subsequently, the orthologue genes *speL* and *speM* genes were identified from *S. pyogenes*.

The increasing number of genome sequencing projects and the availability of the raw sequencing data on the Internet facilitates the rapid identification of novel genes by sequence similarity. We have recently increased the number of known streptococcal SAGs after searching the *S. pyogenes* M1 genome database at Oklahoma University (32, 33). Interestingly, only three *sag* genes (*speG*, *speI*, and *smez*) were found on all analyzed *S. pyogenes* isolates, whereas all other streptococcal SAG genes




FIG. 2. Multiple alignment of streptococcal SAg protein sequences. The protein sequences of mature proteins were aligned using the ClustalX computer program. The clear box near the C terminus represents a primary zinc binding motif (H-X-D), a feature of all streptococcal toxins except SPE-A and SSA. Structural elements of SMEZ-2 and SPE-C, respectively, are shown below.

Downloaded from http://iai.asm.org/ on February 24, 2021 by guest




FIG. 3. Stimulation of human T cells with recombinant toxins. PBLs were isolated from human blood samples and incubated with various concentrations of recombinant toxin (in duplicates). After 3 days, 0.1 μ Ci of [³H]thymidine was added, and cells were incubated for another 24 h before being harvested and counted on a scintillation counter.

occur with different frequencies due to their location on mobile genetic elements. Horizontal gene transfer appears to be the likely mechanism for the distribution of *sag* genes, not only between different *S. pyogenes* isolates, but also between different streptococcal species. The *speJ* and *speI* genes occur with frequencies of 26 and 16%, respectively, within *S. pyogenes* isolates, and orthologues of these genes (*sepe-h* and *sepe-i*) have recently been found in *S. equi* (4). This suggests that many

more SAGs are likely to exist in other streptococcal isolates. Genotyping of 40 *S. pyogenes* isolates showed that *speL* and *speM* occur with low frequencies (15 and 5%, respectively) and are therefore likely to be on mobile DNA element(s). The *S. equi* orthologues *speL_{Se}* and *speM_{Se}* could not be detected in any of the analyzed *S. equi* isolates, suggesting that they are rare in this species. The *speL* gene is identical to a hypothetical SAG gene that has recently been identified on the streptococcal prophage Φ NIH1.1. Ikebe et al. showed that all of 18 M3/T3 isolates recovered from Japan during or after 1992 have the *speL* gene, while none of 10 T3 isolates recovered before 1973 have this gene (22). This suggests a very recent horizontal transfer of *speL* between different *S. pyogenes* isolates and between *S. pyogenes* and *S. equi*. The fact that this prophage does not carry the *speM* gene and the fact that none of our analyzed *S. pyogenes* isolates carry both *sag* genes suggest that *speM* is located on a different prophage. However, the *speI* gene also could not be detected in all *speH*-positive isolates (Table 1), despite the fact that both genes are located on the SF370.2 prophage on the *S. pyogenes* M1 SF370 genome (14). A possible reason for the separation from *speL* is that *speM* might have been lost in some *S. pyogenes* isolates during integration of the prophage into the streptococcal genome.

One of the most interesting outcomes of this study is the finding that the *speL* gene is linked to *S. pyogenes* M89, a serotype that is strongly associated with acute rheumatic fever in New Zealand (28). SAGs have been implicated in acute rheumatic fever, but their role in the pathology of the disease remains elusive (6). Screening of a much larger panel of clinical isolates is currently under way to determine the strength of this linkage between *spe-l* and strains causing acute rheumatic fever.

The novel toxins are closely related to the streptococcal and staphylococcal SAGs. Recombinant forms of SPE-L and SPE-M are extremely potent stimulators of human PBLs at nanomolar concentrations, confirming their role as SAGs. Seroreconversion against SPE-L and SPE-M was observed in several blood samples from healthy donors, suggesting in vivo

TABLE 2. V β specificity of recombinant toxins on human PBLs^a

V β cDNA	% of V β enrichment after stimulation with:		
	ConA	SPE-L	SPE-M
1.1	0.1	<u>8.44</u>	<u>6.75</u>
2.1	3.94	4.46	4.2
3.1	19.57	6.53	9.83
4.1	0.12	0.98	1.41
5.1	1.82	2.78	3.69
5.3	12.06	10.2	13.96
5.8	3.31	4.49	4.43
6.3	1.03	1.57	2.15
6.4	0.63	1.41	1.47
6.9	5.96	4.85	4.92
7.3	0.84	0.94	1.1
7.4	2.31	1.71	1.47
8.1	0.3	0.68	0.79
9.1	1.34	1.78	2.05
12.3	15.36	7.76	8.93
12.5	1.17	1.38	1.94
14.1	0.3	0.79	0.95
15.1	2.26	3.92	4.25
17.1	-0.07	1.05	1.12
18.1	1.51	2.54	3.13
22a	1.1	2.09	2.27
23.1	0.45	1.39	1.41
Total	71.6	78	82.2

^a Human PBLs were incubated with recombinant toxin (20 pg/ml) for 4 days. Relative enrichment of V β cDNAs was analyzed from RNA of toxin-stimulated and ConA-stimulated PBLs by anchored-primer PCR and reverse dot blot to a panel of 22 different V β cDNAs. The figures represent the percentage of each V β with respect to total C β . Significant responses are underlined.




FIG. 4. Competition binding studies with rSPE-L and rSPE-M. LG-2 cells were incubated in duplicate with 1 ng of ^{125}I -labeled recombinant toxin and increasing amounts of unlabeled toxins. After 1 h cells were washed and counted. (A) Competition assay with labeled rSPE-L; (B) competition assay with labeled rSPE-M.

expression of both toxins. Despite their highly conserved core structure, individual SAGs differ remarkably in their surface-exposed regions. This explains why immune responses against SAGs are highly specific, e.g., polyclonal serum raised against a certain SAG does not cross-react with any other SAG (data not shown). Therefore, seroconversion against a particular SAG is

a strong indicator for the actual production of this SAG by the bacteria and also suggests a potential role as a virulence factor.

The data presented here suggest that SPE-L and SPE-M bind MHC class II exclusively at the β -chain via a zinc binding complex. This assumption is supported by the fact that both toxins contain the characteristic zinc binding motif (H-X-D) close to the C terminus and by the results from the competitive binding assays. Excess concentrations of rSPE-C, a toxin that exclusively binds to the MHC class II β -chain (26) clearly inhibited SPE-L and SPE-M from binding to LG-2 cells. In contrast, the typical MHC class II α -chain binding toxins SEB and TSST showed no inhibition effect. This MHC class II binding mode appears to be conserved among all group A members of the SAG family (Fig. 1). SSA and SPE-A are the only streptococcal SAGs known thus far that lack the zinc binding motif and are unable to bind to the MHC class II α -chain. Both toxins belong to the group B clade in the SAG family tree and are more closely related to the staphylococcal enterotoxins than to any other streptococcal toxin.

Interestingly, SPE-L and SPE-M both stimulate mainly V β 1.1-expressing T cells. Usually, SAGs can be distinguished by their characteristic combination of V β targets. This finding might explain why none of the analyzed streptococcal isolates carried both genes, *speL* and *speM*, as the need to activate a single V β can be met by a single toxin and two toxins with the same function would be unlikely to offer a significant selective advantage.

SPE-L and SPE-M differ from their *S. equi* orthologues by only 4 and 8 aa, respectively, and most of the exchanges are conservative. All of the nonconservative exchanges are at amino acid positions that are not well conserved among the SSAs. This suggests that SPE-L_{se} and SPE-M_{se} are likely to function in an identical fashion to their *S. pyogenes* orthologues SPE-L and SPE-M. It is not known yet what role SAGs play in *S. equi*. Artiushin and coworkers recently reported that SePE-I and SePE-H were highly mitogenic on horse PBLs and pyrogenic for rabbits. Furthermore, horses that had recovered from strangles or been immunized with SePE-I were resistant to the pyrogenic effect of SePE-I (4). Interestingly, an epidemic outbreak of nephritis in Nova Serrana, Brazil, has recently been reported, and throat cultures indicated that nephritis was associated with group C *S. equi* (5).

Two novel SAGs were found in streptococcal strains belonging to two different Lancefield groups, group A and group C. The *sag* genes are most likely located on a mobile DNA element that allows horizontal transfer of these genes between different streptococcal species, such as *S. equi* and *S. pyogenes*. The significant association of *speL* to *S. pyogenes* strain M89, which causes acute rheumatic fever, suggests a potential involvement of SPE-L in this disease. However, this has to be reevaluated after screening a larger panel of *S. pyogenes* isolates. It remains to be seen whether *sag* genes will also be found in other non-group A streptococci.

ACKNOWLEDGMENTS

We gratefully acknowledge the *S. equi* Sequencing Group at the Sanger Institute for making their DNA sequences available on the Internet. We thank Diana Martin, Institute of Environmental Sciences and Research, Porirua, New Zealand, for the New Zealand *S. pyogenes* isolates; John Tagg, Otago University, Dunedin, New Zealand, for the

S. equi isolates; and Heide Mueller-Alouf, Pasteur Institute, Lille, France, for the *S. pyogenes* 33117 isolate.

This study was supported by the Health Research Council of New Zealand and by an Early Career Excellence Grant from the University of Auckland.

REFERENCES

- Acharya, K. R., E. F. Passalacqua, E. Y. Jones, K. Harlos, D. I. Stuart, R. D. Brehm, and H. S. Tranter. 1994. Structural basis of superantigen action inferred from crystal structure of toxic-shock syndrome toxin-1. *Nature* **367**: 94–97.
- Alouf, J. E., H. Mueller-Alouf, and W. Koehler. 1999. Superantigenic *Streptococcus pyogenes* erythrogenic/pyrogenic exotoxins, p. 567–588. In J. E. Alouf and J. H. Freer (ed.), *Sourcebook of bacterial protein toxins*. Academic Press, San Diego, Calif.
- Arcus, V. L., T. Proft, J. A. Sigrell, H. M. Baker, J. D. Fraser, and E. N. Baker. 2000. Conservation and variation in superantigen structure and activity highlighted by the three-dimensional structures of two new superantigens from *Streptococcus pyogenes*. *J. Mol. Biol.* **299**:157–168.
- Artiushin, S. C., J. F. Timoney, A. S. Sheoran, and S. K. Muthupalani. 2002. Characterization and immunogenicity of pyrogenic mitogens SePE-H and SePE-I of *Streptococcus equi*. *Microb. Pathog.* **32**:71–85.
- Balter, S., A. Benin, S. W. Pinto, L. M. Teixeira, G. G. Alvim, E. Luna, D. Jackson, L. LaClaire, J. Elliott, R. Facklam, and A. Schuchat. 2000. Epidemic nephritis in Nova Serrana, Brazil. *Lancet* **355**:1776–1780.
- Bathnagar, A., A. Grover, and N. K. Ganguly. 1999. Superantigen-induced T cell responses in acute rheumatic fever and chronic rheumatic heart disease patients. *Clin. Exp. Immunol.* **116**:100–106.
- Bernal, A., T. Proft, J. D. Fraser, and D. N. Posnett. 1999. Superantigens in human disease. *J. Clin. Immunol.* **19**:149–157.
- Betley, M. J., D. W. Borst, and L. B. Regasa. 1992. Staphylococcal enterotoxins, toxic shock syndrome toxin and streptococcal exotoxins: a comparative study of their molecular biology. *Chem. Immunol.* **55**:1–35.
- Bohach, G. A., D. J. Fast, R. D. Nelson, and P. M. Schlievert. 1990. Staphylococcal and streptococcal pyrogenic toxins involved in toxic shock syndrome and related illnesses. *Crit. Rev. Microbiol.* **17**:251–272.
- Cole, B. C., and C. L. Atkin. 1991. The *Mycoplasma arthritis* T-cell mitogen, MAM: a model superantigen. *Immunol. Today* **12**:271–276.
- Cone, L. A., P. M. Woodard, P. M. Schlievert, and G. S. Tomory. 1987. Clinical and bacteriologic observations of a toxic shock-like syndrome due to *Streptococcus pyogenes*. *N. Engl. J. Med.* **317**:146–149.
- Dellabona, P., J. Peccoud, J. Kappler, P. Marrack, C. Benoist, and D. Mathis. 1990. Superantigens interact with MHC class II molecules outside of the antigen groove. *Cell* **62**:1115–1121.
- Fast, D. J., P. M. Schlievert, and R. D. Nelson. 1989. Toxic shock syndrome-associated staphylococcal and streptococcal pyrogenic toxins are potent inducers of tumor necrosis factor production. *Infect. Immun.* **57**:291–294.
- Ferretti, J. J., W. M. McShan, D. Ajdic, D. J. Savic, G. Savic, K. Lyon, C. Primeaux, S. Sezate, A. N. Suvorov, S. Kenton, H. S. Lai, S. P. Lin, Y. Qian, H. G. Jia, F. Z. Najjar, Q. Ren, H. Zhu, L. Song, J. White, X. Yuan, S. W. Clifton, B. A. Roe, and R. McLaughlin. 2001. Complete genome sequence of an M1 strain of *Streptococcus pyogenes*. *Proc. Natl. Acad. Sci. USA* **98**:4658–4663.
- Fleischer, B., and H. Schrezenmeier. 1988. T cell stimulation by staphylococcal enterotoxins. Clonally variable response and requirement for major histocompatibility complex class II molecules on accessory or target cells. *J. Exp. Med.* **167**:1697–1707.
- Fraser, J., V. Arcus, P. Kong, E. Baker, and T. Proft. 2000. Superantigens—powerful modifiers of the immune system. *Mol. Med. Today* **6**:125–132.
- Fraser, J. D. 1989. High-affinity binding of staphylococcal enterotoxins A and B to HLA-DR. *Nature* **339**:221–223.
- Harrington, D. J., I. C. Sutcliffe, and N. Chanter. 2002. The molecular basis of *Streptococcus equi* infection and disease. *Microb. Infect.* **4**:501–510.
- Herman, A., J. W. Kappler, P. Marrack, and A. M. Pullen. 1991. Superantigens: mechanism of T-cell stimulation and role in immune responses. *Annu. Rev. Immunol.* **9**:745–772.
- Hudson, K. R., H. Robinson, and J. D. Fraser. 1993. Two adjacent residues in Staphylococcal enterotoxins A and E determine T cell receptor V β specificity. *J. Exp. Med.* **177**:175–185.
- Hurley, J. M., R. Shimonkevitz, A. Hanagan, K. Enney, E. Boen, S. Malmstrom, B. L. Kotzin, and M. Matsumura. 1995. Identification of class II major histocompatibility complex and T cell receptor binding sites in the superantigen toxic shock syndrome toxin 1. *J. Exp. Med.* **181**:2229–2235.
- Ikebe, T., A. Wada, Y. Inagaki, K. Sugama, R. Suzuki, D. Tanaka, A. Tamari, Y. Fujinaga, Y. Abe, Y. Shimizu, and H. Watanabe. 2002. Dissemination of the phage-associated novel superantigen gene *speL* in recent invasive and noninvasive *Streptococcus pyogenes* M3/T3 isolates in Japan. *Infect. Immun.* **70**:3227–3233.
- Ito, Y., J. Abe, K. Yoshino, T. Takeda, and T. Kohsaka. 1995. Sequence analysis of the gene for a novel superantigen produced by *Yersinia pseudotuberculosis* and expression of the recombinant protein. *J. Immunol.* **154**: 5896–5906.
- Jarraud, S., M. A. Peyrat, A. Lim, A. Tristan, M. Bes, C. Mougel, J. Etienne, et al. 2001. egc, a highly prevalent operon of enterotoxin gene, forms a putative nursery of superantigens in *Staphylococcus aureus*. *J. Immunol.* **166**:669–677.
- Kotzin, B. L., D. Y. Leung, J. Kappler, and P. Marrack. 1993. Superantigens and their potential role in human disease. *Adv. Immunol.* **54**:99–166.
- Li, P. L., R. E. Tiedemann, S. L. Moffat, and J. D. Fraser. 1997. The superantigen streptococcal pyrogenic exotoxin C (SPE-C) exhibits a novel mode of action. *J. Exp. Med.* **186**:375–383.
- Li, Y., H. Li, N. Dimasi, J. K. McCormick, R. Martin, P. Schnuck, P. M. Schlievert, and R. A. Mariuzza. 2001. Crystal structure of a superantigen bound to the high-affinity, zinc-dependent site on MHC class II. *Immunity* **14**:93–104.
- Marrack, P., and J. Kappler. 1990. The staphylococcal enterotoxins and their relatives. *Science* **248**:705–711.
- Martin, D. R., L. M. Voss, S. J. Walker, and D. Lennon. 1994. Acute rheumatic fever in Auckland, New Zealand: spectrum of associated group A streptococci different from expected. *Pediatr. Infect. Dis. J.* **13**:264–269.
- Nielsen, H., J. Engelbrecht, S. Brunak, and G. von Heijne. 1997. Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. *Protein Eng.* **10**:1–6.
- Page, R. D. M. 1996. TREEVIEW: an application to display phylogenetic trees on personal computers. *Comput. Appl. Biosci.* **12**:357–358.
- Papageorgiou, A. C., and K. R. Acharya. 2000. Microbial superantigens: from structure to function. *Trends Microbiol.* **8**:369–375.
- Proft, T., V. L. Arcus, V. Handley, E. N. Baker, and J. D. Fraser. 2001. Immunological and biochemical characterization of streptococcal pyrogenic exotoxins I and J (SPE-I and SPE-J) from *Streptococcus pyogenes*. *J. Immunol.* **166**:6711–6719.
- Proft, T., S. L. Moffatt, C. J. Berkahn, and J. D. Fraser. 1999. Identification and characterization of novel superantigens from *Streptococcus pyogenes*. *J. Exp. Med.* **189**:89–101.
- Proft, T., S. L. Moffatt, K. D. Weller, A. Paterson, D. Martin, and J. D. Fraser. 2000. The streptococcal superantigen SMEZ exhibits wide allelic variation, mosaic structure, and significant antigenic variation. *J. Exp. Med.* **191**:1765–1776.
- Roussel, A., B. F. Anderson, H. M. Baker, J. D. Fraser, and E. N. Baker. 1997. Crystal structure of the streptococcal superantigen SPE-C: dimerization and zinc binding suggest a novel mode of interaction with MHC class II molecules. *Nat. Struct. Biol.* **4**:635–643.
- Seth, A., L. J. Stern, T. H. Ottenhoff, I. Engel, M. J. Owen, J. R. Lamb, R. D. Klausner, and D. C. Wiley. 1994. Binary and ternary complexes between T-cell receptor, class II MHC and superantigen in vitro. *Nature* **369**:324–327.
- Stevens, D. L. 2000. Streptococcal toxic shock syndrome associated with necrotizing fasciitis. *Annu. Rev. Med.* **51**:271–288.
- Sundberg, E. J., H. Li, A. S. Llera, J. K. McCormick, J. Tormo, P. M. Schlievert, K. Karjalainen, and R. A. Mariuzza. 2002. Structures of two streptococcal superantigens bound to TCR β chains reveal diversity in the architecture of T cell signaling complexes. *Structure* **10**:687–699.
- Thompson, J. D., D. G. Higgins, and T. J. Gibson. 1994. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment. *Nucleic Acids Res.* **22**:4673–4680.
- Walker, P. A., L. E. Leong, P. W. Ng, S. H. Tan, S. Waller, D. Murphy, and A. G. Porter. 1994. Efficient and rapid affinity purification of proteins using recombinant fusion proteases. *Bio/Technology* **12**:601–605.