

Osmolarity, a Key Environmental Signal Controlling Expression of Leptospiral Proteins LigA and LigB and the Extracellular Release of LigA

James Matsunaga,^{1,2*} Yolanda Sanchez,¹ Xiaoyi Xu,¹ and David A. Haake^{1,2}

Division of Infectious Diseases, Veterans Affairs Greater Los Angeles Healthcare System,¹ and Division of Infectious Diseases, Department of Medicine, UCLA David Geffen School of Medicine,² Los Angeles, California

Received 14 July 2004/Returned for modification 11 August 2004/Accepted 25 August 2004

The high-molecular-weight leptospiral immunoglobulin-like repeat (Lig) proteins are expressed only by virulent low-passage forms of pathogenic *Leptospira* species. We examined the effects of growth phase and environmental signals on the expression, surface exposure, and extracellular release of LigA and LigB. LigA was lost from stationary-phase cells, while LigB expression was maintained. The loss of cell-associated LigA correlated with selective release of a lower-molecular-weight form of LigA into the culture supernatant, while LigB and the outer membrane lipoprotein LipL41 remained associated with cells. Addition of tissue culture medium to leptospiral culture medium induced LigA and LigB expression and caused a substantial increase in released LigA. The sodium chloride component of tissue culture medium was primarily responsible for the enhanced release of LigA. Addition of sodium chloride, potassium chloride, or sodium sulfate to leptospiral medium to physiological osmolarity caused the induction of both cell-associated LigA and LigB, indicating that osmolarity regulates the expression of Lig proteins. Osmotic induction of Lig expression also resulted in enhanced release of LigA and increased surface exposure of LigB, as determined by surface immunofluorescence. Osmolarity appears to be a key environmental signal that controls the expression of LigA and LigB.

Infection of human beings by pathogenic members of the spirochete genus *Leptospira* results in a potentially fatal infection characterized by jaundice, renal failure, and/or pulmonary hemorrhage (3). The potentially severe nature of acute leptospirosis in humans contrasts with chronic infection in reservoir host animals such as *Rattus norvegicus*, which exhibits lifelong renal tubular carriage and urinary shedding (3). Water is an important vehicle for transmission to new mammalian hosts. Transmission to humans via contaminated water typically occurs in developed countries as a result of recreational or occupational exposures and in developing countries in settings where heavy rainfall results in urban flooding (19, 26). Leptospire enter the host by penetrating mucous membranes or broken skin.

Little is known about the environmental signals that leptospire use to respond to the variety of environmental conditions they encounter inside and outside of the mammalian host. Pathogenic bacteria control the expression of virulence factors in response to environmental cues (25). The presence of 79 two-component regulatory systems and 11 extracytoplasmic function (ECF) sigma factors in the genomes of *L. interrogans* serovar Lai and *L. interrogans* serovar Copenhageni suggest that pathogenic *Leptospira* spp. are capable of responding to a diverse array of environmental signals (28, 30). Growth phase and temperature affect the levels of several leptospiral membrane proteins. For example, expression of the outer membrane lipoprotein LipL36 is decreased during stationary phase (11), while expression of the peripheral outer membrane

protein P31_{LipL45} (Qlp42) is increased during stationary phase (23). Similarly, growth at 37°C instead of the standard 30°C cultivation temperature causes a reduction in the expression of LipL36 and an increase in that of P31_{LipL45} (27). Consistent with these observations, leptospire residing in the kidney tubules of hamsters fail to express LipL36 yet express P31_{LipL45} (2, 23). A global examination of leptospiral outer membrane proteins by two-dimensional electrophoresis revealed several additional proteins whose expression is affected by iron availability and temperature, including pL24 and pL50 (9). The mechanisms by which the expression of these proteins are regulated remain to be determined.

Efforts to identify leptospiral proteins that are expressed during infection led to the discovery of genes encoding the leptospiral immunoglobulin (Ig)-like repeat (Lig) family of proteins (22,29). Lig proteins contain imperfect tandem repeats of a ~90-amino-acid residue sequence that is predicted to form an Ig-like fold (22, 29). In *L. interrogans* strain Fiocruz L1-130 and *L. kirschneri* strain RM52, LigA contains 13 copies of the Ig-like sequence and LigB contains 12 copies followed by a unique C-terminal domain. Both LigA and LigB contain a lipobox, and LigA was lipidated by palmitate when expressed in *E. coli* (20). LigB has been shown to be surface exposed; whether LigA is surface exposed is unknown (22). An intact *ligC* gene is present in the genome of *L. interrogans* serovar Lai, but *ligC* is likely to be a pseudogene in *L. interrogans* serovar Copenhageni and *L. kirschneri* serovar Grippotyphosa (22). Expression of Lig proteins is lost during culture attenuation of *L. kirschneri* and *L. interrogans*, suggesting that these proteins are associated with virulence (22). It is thought that the Lig proteins may function as adhesins, given the similarity of their domain organization to those of the well-characterized ad-

* Corresponding author. Mailing address: VA Greater Los Angeles Healthcare System, B113/R225, Los Angeles, CA 90073. Phone: (310) 478-3711 ext. 42144. Fax: (310) 268-4142. E-mail: jamesm@ucla.edu.

hesins invasins and intimin, found in *Yersinia* species and enteropathogenic *Escherichia coli*, respectively (12, 21).

Several lines of evidence indicate that expression of Lig proteins is increased during infection of the mammalian host. First, LigA could not be detected in cultivated *L. interrogans* strain kennewicki but was detected by immunohistochemistry in leptospires residing in the kidneys of infected hamsters (29). Second, rats immunized with killed *L. interrogans* grown in vitro failed to produce antibody to Lig, most probably a result of insufficient Lig expression (22). In contrast, rats infected with *L. interrogans* produced Lig antibody, consistent with a higher level of Lig expression during infection than during in vitro growth. In this study, we sought to identify environmental conditions that affect expression of Lig proteins in *L. kirschneri* RM52 and *L. interrogans* Fiocruz L1-130. We found that the expression of Lig proteins is regulated by osmolarity. We also demonstrate that both *Leptospira* strains release LigA into the extracellular fluid and that the release of LigA is enhanced by salt.

MATERIALS AND METHODS

Bacterial strains and cultivation. *L. kirschneri* serovar Grippotyphosa strain RM52 was isolated during an outbreak of porcine abortion in the United States (33). *L. interrogans* serovar Copenhageni strain L 1-130 is a human blood isolate obtained during an outbreak of leptospirosis in Salvador, Brazil (22). All experiments were performed with virulent, low-passage forms of these strains obtained by infection and reisolation from Golden Syrian hamsters. The spirochetes were maintained in Ellinghausen-McCullough-Johnson-Harris (EMJH) medium (pH 7.2) supplemented with 1% rabbit serum and 100 μ g of 5-fluorouracil (Sigma, St. Louis, Mo.) per ml at 30°C (17). Albumin was purchased from Intergen (Purchase, N.Y.; catalog no. 31-003-3) and from Sigma (catalog no. A7906) for cultivation of *L. kirschneri* RM52 and *L. interrogans* Fiocruz L1-130, respectively. Modified Eagle medium (MEM), which was purchased from the American Type Culture Collection (Manassas, Va.), contains Earle's salts and both essential and nonessential amino acids. Fetal bovine serum (FBS) was purchased from Atlanta Biologicals (Norcross, Ga.).

Plasmid DNA. The portion of the *flaA1* gene beginning from the codon following the segment encoding the signal peptide was amplified by PCR with ExTaq DNA polymerase (Takara, Madison, Wis.) using the forward primer 5'-GACTCGAGAATGGACAAAACATCATCAAAGGCAAAC-3' and the reverse primer 5'-GACTCGAGAATGGACAAAACATCATCAAAGGCAAAC-3'. The primers included an XhoI site and a HindIII site, respectively (underlined), and *L. interrogans* Fiocruz L1-130 genomic DNA was used as template for PCR. The amplified *flaA1* gene fragment was digested with XhoI and HindIII and ligated to the His₆ vector pRSET A (Invitrogen, Carlsbad, Calif.) with T4 DNA ligase to generate the plasmid pRSETA-FlaA1. Restriction and modifying enzymes were purchased from New England Biolabs (Beverly, Mass.).

Antisera. LigA-B and LipL41 antisera were prepared by immunizing New Zealand White rabbits (Harlan, Indianapolis, Ind.) with His₆-Lig and His₆-LipL41 recombinant proteins as described previously (22, 31). LigA-B antiserum was raised against amino acid residues 342 to 1224 of LigA, which includes a region shared with LigB. LigA antiserum raised against the last 289 amino acid residues of LigA (residues 936 to 1224) does not react with LigB in immunoblots. Likewise, the LigB-specific antiserum was raised against the C-terminal domain of LigB (residues 1113 to 1886). FlaA1 antiserum was generated as follows. Plasmid pRSETA-FlaA1 was transformed into *Escherichia coli* BLR(DE3)/pLysS (EMD Biosciences, Madison, Wis.), and expression of the His₆-FlaA1 fusion protein was induced with 0.5 mM isopropyl- β -D-thiogalactoside. The recombinant protein was purified by affinity chromatography with Ni²⁺-nitrilotriacetate agarose resin (Qiagen, Valencia, Calif.) and loaded onto a preparative sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) gel (12% polyacrylamide). Following electrophoresis, the band containing approximately 150 μ g of His₆-FlaA1 protein was cut out of the gel, desiccated, ground into powder, and mixed with 1 ml of sterile water. The suspension was mixed with 1 ml of complete Freund's adjuvant (Sigma) and inoculated subcutaneously and intramuscularly into a New Zealand White male rabbit. Additional immunizations with approximately 150 μ g of His₆-FlaA1 in 1 ml of sterile water mixed with 1 ml of incomplete Freund's adjuvant were given 4 weeks and 8 weeks following

the primary immunization. The rabbit was bled 10 weeks after the primary immunization. The immunization protocol was approved by the West Los Angeles Veterans Affairs Animal Research Committee.

Immunoblot analysis. Culture supernatant fluid was collected by centrifugation of 10⁹ leptospires for 4 min at 9,000 \times g in a Beckman (Fullerton, Calif.) Coulter Microfuge 18 centrifuge followed by collection of the supernatant fluid, which was subsequently examined for the presence of Lig and other leptospiral proteins by immunoprecipitation (see below). The cell pellet was washed once in phosphate-buffered saline (PBS)-5 mM MgCl₂ and resuspended in 100 μ l of final sample buffer consisting of 50 mM Tris-HCl (pH 6.8), 100 mM dithiothreitol, 2% sodium dodecyl sulfate, 0.25 mM phenylmethylsulfonyl fluoride, and 0.1% bromophenol blue in 20% glycerol and boiled for 3 min. Proteins were separated in 10 or 12% PAGER Gold precast Tris-glycine gels (Cambrex, Walkersville, Md.). Following electrophoresis, the material in each gel was transferred electrophoretically to a polyvinylidene difluoride membrane (Millipore, Bedford, Mass.). The membranes were incubated in blocking solution (5% skim milk in PBS-0.1% Tween 20 [PBS-T]) for 30 min and then incubated with rabbit serum (diluted 1:2000 to 1:10,000) for 30 min and washed three times with PBS-T. They were then incubated with donkey anti-rabbit antibody (Amersham Biosciences, Piscataway, N.J.) or protein A-horseradish peroxidase conjugate (Amersham) at a dilution of 1:5,000 or 1:2,000, respectively, in blocking solution for 30 min and again washed three times in PBS-T. The membranes were developed with the ECL Western blot detection system (Amersham), and the bands were visualized with Hyperfilm (Amersham). Band intensities of Lig proteins were quantitated with NIH Image 1.62 (National Institutes of Health; <http://rsb.info.nih.gov/nih-image>) and normalized against the intensity of the LipL41 band in the same lane.

Immunoprecipitation. Culture supernatant (995 μ l) was mixed with 5 μ l of rabbit serum and left on ice overnight. A 25- μ l volume of EZview Red protein A affinity gel (Sigma) was then added, and the mixture was placed on an orbital mixer at 4°C for 2 h. The immune complex bound to the protein A was recovered by centrifugation for 7 s, washed twice with 800 μ l of 10 mM Tris-HCl-0.4 M NaCl (pH 8.0), and finally washed once with 800 μ l of 10 mM Tris-HCl (pH 8.0). The pellet was resuspended in 50 to 100 μ l of final sample buffer. Samples were boiled for 3 min and centrifuged for 7 s. A 10- μ l volume was used for immunoblot analysis.

RT-PCR. *L. kirschneri* strain RM52 (5×10^9 cells) was transferred into an Erlenmeyer flask, chilled in a dry ice-ethanol bath, and centrifuged at 10,000 rpm for 10 min in a Sorvall SM-24 rotor. RNA was extracted from the bacteria with hot phenol, precipitated with ethanol, and resuspended in 50 μ l of diethylpyrocarbonate-treated water purchased from ICN (5). RNA (4 μ g) was treated with 2 U of Turbo-DNase in a final volume of 40 μ l for 30 min at 37°C as directed by the manufacturer (Ambion, Austin, Tex.).

A 2- μ g portion of DNase-treated RNA was hybridized to random hexamer primers (Promega, Madison, Wis.), and cDNA was synthesized with Omniscript reverse transcriptase (RT) as specified by the manufacturer (Qiagen). The cDNA (representing 50 ng of RNA per reaction) was amplified with *Taq* DNA polymerase (Qiagen) with gene-specific primer pairs. The *ligA*, *ligB*, and *ligC* primers have been described previously (22). The sequences of the *lipL41* forward and reverse primers are 5'-TCGGTGAAGGTTCCAGTTTTATTGAT-3' and 5'-TACTTCTCCGGTTTCTACTTTGATGA-3', respectively. Those of the *ligC* primers that amplify a segment upstream of the frameshift in the pseudogene are 5'-TATTCTTTTTTCTAACATTACAGCCTAT-3' (forward) and 5'-AAGTCCGAACTAAACCTGTGGTGT-3' (reverse).

Immunofluorescence. Sodium azide was added to a culture of *L. interrogans* Fiocruz L1-130 to a final concentration of 0.5% (wt/vol), and the nucleic acid fluorescent stain SYTO 83 (Molecular Probes, Eugene, Oreg.) was added to the culture at a 1:1,000 dilution. After incubation for 1 hr at 30°C, the bacteria were collected by centrifugation at 1,000 \times g for 10 min in a microcentrifuge, resuspended in 50 mM carbonate buffer (pH 9.6), placed in two-well glass slide chambers (0.5×10^9 to 1.0×10^9 leptospires per well) (Nunc, Rochester, N.Y.), and incubated at 30°C for 2 h. Next, the bacteria were treated with blocking solution (0.5% bovine serum albumin in PBS-1% Tween 20) for 2 hrs at 30°C. They were then fixed with 4% paraformaldehyde for 1 h at 30°C, washed three times with PBS, treated with a 1:250 dilution of rabbit antiserum in blocking buffer for 1 h at 30°C, washed three times with PBS, treated with a 1:1,000 dilution of Alexa Fluor 488-labeled goat anti-rabbit IgG (Molecular Probes; catalog no. A-11034) in blocking solution for 1 hr at 30°C, washed three times with PBS, and washed twice with water. Finally, mounting medium (ProLong antifade kit; Molecular Probes) and a coverslip were added, and the slide was left overnight. Staining was visualized by confocal microscopy with an LSM 510 microscope (Carl Zeiss, Inc., Jena, Germany).




FIG. 1. Growth phase-dependent expression of Lig proteins. (A) *L. kirschneri* RM52 was inoculated into EMJH and incubated at 30°C. Samples of the culture were collected daily, and the bacterial densities were determined. (B) Leptospires from each sample collected for the experiment in panel A were loaded onto an SDS-PAGE gel (12% polyacrylamide), and Lig and LipL41 expression was examined by immunoblot analysis. The density of the culture (number of motile bacteria per milliliter) is shown above the panel; 10⁸ bacteria were loaded in each lane. The first lane contains proteins from leptospires collected from the stationary-phase culture that was the source of the starting material for the culture whose growth is plotted in panel A.

RESULTS

LigA is released into the culture medium. In previous studies, we showed that the levels of several leptospiral proteins change with growth phase (see Introduction for details). We therefore wondered whether the expression of Lig proteins also varies with the growth phase. To examine this issue, we initiated a culture of *L. kirschneri* strain RM52 in EMJH at 30°C from a stationary-phase culture and incubated the culture until it reached stationary phase. During the incubation, samples of the culture were collected at various times (Fig. 1A). The leptospires collected at each time point were examined by immunoblot analysis for expression of LigA and LigB (Fig. 1B). Lysates from equal numbers of leptospires were loaded into each lane. The membrane was probed with LigA-B and LipL41 antisera, the latter included as a control for loading. LipL41 levels do not change throughout the growth curve (11). The immunoblot shows that the LigA levels were greatly reduced during stationary phase whereas LigB was easily detected at all phases of growth (Fig. 1B).

Since the DNA sequence in the promoter regions of *ligA* and *ligB* are identical (22), we wondered whether the difference in expression of the two proteins could have resulted from alternative fates of the two proteins during stationary phase. One possible explanation for the disappearance of LigA during entry into stationary phase is its selective release into the culture medium. To examine this possibility, experiments were performed in which, starting from late log phase, samples were collected at various time points from cultures of *L. kirschneri* RM52. Leptospires were collected by centrifugation, and both the bacteria and culture supernatant fluid were obtained. To avoid distortion of the bands in the SDS-PAGE gel due to the high concentration of albumin in EMJH, immunoprecipitation of the culture supernatant was performed with antiserum specific for the shared regions of LigA and LigB. The immunoprecipitated material was examined by immunoblotting, revealing a species that migrated slightly faster than cell-associated LigA, which we designated LigA' (Fig. 1C, lanes 4 to 6). In contrast, neither LigB nor LipL41 was detected in the culture supernatant (lanes 4 to 6; data not shown). *L. interrogans* strain Fiocruz L1-130 also released LigA' into the culture supernatant (see Fig. 2C, lane 2).

To verify that LigA' represented a smaller form of LigA and not LigB, the immunoprecipitated material was probed separately with LigA- and LigB-specific antisera. LigA' reacted with LigA antiserum but not with LigB antiserum, indicating that LigA' is derived from LigA (data not shown).

Lig expression is induced by MEM. Immunoblots with sera from infected humans and rats indicate that Lig proteins are expressed during infection of the mammalian host (22). For

(C) *L. kirschneri* RM52 was grown in EMJH to late log phase, and bacteria were collected on three consecutive days as the culture went into saturation. Leptospires were collected by centrifugation, and the supernatant fluid was subjected to immunoprecipitation with anti-Lig serum. The leptospires were loaded onto lanes 1 to 3 of a 4 to 20% gradient SDS-PAGE gel; Lig protein immunoprecipitated from the culture supernatant (supe) was loaded onto lanes 4 to 6. A total of 10⁸ leptospires were loaded in each lane. *, reaction of the protein A-HRP conjugate with the immunoblobulin heavy chain.




FIG. 2. Induction of Lig expression and release by tissue culture medium. (A) *L. kirschneri* RM52 was grown to late log phase. Samples of the culture were mixed with an equal volume of EMJH or tissue culture medium, which consisted of MEM with 10% FBS. The leptospire was incubated for 24 h, and the leptospire and culture supernatant were examined for Lig protein by immunoblot analysis. c, cells;

this reason, we examined whether conditions likely to be encountered within the mammalian host affect Lig protein expression. *L. kirschneri* strain RM52 was incubated overnight in MEM-10% FBS, which models in vivo conditions. Leptospire incubated overnight in MEM-10% FBS instead of the leptospiral medium EMJH lost motility and appeared to degenerate, as determined by dark-field microscopy. However, *L. kirschneri* RM52 incubated overnight in a 50:50 mixture of EMJH and MEM-10% FBS retained motility and morphology. Bacteria and culture supernatant were collected at the end of the incubation period and examined for Lig protein expression by immunoblot analysis. MEM-10% FBS, when mixed with EMJH, enhanced the levels of LigA and LigB by 10- to 15-fold and enhanced the level of LigA' by 50-fold (Fig. 2A, compare lanes 2 and 4). Although addition of FBS to EMJH partially induced Lig expression (Fig. 2B, lanes 7 and 8), Lig expression continued to be fully induced when FBS was omitted from the MEM-EMJH mixture (lanes 5 and 6). In four independent experiments, MEM induction was 4- to 24-fold for cell-associated LigA and LigB and 18- to 53-fold for LigA'. MEM also enhanced LigA, LigB, and LigA' levels in *L. interrogans* strain Fiocruz L1-130 (Fig. 2C), demonstrating that regulation of Lig expression by MEM is not limited to *L. kirschneri* strain RM52. To eliminate the alternative possibility that mixing in the MEM diluted an inhibitory substance in EMJH, *L. kirschneri* RM52 was grown in EMJH that had been diluted twofold with water instead of MEM. Dilution of EMJH had no effect on Lig protein levels or release in comparison to growth in full-strength EMJH (data not shown), indicating that the increase in Lig expression resulted from an inducing ingredient in MEM.

The transcript levels of *lig* were examined by RT-PCR. The *ligA* and *ligB* transcripts were barely detectable in *L. kirschneri* RM52 grown in EMJH (Fig. 3, lanes 2 and 6). In contrast, *ligA* and *ligB* transcripts were easily detected when *L. kirschneri* RM52 was grown in a 50:50 mixture of EMJH and MEM (lanes 4 and 8). These results demonstrate that the presence of MEM elevates *ligA* and *ligB* transcript levels. Primers that hybridize downstream (lanes 10 and 12) or upstream (lanes 18 and 20) of the frameshift in *ligC* could not detect transcript from the *ligC* pseudogene, whether or not MEM was present. In all cases, the lack of reverse transcriptase in the reaction mixture prevented the formation of a PCR product, indicating that contaminating DNA in the RNA preparations was efficiently removed. The positive control *lipL41* transcript was detectable whether or not MEM was present in the culture medium (lanes 14 and 16).

su, culture supernatant. (B) *L. kirschneri* RM52 was grown as described for panel A, and samples of the culture were mixed with an equal volume of EMJH (lanes 1 and 2), MEM with 10% FBS (lanes 3 and 4), MEM (lanes 5 and 6), or 10% FBS (lanes 7 and 8). The cultures were incubated for 24 h. The leptospire and culture supernatant were examined for Lig protein by immunoblot analysis as for the experiment in panel A. (C) *L. interrogans* Fiocruz L1-130 was grown to late log phase. The bacteria were mixed with an equal volume of EMJH or MEM and incubated for 24 h. The leptospire and culture supernatant were examined for Lig protein by immunoblot analysis as for the experiment in panel A.




FIG. 3. Regulation of *lig* transcript levels by MEM. A culture of *L. kirschneri* RM52 at late log phase in EMJH was mixed with an equal volume of EMJH or MEM and incubated for 24 h. RNA was extracted from each culture and was examined for the presence of *lig* and *lipL41* transcripts by RT-PCR. Lane 1, 1-kb ladder from New England Biolabs; +, RT present; -, RT omitted.

Lig expression is regulated by osmolarity. We were interested in determining the ingredient in MEM that induced LigA' release. Since MEM is a completely defined medium and FBS was not required for full induction of Lig protein expression, it was possible to determine which MEM component was responsible for the induction effect by systematically adding the individual components of MEM to EMJH at the concentrations found in MEM. We focused on the components that were at a higher concentration in MEM than in EMJH. Addition of the MEM vitamins, amino acids, sodium pyruvate, or glucose had little or no effect on LigA' release; sodium bicarbonate had a slight effect (data not shown). On the other hand, a mixture of the salts calcium chloride, magnesium sulfate, and sodium chloride added to EMJH at the concentrations found in MEM substantially enhanced LigA' release (Fig. 4, lane 6). When the salts were examined individually, sodium chloride alone accounted for the effect on LigA' release (lanes, 8, 10, and 12). Cell-associated LigA and LigB levels increased slightly in EMJH supplemented with sodium chloride (lane 11).

Because sodium chloride accounts for the majority of the osmolarity of MEM, we wondered whether Lig release was regulated by the osmolarity of the medium rather than the sodium or chloride ions themselves. To test this possibility, *L. interrogans* strain Fiocruz L1-130 was collected by centrifugation, resuspended in EMJH without any additions or in EMJH supplemented with 120 mM NaCl, 120 mM KCl, or 80 mM Na₂SO₄, and incubated overnight. The osmolarity in the cultures supplemented with salt approximated physiological conditions (~300 mosmol per liter), whereas the osmolarity of EMJH was about 70 mosmol. The immunoblot shows that all three salts induced Lig protein expression by 16- to 21-fold and induced LigA' release by 200- to 300-fold (Fig. 5), indicating that Lig expression responds to the osmolarity of the growth medium and not to a specific ion. Similar results were obtained with *L. kirschneri* RM52 (data not shown). In four independent experiments, induction by sodium chloride was 5- to 30-fold for cell-associated LigA and LigB and 50- to 320-fold for extracellular LigA'.

LigA' release appears to be more sensitive to the presence of sodium chloride than is the release of cell-associated Lig




FIG. 4. Induction of Lig expression and release by sodium chloride. A culture of *L. kirschneri* RM52 grown to late log phase in EMJH was mixed with an equal volume of EMJH (lanes 1 and 2); MEM (lanes 3 and 4); a solution of 0.1 mM MgSO₄, 1.8 mM CaCl₂, and 116 mM NaCl (lanes 5 and 6); 0.1 mM MgSO₄ (lanes 7 and 8); 1.8 mM CaCl₂ (lanes 9 and 10); or 116 mM NaCl (lanes 11 and 12). The cultures were incubated for 24 h and examined for Lig expression and release following separation of proteins on an SDS-PAGE gel (10% polyacrylamide). c, cells; s, culture supernatant fluid.

proteins (Fig. 4, lanes 1 and 2 and lanes 11 and 12). To confirm this observation, Lig proteins were examined following overnight incubation of *L. interrogans* Fiocruz L1-130 in EMJH containing different concentrations of sodium chloride. When 40 mM sodium chloride was added to EMJH, a nine-fold increase in the release of LigA' was observed in contrast to the one- to twofold increase in the release of cell-associated Lig proteins (Fig. 6, compare lanes 1 and 2 with lanes 3 and 4). When 80 or 120 mM sodium chloride was added, a 6-fold




FIG. 5. Induction of Lig expression and release is independent of the salt responsible for osmolarity. *L. interrogans* Fiocruz L1-130 grown in EMJH was collected by centrifugation and resuspended in EMJH (lanes 1 and 2) or EMJH containing an additional 120 mM NaCl (lanes 3 and 4), 120 mM KCl (lanes 5 and 6), or 80 mM Na₂SO₄ (lanes 7 and 8). The cultures were incubated for 24 h and examined for Lig expression and release following separation of proteins on an SDS-PAGE gel (10% polyacrylamide). c, cells; s, culture supernatant fluid.




FIG. 6. Sensitivity of Lig expression and release to sodium chloride concentrations. *L. interrogans* Fiocruz L1-130 grown in EMJH was collected by centrifugation and resuspended in EMJH (lanes 1 and 2) or EMJH containing an additional 40 mM NaCl (lanes 3 and 4), 80 mM NaCl (lanes 5 and 6), or 120 mM NaCl (lanes 7 and 8). The cultures were incubated for 24 h and examined for Lig expression and release following separation of proteins on an SDS-PAGE gel (10% polyacrylamide) gel. c, cells; s, culture supernatant fluid.

induction of cell-associated LigA and LigB was observed and release of LigA' was enhanced 30- to 50-fold (lanes 5 to 8). Incubation of *L. interrogans* at higher concentrations of sodium chloride resulted in the appearance of nonmotile, shriveled leptospires when observed by dark-field microscopy.

Sodium chloride increases the surface exposure of LigB. Increased concentrations of sodium chloride may increase the availability of Lig proteins for direct interaction with the host by enhancing their surface exposure. Indirect immunofluorescence was used to determine whether the induction of Lig expression leads to an increase in the amounts of surface-exposed Lig proteins. LipL41, which has been shown to be surface exposed by surface immunoprecipitation, was used as a

positive control (31). FlaA1 antiserum, which reacts with the periplasmic flagellar sheath, was used as a negative control. As expected, *L. kirschneri* RM52 incubated in EMJH and EMJH containing extra sodium chloride reacted with LipL41 antiserum (Fig. 7A and D) and failed to react with FlaA1 antiserum (Fig. 7B and E). LigA-B antiserum reacted weakly with *L. kirschneri* incubated in EMJH (Fig. 7C). In contrast, strong reactivity with the LigA-B antiserum was observed when Lig expression was induced with sodium chloride (Fig. 7F).

The effect of osmolarity on surface exposure of LigA and LigB was also examined with specific antiserum to each protein. When *L. interrogans* Fiocruz L1-130 was incubated in EMJH, weak reactivity with the LigB-specific antiserum was observed (Fig. 8A) (22). Incubation of *L. interrogans* in EMJH containing extra sodium chloride resulted in an increase in the signal observed (Fig. 8C). In contrast, FlaA1 antiserum failed to react with the spirochetes, regardless of the growth conditions (Fig. 8B and D). Treatment of the leptospires with methanol caused reactivity with the FlaA1 antiserum, indicating that the FlaA1 serum was functional (data not shown). *L. interrogans* reacted with the LipL41 antiserum, as expected (data not shown). Uninduced and salt-induced leptospires failed to react with LigA-specific antiserum, even after the bacteria were fixed with methanol (data not shown).

DISCUSSION

We have previously shown that expression of LigA and LigB is associated with virulence (22). In this study, we investigated the environmental signals controlling Lig protein expression. Since the expression of the outer membrane proteins LipL36 and P31_{Lip45} is altered in stationary phase (11, 23), we first examined whether Lig protein expression was controlled by growth phase. While LigB was consistently expressed at all stages of leptospiral growth, LigA expression was dramatically reduced in stationary phase (Fig. 1B). The loss of cell-associated LigA occurred concomitantly with continued release of LigA into the culture medium (Fig. 1C). Therefore, LigA joins the hemolysins as the only characterized leptospiral proteins known to be released extracellularly (35).




FIG. 7. Salt-induced surface exposure of Lig proteins demonstrated by indirect immunofluorescence. *L. kirschneri* RM52 grown in EMJH was collected by centrifugation and resuspended in EMJH (A to C) or EMJH containing an additional 120 mM NaCl (D to F). After a 24-h incubation period, leptospires were stained with SYTO 83 (red) and treated with LipL41 (A and D), FlaA1 (B and E), or LigA-B (C and F) antiserum. Antibody bound to the surface of the leptospires was detected with Alexa-labeled anti-rabbit IgG (green).




FIG. 8. Salt-induced surface exposure of LigB demonstrated by indirect immunofluorescence. *L. interrogans* Fiocruz L1-130 grown in EMJH was collected by centrifugation and resuspended in EMJH (A and B) or EMJH containing an additional 120 mM NaCl (C and D). After a 24-h incubation period, leptospire were stained with SYTO 83 (red) and examined for surface exposure of LigB (A and C) and the periplasmic flagellar protein FlaA1 (B and D) by indirect immunofluorescence. Antibody bound to the surface of the leptospire was detected with Alexa-labeled anti-rabbit IgG (green).

Careful examination of immunoblots revealed that LigA immunoprecipitated from culture supernatants was slightly smaller than cell-associated LigA (Fig. 1C). We have designated the released form of LigA as LigA'. The *ligA* gene encodes a lipoprotein signal peptide, and LigA expressed in *E. coli* is lipidated (20), indicating that there is *sec*-dependent export of LigA across the cytoplasmic membrane followed by lipidation of the lipobox cysteine by leptospiral lipoprotein signal peptidase. It is possible that once LigA reaches the surface of the spirochete as a lipoprotein, it is subject to proteolytic removal of its lipid anchor, resulting in the formation of LigA', which can be recovered from the culture supernatant.

We discovered that tissue culture medium is a valuable *in vitro* model for examining the expression of Lig expression under conditions that may be encountered by the spirochete during infection of the mammalian host. MEM with serum did not sustain the overnight viability of leptospire, most probably a consequence of the strict requirement of long-chain fatty acids as a carbon source (16). However, incubation in a 50:50 mix of leptospiral culture medium and MEM not only allowed leptospire to remain viable but also resulted in an increase in expression of both of cell-associated LigA and LigB and of LigA' released by *L. kirschneri* strain RM52 (Fig. 2). MEM also caused an increase in the release of Lig proteins by *L. interrogans* strain L1-130 (Fig. 2C). Tissue culture media have also been used to examine the regulated secretion of virulence gene products in enteropathogenic *E. coli* (18). The discovery that incubation in MEM caused a dramatic increase in Lig expression facilitated the identification of sodium chloride as the primary molecular signal that controls *lig* expression *in vitro*. Control of Lig expression and release could also be

achieved with KCl and Na₂SO₄, indicating that sodium chloride was acting through its effect on osmolarity (Fig. 5). These data demonstrate that osmolarity is an important environmental cue that determines the level of cell-associated Lig proteins. Lig protein expression regulation is likely to be complex, and additional regulatory factors may yet be discovered. Furthermore, the results described here may not be completely relevant to all pathogenic *Leptospira* spp. For example, *L. interrogans* serovar Lai lacks a homolog of LigA. On the other hand, LigA has been found in *L. interrogans* serovar Pomona (29) and two forms of LigA have been found in *L. interrogans* serovar Manilae (20).

The increase in osmolarity experienced by pathogenic *Leptospira* spp. entering a mammalian host from the freshwater environment would be a particularly convenient signal for pathogenic *Leptospira* species to up-regulate the expression of virulence genes required during the early stages of infection of the host. Lig antibody is detected during the acute phase of leptospirosis (J. H. R. Croda and A. I. Ko, personal communication), suggesting that Lig is expressed early during infection. The putative adhesin activity of Lig proteins may be important in the dissemination of leptospire from the circulation to different organs. Osmolarity regulates the availability of LigA and LigB to directly interact with the host environment by enhancing release and surface exposure, respectively. We were unable to determine the surface exposure of LigA because of the poor reactivity of the LigA-specific antiserum in the immunofluorescence assay, even after the leptospire were fixed with methanol.

The osmotic control of Lig protein expression appears to involve transcriptional regulation. The increased Lig expres-

sion resulting from incubation in MEM was associated with increased transcript levels of the *ligA* and *ligB* genes, as demonstrated by RT-PCR (Fig. 3). Candidate proteins for regulation of *lig* gene transcription would include 1 or more of the 79 genes encoding two-component sensor histidine kinase response regulator proteins or 11 ECF-type sigma factors recently identified by leptospiral genome sequencing (28, 30). Leptospiral genomes encode homologs of the ProP and Kdp transporters (8, 13) and the two-component system KdpDE (34), suggesting that pathogenic leptospires are equipped with osmosensor and osmoregulator proteins. Changes in gene expression by osmolarity in other spirochetes have not been examined. In this context, it is worth noting that resistance of *Borrelia burgdorferi* at stationary phase to high concentrations of sodium chloride is reduced in an *rpoS* mutant, suggesting the presence of a borrelial RpoS regulon that is regulated by environmental sodium chloride (10). The *rpoS* gene is not present in the genome of either sequenced strain of *L. interrogans*, indicating that the mechanism of osmotic regulation of gene expression in *Leptospira* spp. is likely to differ from the RpoS-based mechanism examined extensively in *E. coli* (15).

The data presented here do not address the role of LigA' release or even whether LigA' is released during infection of the mammalian host or instead remains cell associated. A recent study examined the importance of release of the adhesin filamentous hemagglutinin (FHA) on *B. pertussis* pathogenesis in a mouse model (6). Following the delivery of the FHA proprotein to the surface of *B. pertussis*, cleavage by the protease SphB1 releases FHA from the bacterium (7). Disruption of the gene that encodes the SphB1 protease results in enhanced attachment of *B. pertussis* to monolayers in vitro relative to the strain with the wild-type gene. Nevertheless, colonization of mouse lungs following intranasal inoculation of the mutant bacteria was diminished, demonstrating that FHA secretion was essential for pathogenesis in the mouse model (6). It was proposed that secretion of the adhesin permits dissemination of the bacteria from the initial site of colonization into deeper tissues. A second possibility is that secreted LigA' stimulates host cell signaling pathways that are advantageous to leptospiral infection. An adhesin of the intracellular pathogen *Listeria monocytogenes*, InlB, which is found in both cell-associated and extracellular forms, stimulates phagocytosis of the bacteria by nonphagocytic cells by activating host signaling cascades. Furthermore, InlB stimulates additional host intracellular signaling pathways that possibly enhance the survival of the host cell that harbors the bacteria (4). Third, released LigA' may have immunomodulatory properties. FHA was shown to initiate apoptosis of human macrophages in vitro and in another study to suppress interleukin-12 production by macrophages (1, 24). Other examples of high-molecular-weight adhesins that are released include the Hap and HMW adhesins of *Haemophilus influenzae* (14, 32). Additional studies are necessary to address the role of Lig proteins in the interaction of pathogenic leptospires with the mammalian host.

ACKNOWLEDGMENTS

This work was supported by VA Medical Research Funds (to J.M. and D.A.H.) and Public Health Service grant AI-34431 (to D.A.H.) from the National Institute of Allergy and Infectious Diseases.

We thank George Sachs and David Scott for providing access to the confocal microscope and assistance in its use.

REFERENCES

- Abramson, T., H. Kedem, and D. A. Relman. 2001. Proinflammatory and proapoptotic activities associated with *Bordetella pertussis* filamentous hemagglutinin. *Infect. Immun.* **69**:2650–2658.
- Barnett, J. K., D. Barnett, C. A. Bolin, T. A. Summers, E. A. Wagar, N. F. Chevillat, R. A. Hartskeerl, and D. A. Haake. 1999. Expression and distribution of leptospiral outer membrane components during renal infection of hamsters. *Infect. Immun.* **67**:853–861.
- Bharti, A. R., J. E. Nally, J. N. Ricaldi, M. A. Matthias, M. M. Diaz, M. A. Lovett, P. N. Levett, R. H. Gilman, M. R. Willig, E. Gotuzzo, and J. M. Vinetz. 2003. Leptospirosis: a zoonotic disease of global importance. *Lancet Infect. Dis.* **3**:757–771.
- Bierne, H., and P. Cossart. 2002. InlB, a surface protein of *Listeria monocytogenes* that behaves as an invasin and a growth factor. *J. Cell Sci.* **115**:3357–3367.
- Case, C. C., E. L. Simons, and R. W. Simons. 1990. The *IS10* transposase mRNA is destabilized during antisense RNA control. *EMBO J.* **9**:1259–1266.
- Coutte, L., S. Alonso, N. Reveneau, E. Willery, B. Quatannens, C. Lochet, and F. Jacob-Dubuisson. 2003. Role of adhesin release for mucosal colonization by a bacterial pathogen. *J. Exp. Med.* **197**:735–742.
- Coutte, L., R. Antoine, H. Drobecq, C. Lochet, and F. Jacob-Dubuisson. 2001. Subtilisin-like autotransporter serves as maturation protease in a bacterial secretion pathway. *EMBO J.* **20**:5040–5048.
- Culham, D. E., J. Henderson, R. A. Crane, and J. M. Wood. 2003. Osmosensor ProP of *Escherichia coli* responds to the concentration, chemistry, and molecular size of osmolytes in the proteoliposome lumen. *Biochemistry* **42**:410–420.
- Cullen, P. A., S. J. Cordwell, D. M. Bulach, D. A. Haake, and B. Adler. 2002. Global analysis of outer membrane proteins from *Leptospira interrogans* serovar Lai. *Infect. Immun.* **70**:2311–2318.
- Elias, A. F., J. L. Bono, J. A. Carroll, P. Stewart, K. Tilly, and P. Rosa. 2000. Altered stationary-phase response in a *Borrelia burgdorferi* *rpoS* mutant. *J. Bacteriol.* **182**:2909–2918.
- Haake, D. A., C. Martinich, T. A. Summers, E. S. Shang, J. D. Pruetz, A. M. McCoy, M. K. Mazel, and C. A. Bolin. 1998. Characterization of leptospiral outer membrane lipoprotein LipL36: downregulation associated with late-log-phase growth and mammalian infection. *Infect. Immun.* **66**:1579–1587.
- Hamburger, Z. A., M. S. Brown, R. R. Isberg, and P. J. Bjorkman. 1999. Crystal structure of invasin: a bacterial integrin-binding protein. *Science* **286**:291–295.
- Heermann, R., and K. Jung. 2004. Structural features and mechanisms for sensing high osmolarity in microorganisms. *Curr. Opin. Microbiol.* **7**:168–174.
- Hendrixson, D. R., M. L. de la Morena, C. Stathopoulos, and J. W. St. Geme III. 1997. Structural determinants of processing and secretion of the *Haemophilus influenzae* hap protein. *Mol. Microbiol.* **26**:505–518.
- Hengge-Aronis, R. 1996. Back to log phase: sigma S as a global regulator in the osmotic control of gene expression in *Escherichia coli*. *Mol. Microbiol.* **21**:887–893.
- Johnson, R. C., and N. D. Gary. 1963. Nutrition of "*Leptospira pomona*." II. Fatty acid requirements. *J. Bacteriol.* **85**:976–982.
- Johnson, R. C., and V. G. Harris. 1967. Differentiation of pathogenic and saprophytic leptospires. I. Growth at low temperatures. *J. Bacteriol.* **94**:27–31.
- Kenny, B., and B. B. Finlay. 1995. Protein secretion by enteropathogenic *Escherichia coli* is essential for transducing signals to epithelial cells. *Proc. Natl. Acad. Sci. USA* **92**:7991–7995.
- Ko, A. I., M. Galvao Reis, C. M. Ribeiro Dourado, W. D. Johnson, Jr., L. W. Riley, and the Salvador Leptospirosis Study Group. 1999. Urban epidemic of severe leptospirosis in Brazil. *Lancet* **354**:820–825.
- Koizumi, N., and H. Watanabe. 2004. Leptospiral immunoglobulin-like proteins elicit protective immunity. *Vaccine* **22**:1545–1552.
- Luo, Y., E. A. Frey, R. A. Pfuetzner, A. L. Creagh, D. G. Knoechel, C. A. Haynes, B. B. Finlay, and N. C. Strynadka. 2000. Crystal structure of enteropathogenic *Escherichia coli* intimin-receptor complex. *Nature* **405**:1073–1077.
- Matsunaga, J., M. A. Barocchi, J. Croda, T. A. Young, Y. Sanchez, I. Siqueira, C. A. Bolin, M. G. Reis, L. W. Riley, D. A. Haake, and A. I. Ko. 2003. Pathogenic *Leptospira* species express surface-exposed proteins belonging to the bacterial immunoglobulin superfamily. *Mol. Microbiol.* **49**:929–945.
- Matsunaga, J., T. A. Young, J. K. Barnett, D. Barnett, C. A. Bolin, and D. A. Haake. 2002. Novel 45-kilodalton leptospiral protein that is processed to a 31-kilodalton growth-phase-regulated peripheral membrane protein. *Infect. Immun.* **70**:323–334.
- McGuirk, P., and K. H. Mills. 2000. Direct anti-inflammatory effect of a bacterial virulence factor: IL-10-dependent suppression of IL-12 production by filamentous hemagglutinin from *Bordetella pertussis*. *Eur. J. Immunol.* **30**:415–422.
- Mekalanos, J. J. 1992. Environmental signals controlling expression of virulence determinants in bacteria. *J. Bacteriol.* **174**:1–7.
- Morgan, J., S. L. Bornstein, A. M. Karpati, M. Bruce, C. A. Bolin, C. C.

- Austin, C. W. Woods, J. Lingappa, C. Langkop, B. Davis, D. R. Graham, M. Proctor, D. A. Ashford, M. Bajani, S. L. Bragg, K. Shutt, B. A. Perkins, and J. W. Tappero. 2002. Outbreak of leptospirosis among triathlon participants and community residents in Springfield, Illinois, 1998. *Clin. Infect. Dis.* **34**:1593–1599.
27. Nally, J. E., S. Artiushin, and J. F. Timoney. 2001. Molecular characterization of thermoinduced immunogenic proteins Qlp42 and Hsp15 of *Leptospira interrogans*. *Infect. Immun.* **69**:7616–7624.
28. Nascimento, A. L., A. I. Ko, E. A. Martins, C. B. Monteiro-Vitorello, P. L. Ho, D. A. Haake, S. Verjovski-Almeida, R. A. Hartskeerl, M. V. Marques, M. C. Oliveira, C. F. Menck, L. C. Leite, H. Carrer, L. L. Coutinho, W. M. Degraive, O. A. Dellagostin, H. El-Dorry, E. S. Ferro, M. I. Ferro, L. R. Furlan, M. Gamberini, E. A. Giglioti, A. Goes-Neto, G. H. Goldman, M. H. Goldman, R. Harakava, S. M. Jeronimo, I. L. Junqueira-de-Azevedo, E. T. Kimura, E. E. Kuramae, E. G. Lemos, M. V. Lemos, C. L. Marino, L. R. Nunes, R. C. de Oliveira, G. G. Pereira, M. S. Reis, A. Schriefer, W. J. Siqueira, P. Sommer, S. M. Tsai, A. J. Simpson, J. A. Ferro, L. E. Camargo, J. P. Kitajima, J. C. Setubal, and M. A. Van Sluys. 2004. Comparative genomics of two *Leptospira interrogans* serovars reveals novel insights into physiology and pathogenesis. *J. Bacteriol.* **186**:2164–2172.
29. Palaniappan, R. U., Y. F. Chang, S. S. Jusuf, S. Artiushin, J. F. Timoney, S. P. McDonough, S. C. Barr, T. J. Divers, K. W. Simpson, P. L. McDonough, and H. O. Mohammed. 2002. Cloning and molecular characterization of an immunogenic LigA protein of *Leptospira interrogans*. *Infect. Immun.* **70**:5924–5930.
30. Ren, S. X., G. Fu, X. G. Jiang, R. Zeng, Y. G. Miao, H. Xu, Y. X. Zhang, H. Xiong, G. Lu, L. F. Lu, H. Q. Jiang, J. Jia, Y. F. Tu, J. X. Jiang, W. Y. Gu, Y. Q. Zhang, Z. Cai, H. H. Sheng, H. F. Yin, Y. Zhang, G. F. Zhu, M. Wan, H. L. Huang, Z. Qian, S. Y. Wang, W. Ma, Z. J. Yao, Y. Shen, B. Q. Qiang, Q. C. Xia, X. K. Guo, A. Danchin, I. Saint Girons, R. L. Somerville, Y. M. Wen, M. H. Shi, Z. Chen, J. G. Xu, and G. P. Zhao. 2003. Unique physiological and pathogenic features of *Leptospira interrogans* revealed by whole-genome sequencing. *Nature* **422**:888–893.
31. Shang, E. S., T. A. Summers, and D. A. Haake. 1996. Molecular cloning and sequence analysis of the gene encoding LipL41, a surface-exposed lipoprotein of pathogenic *Leptospira* species. *Infect. Immun.* **64**:2322–2330.
32. St. Geme, J. W., III, and S. Grass. 1998. Secretion of the *Haemophilus influenzae* HMW1 and HMW2 adhesins involves a periplasmic intermediate and requires the HMWB and HMWC proteins. *Mol. Microbiol.* **27**:617–630.
33. Thiermann, A. B., R. D. McClellan, and H. T. Hill. 1984. Improved techniques for the isolation of leptospires from swine abortion cases. *Ann. Proc. Am. Assoc. Vet. Lab. Diagn.* **27**:233–244.
34. Walderhaug, M. O., J. W. Polarek, P. Voelkner, J. M. Daniel, J. E. Hesse, K. Altendorf, and W. Epstein. 1992. KdpD and KdpE, proteins that control expression of the *kdpABC* operon, are members of the two-component sensor-effector class of regulators. *J. Bacteriol.* **174**:2152–2159.
35. Zuerner, R. L., W. Knudtson, C. A. Bolin, and G. Trueba. 1991. Characterization of outer membrane and secreted proteins of *Leptospira interrogans* serovar pomona. *Microb. Pathog.* **10**:311–322.

Editor: D. L. Burns