

Genetics and Virulence Association of the *Shigella flexneri* Sit Iron Transport System[∇]

Carolyn R. Fisher,¹ Nicola M. L. L. Davies,¹ Elizabeth E. Wyckoff,¹ Zhengyu Feng,¹
Edwin V. Oaks,² and Shelley M. Payne^{1*}

Section of Molecular Genetics and Microbiology, Institute for Cellular and Molecular Biology, 1 University Station A5000,
The University of Texas at Austin, Austin, Texas 78712,¹ and Department of Enteric Infections/DCD&I,
Walter Reed Army Institute of Research, Silver Spring, Maryland 20910-7500²

Received 18 January 2009/Returned for modification 16 February 2009/Accepted 5 March 2009

The *sit*-encoded iron transport system is present within pathogenicity islands in all *Shigella* spp. and some pathogenic *Escherichia coli* strains. The islands contain numerous insertion elements and sequences with homology to bacteriophage genes. The *Shigella flexneri sit* genes can be lost as a result of deletion within the island. The formation of deletions was dependent upon RecA and occurred at relatively high frequency. This suggests that the *sit* region is inherently unstable, yet *sit* genes are maintained in all of the clinical isolates tested. Characterization of the *sitABCD* genes in *S. flexneri* indicates that they encode a ferrous iron transport system, although the genes are induced aerobically. The *sit* genes provide a competitive advantage to *S. flexneri* growing within epithelial cells, and a *sitA* mutant is outcompeted by the wild type in cultured epithelial cells. The Sit system is also required for virulence in a mouse lung model. The *sitA* mutant was able to infect the mice and induce a protective immune response but was avirulent compared to its wild-type parent strain.

The *Shigella* species are invasive enteric pathogens that cause dysentery in their human host. Following ingestion, the bacteria invade colonic epithelial cells and multiply within the cytoplasm. They spread from cell to cell, eventually causing death of the host cells and provoking an intense inflammatory response (39, 51).

Shigella spp. are closely related to and considered part of the same species as *Escherichia coli* (40, 41). The patterns that have emerged from analysis of the *E. coli* and *Shigella* genomes are that functional genes that are present in two or more members of this enteric group show a high degree of sequence conservation, but the overall organization of the genomes is different. Compared with the *E. coli* K-12 genome, each of the *Shigella* genomes contains a significant number of rearrangements, as well as insertions and deletions. Some of the insertions, termed pathogenicity islands (PAI), are quite large and include genes that increase virulence or fitness. Deletions in the chromosome may also affect pathogenicity; for example, the deletion of lysine decarboxylase genes in *S. flexneri* is associated with increased virulence (25, 26).

One class of genes that shows significant variability among *Shigella* and *E. coli* strains are those encoding high-affinity transport systems for iron. Iron is an essential element for *Shigella*, but the acquisition of iron is complicated by its insolubility in aerobic environments at neutral pH. Bacteria often have multiple pathways for importing iron, allowing the utilization of Fe⁺⁺, Fe⁺⁺⁺, and iron bound to a variety of carriers.

Ferric iron is efficiently transported into *E. coli* K-12 by the siderophore enterobactin (Ent) (11). This low-molecular-

weight iron chelator is synthesized and secreted into the environment, where it binds ferric iron with high affinity (32), and the ferri-siderophore complex is transported back into the cell via a specific transporter system (Fep). Most *E. coli*, *Shigella dysenteriae* type 1, and *Shigella sonnei* strains synthesize and transport enterobactin (11, 34, 36, 38). However, many *Shigella boydii* and *Shigella flexneri* strains are Ent⁻ due to deletions and point mutations within the *ent/fep* operons (36, 52). *Shigella* strains that fail to produce enterobactin produce a different siderophore, aerobactin, and some strains produce both siderophores (22, 36). Aerobactin is a secondary hydroxamate, and genes for its synthesis (*iucABCD*) and receptor (*iutA*) are located within a single operon (4, 22). These genes may be on either the chromosome (22, 24) or a plasmid (4, 8). The chromosomal genes are located within PAI in *S. flexneri* and *S. boydii* (31, 42, 57).

The *Shigella* species have additional iron transport systems that are not found in *E. coli* K-12. A transport system with homology to the *Salmonella enteritidis* Sit system is found in all of the *Shigella* species. Some strains have heme transport systems, the best characterized of which are the *S. dysenteriae* Shu heme transporter (30, 61) and a nearly identical system in *E. coli* O157:H7 (54). *S. dysenteriae* type 1 strains also have the *iro* genes (43) for the biosynthesis of salmochelin, a modified form of enterobactin first described in *Salmonella* (13). Additional transport systems for iron are present in enteric pathogens, and some of these are completely uncharacterized.

The only iron transport system that appears to be common to all members of the *E. coli/Shigella* group is Feo. This is a ferrous iron cytoplasmic membrane transporter encoded by the *feoABC* genes (5, 19). FeoB is a cytoplasmic membrane protein with GTPase activity (23), but the mechanism of transport and the functions of FeoA and FeoC have not been fully determined.

Despite the apparent variation in the iron transport systems

* Corresponding author. Mailing address: The University of Texas at Austin, Molecular Genetics & Microbiology, 1 University Station A5000, Austin, TX 78712-0162. Phone: (512) 471-9258. Fax: (512) 471-7088. E-mail: payne@mail.utexas.edu.

[∇] Published ahead of print on 16 March 2009.

found in *Shigella*, there are consistent patterns. All express at least one siderophore and have the Feo and Sit transporters. *S. flexneri* serotype 2a strains SA100 and 2457T have only these three systems and thus were chosen for analysis of iron transport in vitro and within the host cell cytoplasm.

Analysis of isogenic strains lacking one or more of the iron transporters showed that no single mutation eliminated intracellular growth, as the single *iucD*, *feoB*, and *sitA* mutants all produced plaques in cultured cells (48). The double mutants produced smaller plaques, but only the triple mutant was completely defective in growth and plaque formation (48).

Additional information about the roles of these three systems in iron acquisition comes from our previous studies of their regulation. All of the iron transport systems are negatively regulated by Fur (1, 19), an iron-binding repressor protein. However, analysis of the expression of iron transport genes by wild-type *S. flexneri* growing in the intracellular environment indicated that only the *sit* genes were highly expressed in this environment (47). The aerobactin genes were downregulated in the intracellular environment (14), and *feo* expression appeared unchanged. The fact that, of the three iron transport systems, only *sit* is normally induced intracellularly suggests that it plays an important role when the bacteria inhabit the host cell cytoplasm. The failure of a *sit* mutant to show a defect in plaque formation may indicate that iron starvation induced by loss of the Sit system resulted in upregulation of the *feo* and *iuc* genes when they would not normally have been expressed. Further, since all of these genes appear to be regulated similarly by Fur in vitro, this differential expression of the iron transport genes in the intracellular environment suggested that factors other than iron were contributing to their regulation.

An additional environmental factor that controls the expression of these genes is oxygen (3). As previously noted in *E. coli* K-12 (19), expression of the *feo* ferrous iron transport operon was induced under anaerobic conditions, where ferrous iron should predominate over ferric iron. In contrast, the aerobactin genes were induced aerobically, consistent with their role in ferric iron uptake (3). Surprisingly, the *S. flexneri sit* genes were repressed when the cells were grown anaerobically (3), although the homologous *sitABCD* genes in *Salmonella* encode a ferrous iron uptake system (20, 63). Consistent with aerobic induction, the Sit system was sufficient to support plaque formation by *S. flexneri* under aerobic, but not anaerobic, conditions (3).

Because the Sit iron transport system was found in all of the *Shigella* species and was induced when the bacteria were growing intracellularly, we undertook a characterization of its genetics and role in the virulence of *S. flexneri*.

MATERIALS AND METHODS

Strains and media. Bacterial strains and mutants are described in Table 1. Strains were maintained in Trypticase soy broth (TSB) with 20% glycerol at -80°C . Cultures were routinely grown at 37°C in Luria-Bertani (LB) broth (1% tryptone, 0.5% yeast extract, 1% NaCl) or on LB agar. Congo red agar (35), which contains TSB, 0.01% Congo red, and 1.5% agar, was used to distinguish virulent isolates by their red color. For growth of *S. flexneri* strains in minimal medium, MM9 (53) without added iron and containing 2 μg of nicotinic acid per ml and 0.4% (wt/vol) glucose was used. The aerobactin for supplementing *S. flexneri* iron transport mutants was sterile culture supernatant of *S. flexneri* SA101 prepared as previously described (48). Antibiotics were used at the following

concentrations per milliliter: 250 μg of carbenicillin, 50 μg of kanamycin, and 30 μg of chloramphenicol. Expression of T7 RNA polymerase was induced with 500 μM isopropyl- β -D-thiogalactopyranoside (IPTG; Promega, Madison, WI).

Construction of Lac⁺ *S. flexneri* strains. SM100 Lac and SM166 Lac were constructed by inserting the *E. coli lacZ* gene into the intergenic region between the *glmS* and *pstS* genes with Tn7 transposase encoded on plasmid pGRGlac1, which was constructed as follows. *E. coli lacZ* was amplified from strain ARM110 (29) with Pfx polymerase (Invitrogen, Carlsbad, CA) and primers ec.lacZ.1293 (5'-ATACGCAAACCGCTCTCCC-3') and ec.lacZ.5024.rev (5'-CTGACTTTCTCAATAAATGCCTCTACTG-3'). The resulting PCR product was cloned into the SmaI site of pGRG36 (27) to give pGRGlac1. This plasmid was introduced into the indicated *S. flexneri* strains by electroporation, and colonies that were blue on 5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside (X-Gal) plates were selected following growth at 37°C . The proper integration of *lacZ* was confirmed by PCR with a primer hybridizing within the *glmS* gene and one hybridizing within *lacZ*.

PCR detection of *sit* and T7 promoters. The presence of the *sit* operon was detected by amplification with primers sitAB forward (5'-CTCTTGAGCACTGAAGGAG-3') and sitAB reverse (5'-CGCACAAATCCCATAATC-3'). The T7 promoter region was detected by PCR with primers SMP028 (5'-GTGTCCCTTCTCCCTATAGTG-3') and SMP033 (5'-CTTACTACAGACCTGTGTG-3').

DNA sequence analysis. DNA sequencing was performed by the University of Texas Institute for Cellular and Molecular Biology DNA Core Facility with an ABI Prism 3700 DNA sequencer. Analysis of DNA sequences was carried out with MacVector 7.2 and Clone Manager 7.04. BLAST searches and other bioinformatic analyses were done with the National Center for Biotechnology and Enteropathogen Resource Integration Center (www.ERICBRC.org) databases. Pairwise alignments were carried out with ClustalW from within MacVector 7.2, and genomic alignments were done with MAUVE (9).

Iron transport assay. Cultures of *S. flexneri* SM193w containing the indicated plasmid were grown overnight in LB broth supplemented with 1/10 volume of sterile MM9 supernatant from *S. flexneri* strain SA101 as a source of aerobactin. Overnight cultures were diluted 1/25 into MM9 with aerobactin and grown to late exponential phase with aeration at 37°C . Transport assays were performed in triplicate at room temperature in the presence or absence of 5 mM sodium ascorbate, as previously described (62).

T7 polymerase sensitivity assay. Serial dilutions were made from cultures of SM100/pAR1219, SM160/pAR1219, or *E. coli* HB101/pEG1/pAR1219, and each dilution was plated on both TSB agar containing carbenicillin (viable count) and the same medium with 1 mM IPTG. The frequency of resistance to T7 RNA polymerase was calculated by dividing the number of colonies on the plates containing IPTG by the viable count. The frequency of loss of the *sit* operon in T7 RNA polymerase-resistant isolates was determined by screening T7 RNA polymerase-resistant colonies by PCR.

Henle cell plaque assay. Monolayers of Henle cells (intestinal 407 cells; American Type Culture Collection, Manassas, VA) were maintained in Eagle's minimum essential medium with 2 mM glutamine, 10% fetal bovine serum, tryptone phosphate broth, and minimal essential amino acids in a 5% CO₂ atmosphere at 37°C . Plaque assays were performed as described previously (16, 33), with the following modifications. Confluent Henle cell monolayers grown in 35-mm-diameter plates were infected with 2×10^4 bacteria. After 60 min of incubation, the medium overlying the Henle cells was removed and replaced with fresh medium plus 0.45% (wt/vol) glucose and 20 μg of gentamicin per ml. The cells were then incubated for 72 h.

Competition assay. Henle cell monolayers were infected with a mixture of equal numbers of SM166 Lac and SM100 or SM166 and SM100 Lac bacteria. The exact ratio of the two strains in each experiment was determined by plating dilutions of the inoculum on agar medium with X-Gal and counting the Lac⁺ and Lac⁻ colonies. After 72 h of incubation, the infected monolayers were detached with 0.025% (wt/vol) trypsin and lysed with 0.5% (wt/vol) sodium deoxycholate as described previously (16) to recover the intracellular bacteria. Serial dilutions of the recovered bacteria were plated on TSB agar with X-Gal, and the ratio of SM100 to SM166 bacteria recovered was determined by counting the Lac⁺ and Lac⁻ colonies. The competitive index was calculated as the ratio of SM166 to SM100 bacteria recovered from the cells divided by the ratio of SM166 to SM100 bacteria in the inoculum.

Mouse virulence. Eight-week-old BALB/cJ female mice weighing approximately 25 g (Jackson Laboratory, Bar Harbor, ME) were sedated by intramuscular injection of a mixture of xylazine hydrochloride (40 mg/kg) (Rompun; Mobay Corp., Shawnee, KS) and ketamine hydrochloride (12 mg/kg) (Ketaset; Aveco Co., Fort Dodge, IA) in 50 μl of saline. Groups of five mice were inoculated with 30 μl of a suspension containing 10^7 wild-type or mutant *S.*

TABLE 1. Bacterial strains and plasmids used in this study

Strain or plasmid	Characteristic(s)	Reference or source
<i>E. coli</i> strains		
DH5 α	<i>endA1 hsdR17 supE44 thi-1 recA1 gyrA96 relA1</i> Δ (<i>lacZYA-argF</i>) <i>U169 deoR</i> [ϕ 80 <i>dlac</i> Δ (<i>lacZ</i>)M15]	50
HB101	F ⁻ Δ (<i>gpt-proA</i>)62 <i>leu supE44 ara-14 galK2 lacY1</i> Δ (<i>mcrC-mrr</i>) <i>rpsL20 xyl-5 mtl-1 recA13</i>	50
<i>S. flexneri</i> strains		
2457T	<i>S. flexneri</i> wild-type serotype 2a	60
SA100	<i>S. flexneri</i> wild-type serotype 2a	36
SA101	SA100, Vir ⁻ , source of aerobactin	21
SA240	SA100 <i>iucD::kan</i>	21
SM100	SA100 Str ^r	Stefan Seliger
SM160	SM100 <i>recA::cam</i>	Laura Runyen-Janecky
SM166	SM100 <i>sitA::cam</i>	48
SM193w	SM100 <i>sitA::cam feoB::tmp iucD::kan</i>	48, 62
SM100 Lac	SM100 with <i>lacZ</i> integrated into chromosome	This study
SM166 Lac	SM166 with <i>lacZ</i> integrated into chromosome	This study
Plasmids		
pACYC184	Plasmid cloning vector	6
pWKS30	Plasmid cloning vector	58
pLAFR1	Cosmid cloning vector	12
pEG1	Cosmid vector pLAFR1 carrying SA100 <i>sit</i> operon and surrounding sequence	48
pEG3	pWKS30 carrying 8-kb HindIII fragment from pEG1 including <i>sitABCD</i>	This study
pAR1219	T7 RNA polymerase gene under control of <i>lacUV5</i> promoter in pBR322	10
pFZ68	pACYC184 carrying 4.6-kb EcoRI fragment of pEG1; includes entire <i>sit</i> operon	This study
pFZ69	pACYC184 carrying 6.4-kb EcoRI fragment of pEG1	This study
pFZ70	pACYC184 carrying 13-kb EcoRI fragment of pEG1; includes T7 promoter region	This study
pFZ74	pACYC184 carrying 5.6-kb EcoRV fragment of pFZ70; includes T7 promoter region	This study
pFZ76	pACYC184 carrying 2-kb EcoRV/DraI fragment of pFZ74; includes T7 promoter region	This study
pGRG36	Tn7 insertion vector	27
pGRG1ac1	<i>E. coli lacZ</i> cloned into pGRG36	This study

flexneri bacteria, which was applied dropwise to the external nares of each mouse with a 100- μ l Hamilton syringe. Mice were observed for 10 days for deaths. Weight loss and rebound were recorded. Three weeks after the initial infection, surviving mice were challenged intranasally with a lethal dose of *S. flexneri* serotype 2a strain 2457T (10^7 CFU/30 μ l). The mouse challenge dose was prepared from a frozen lot of *S. flexneri* serotype 2a bacteria that had been harvested during the log phase of growth and then stored in liquid nitrogen.

RESULTS

Sit iron transport. While the *Salmonella* Sit system has been shown to be a ferrous iron transporter (20), the iron ligand preference of the related *S. flexneri* Sit system has not been determined experimentally. Our previous data indicated that expression of the *S. flexneri sit* genes is induced, and it contributes to intracellular growth most effectively, under aerobic conditions (3) where iron should normally be in the ferric form. Thus, while the *S. flexneri* genes were classified as *sit* genes based on amino acid identity (64 to 77%) of the predicted gene products to the *Salmonella* Sit proteins (48), there could be differences in the structure and function of the proteins and in the metal ligands they transport. Therefore, we determined the uptake of ferric and ferrous iron by SM193w carrying *sitABCD* cloned into a low-copy-number vector, pWKS30. SM193w has mutations in *iucD*, *sitA*, and *feoB* and does not grow or transport significant amounts of iron in the absence of added siderophore or expression of a cloned iron transport system (48, 62). Thus, there is minimal background iron transport in this strain. Transport by the Sit system was measured in the presence or absence of ascorbate, which keeps the iron in the reduced ferrous form (Fig. 1). There was min-

imal difference in the uptake of radiolabeled iron in the presence or absence of the *sit* genes when no ascorbate was present in the transport buffer. This indicates that Sit did not enhance the uptake of ferric iron. In contrast, the *sit* genes greatly increased the transport of iron when ascorbate was present, showing that the Sit system is a ferrous iron transporter.

Genomic location and organization of the *sit* genes. The *sit* genes were present in the chromosomes of all of the *Shigella* strains that we tested (48), but as noted by Sabri et al. (49), the genes had different locations in different strains. Therefore, we examined the sequenced genomes of additional *Shigella* strains to determine the location and organization of the *sit* genes (Fig. 2). The genes are found within islands and are flanked by a variety of insertion sequences and phage-like genes. The locations of the islands relative to the *E. coli* K-12 genome are different, even in closely related strains such as the two sequenced *S. flexneri* serotype 2a strains 2457T and 301 (Fig. 2). It is not clear whether the genes were acquired by horizontal transmission multiple times in *Shigella* lineages or whether the genes have moved in some strains subsequent to their acquisition. In the *E. coli/Shigella* group, the Sit island insertions fall into five clusters. The uropathogenic and avian pathogenic *E. coli* strains have the island at a common position, compared to the *E. coli* K-12 genome, but the sizes of the islands and the orientation of the *sit* operon within the islands differ (Fig. 2). *E. coli* SMS-3-5, an environmental isolate that has numerous virulence factors and multiple drug resistance genes (8), has two copies of the *sit* genes (Fig. 2). One copy is on the chromosome at a unique site, while the other is on a plasmid. A plasmid-




FIG. 1. The *S. flexneri* Sit system transports ferrous iron. *S. flexneri* strain SM193w carrying either plasmid vector pWKS30 or the cloned *sitABCD* genes (pEG3) was grown to late exponential phase. The transport of ⁵⁵Fe was measured as described in Materials and Methods after 5 min of incubation in transport buffer with or without supplementation with 5 mM sodium ascorbate. The data shown are the average of three experiments, and the error bars represent 1 standard deviation.

encoded Sit system has also been noted in an avian pathogenic *E. coli* strain (18, 49). There also is considerable variation in the sizes of the islands and the orientation of the genes among sequenced *Shigella* isolates (Fig. 2). Two of the *S. flexneri* serotype 2a strains, 2457T and SA100, have the island adjacent to the *S. flexneri* homolog of *E. coli* K-12 *aspS*, a site 500 kb away and in the opposite orientation from the other *Shigella* isolates. As described by Chen and Schneider (7), comparison of the *sit* regions in *S. flexneri* 2457T and 301 suggests that there may have been rearrangements subsequent to the acquisition of the *sit* island in the ancestral strain, since both strains have the same insertion, which includes *ipaH*, homologs of *recE* and *intR*, and the *sit* genes (Fig. 3). The uniform presence of the *sit* genes in *Shigella* strains, despite the apparent instability of the islands, suggests a selective advantage for their retention.

Instability of the *sit* island. To measure the stability of the *sit* island, we took advantage of the fact that the *sit* genes in *S. flexneri* serotype 2a strains 2457T and SA100 were linked to tandem repeats of a sequence with a perfect match to a strong T7 polymerase promoter (7). We observed that expression of T7 polymerase in these strains is lethal and that a majority of the T7 polymerase-resistant colonies had deletions in the *sit* PAI, as determined by DNA hybridization to *Shigella* genomic microarrays (data not shown). To understand the basis of this lethality, cosmid pEG1, containing much of the *sit* PAI, was mated into *E. coli* strain HB101 and found to cause lethality when T7 polymerase was induced from the *lacUV5* promoter on plasmid pAR1219 (Fig. 4). Various subclones obtained from pEG1 were tested in this assay, and only fragments containing the T7 promoter repeats caused lethality (Fig. 3). The shortest fragment had no open reading frame (ORF) downstream of the T7 promoters and only a portion of the integrase ORF upstream of the promoters. This, combined with the observation that lethality was only observed upon induction of the T7 polymerase genes, indicates that it is the expression of the T7 promoters themselves, and not the expression of a specific gene, that is responsible for cell death. The mechanism of killing is not known, although it is possible that the high level




FIG. 2. (A) PAI containing *sit* islands in *E. coli* and *Shigella*. The line indicates the *E. coli* K-12 chromosome with the relative positions of the Sit islands for each sequenced strain shown above the line. Strains a to n contain genomic islands, while strains d and o also have *sit* genes located on a plasmid, indicated by the circle above the chromosome. The values below the line are distances in megabase pairs. (B) The Sit islands from specific strains are shown, with the length of the line indicating the size of the island and the bold arrow indicating the relative location and direction of transcription of the *sitABCD* genes. Lines crossed by the short double lines are not to scale. Abbreviations: UPEC, uropathogenic *E. coli*; Apec, avian pathogenic *E. coli*; S. f., *S. flexneri*; S. b., *S. boydii*; S. s., *S. sonnei*; S. d., *S. dysenteriae*; E.c., *E. coli*.

of activity of these strong promoters leads to depletion of nucleotide pools within the cell.

In *S. flexneri*, sensitivity to T7 polymerase also correlated with deletion of the promoter repeats, and PCR analysis showed that resistant isolates had deletions of the T7 promoters (Table 2 and Fig. 3). To determine the frequency of deletions, the numbers of colonies in the presence and absence of T7 induction were compared (Table 2). The frequency of deletion in either the wild-type strain or a derivative carrying a chloramphenicol resistance cassette in the *sit* operon was approximately 10⁻⁴. Additional PCR analysis showed that many of the T7 promoter deletions included a large portion of the PAI containing the *sit* genes (Table 2 and Fig. 3).

Mapping of several of the deletions showed that they varied in size and were not simply excisions of the entire island (Fig. 3). The large number of insertion sequences and rearrangements within the islands made it difficult to determine the precise endpoints of the deletions, but it appeared that they might result from recombination between two copies of the

TABLE 2. Frequency of T7 RNA polymerase resistance and loss of *sitA* in RecA⁺ and RecA⁻ *S. flexneri*

Strain (genotype)	Avg frequency of T7 polymerase resistance ^a	Range	% of resistant isolates with deletion of:	
			T7 promoters ^b	<i>sit</i> genes ^c
SM100/pAR1219 (wild type)	2.7×10^{-4}	1.0×10^{-4} – 4.1×10^{-4}	100	75
SM166/pAR1219 (<i>sit::cam</i>)	8.9×10^{-3}	1.6×10^{-3} – 5.6×10^{-4}	100	60
SM160/pAR1219 (<i>recA::cam</i>)	2.9×10^{-5}	2.0×10^{-5} – 5.7×10^{-6}	0	0

^a Strains were grown in LB broth and plated on medium containing IPTG to induce expression of T7 polymerase from pAR1219. The frequency is the ratio of colonies on IPTG-containing medium to the number of colonies from the same culture plated on medium without IPTG. The average and range of values for 4 to 10 separate experiments are shown.

^b A minimum of 10 individual colonies was selected and tested by PCR with primers in the T7 promoters.

^c Individual colonies were selected and tested by PCR with primers in the *sitAB* genes (SM100 and SM160) or by plating on chloramphenicol (SM166) to determine the presence or absence of the *sit* operon.

($P = 0.24$, two-tailed t test). Thus, the presence of the *lacZ* gene as a marker did not significantly influence the competitive index of the strains.

Sit is required for virulence in the mouse lung model. To further determine whether Sit plays a role in the virulence of *S. flexneri*, mice were infected with the wild type or the *sitA* mutant in a lung model of *Shigella* virulence (56) (Table 3). In this model, both the wild type and the streptomycin-resistant parent of the mutants were virulent; all five mice in each group died. In contrast, the *sitA* mutant was attenuated (Table 3). Only one of the five mice died, and the other four recovered within 10 days. The aerobactin (*iucD*) mutant, which had previously been shown to be indistinguishable from the wild type in plaque and Sereny (37) assays, was virulent in this model. Four of five mice inoculated with the *iucD* mutant died. This is consistent with our observation that the aerobactin genes are repressed during intracellular growth and do not play a significant role in iron acquisition after the invasion of epithelial cells (15, 21).

To ensure that the mice that survived infection with the *sit* mutant had been successfully inoculated with the mutant strain, a back challenge was performed (Table 3). Surviving

mice were challenged with a lethal dose of the wild-type serotype 2a strain 21 days after the initial infection. All of the mice that had initially received the *sitA* strain survived a challenge with the wild type, while most of the mice (four of five) that had not previously been infected succumbed to this dose. Therefore, the mice that survived infection with the *sit* mutant had been successfully inoculated and had mounted an effective immune response that prevented reinfection. Similarly, the one animal that survived infection with the *iucD* mutant (SA240) was also protected against a subsequent challenge.

DISCUSSION

Shigella spp. have a variety of iron transport systems, but three of these, Feo, Sit, and a siderophore, are common to all of the clinical isolates examined. Differences were noted in the type of siderophore produced. Enterobactin, salmochelin, aerobactin, or combinations of these were found in different strains. This suggests that siderophore synthesis is important in some stage of the *Shigella* lifestyle but virulence is not dependent on the production of any particular siderophore. In contrast, studies of other pathogenic *E. coli* strains have shown that aerobactin production is advantageous in certain pathogens (28, 55, 59).

The Sit system, which was first described in *Salmonella* (63) and was shown to be required for full virulence in this pathogen (17), is primarily present in bacterial species that invade and multiply within host cells. In *Salmonella*, Sit has a higher affinity for manganese than for iron and is thought to be primarily a manganese transporter under physiological conditions




FIG. 5. Competition between the wild type and the *sitA* mutant in a plaque assay. Henle cells were infected with a mixture of equal numbers of wild-type (SM100) and *sitA* mutant (SM166) *S. flexneri* bacteria and incubated in the presence of gentamicin to kill extracellular bacteria. In each set of experiments, one of the two strains was marked with *lacZ*. Bacteria were harvested from the plaques after 72 h of incubation, and the competitive index (ratio of *sitA* mutant to wild-type bacteria recovered) was determined. Each symbol represents the competitive index in a single experiment. The average competitive index is shown by a horizontal line. The asterisks indicate that the P value (two-tailed t test) for the difference between the observed competitive index and an expected index of 1.0 is <0.02 for SM166 Lac/SM100 and <0.04 for SM166/SM100 Lac.

TABLE 3. Virulence of wild type and iron transport mutants of *S. flexneri* in the mouse lung model

<i>S. flexneri</i> strain	Primary challenge		No. of survivors/total after back challenge with <i>S. flexneri</i> serotype 2a strain 2457T
	Time of death	No. of survivors/total	
SA100 (wild type)	All dead by day 9	0/5	ND ^a
SM100 (Str ^r)	All dead by day 4	0/5	ND
SA240 (<i>iucD</i>)	Last death on day 6	1/5	1/1
SM166 (<i>sitA</i>)	Last death on day 5	4/5	4/4
None			1/5

^a ND, not done.

(20). Similarly, the *S. flexneri* Sit system transports both iron and manganese (46) but functions well to provide iron when the bacteria are growing in the host cell cytoplasm (48). An *S. flexneri* mutant that lacks all iron transporters other than Sit grows normally in the intracellular environment and produces wild-type-size plaques in Henle cells in an aerobic environment. The *sit* mutant also grew intracellularly, presumably using the Feo and Iuc systems for iron acquisition. However, the plaques were slightly smaller, and in this work, we confirm that the *sit* mutant was at a competitive disadvantage by coinfecting the monolayers with the wild type and the *sit* mutant and directly comparing them for growth and intercellular spreading in a plaque assay. The wild type outcompeted the mutant and was recovered in a higher proportion from the infected monolayers. Further, the mutant was much less virulent in a mouse lung model of infection. In this model, the bacteria must be able to infect mice, invade lung epithelial cells, and provoke an inflammatory response. The mutant was able to infect the mice, since the survivors mounted an immune response and were resistant to challenge with the wild type, but the mutant was defective at some stage of invasion and intracellular replication. These data also indicate that a *sit* mutation may be appropriate to include in vaccine strains, since loss of *sit* causes attenuation and the *sit* mutant induces protective immunity.

It is unclear why the Sit system, a ferrous iron transporter, would be induced aerobically and repressed under anaerobic conditions, where ferrous iron should be more available. This regulation may reflect its dual role in iron and manganese transport, allowing increased transport of manganese, and consequently iron, when the bacteria are growing aerobically. Since the Sit system was sufficient for iron acquisition under aerobic conditions, it suggests that there is an accessible pool of ferrous iron in the presence of oxygen within the host cell cytoplasm. The cytoplasm is a reducing environment (44) which likely keeps a portion of the small pool of accessible iron in the ferrous form, even in the presence of oxygen. This is consistent with our earlier observation that the Sit system provided iron to intracellular bacteria when the cultured cells were growing aerobically but not in an anaerobic environment (3).

Both the Henle cell competition assay and the mouse virulence results indicate that the presence of Sit provides a selective advantage in the host. This is supported by the maintenance of this locus and the conservation of *sit* genes in virulent strains, despite the relatively high frequency of deletion. It also appears that the locus was acquired more than once in the evolution of *Shigella* since it is located at different sites and with different flanking sequences in various *Shigella* isolates. The Sit transport system is also present in enteroinvasive *E. coli* (48), which are derived from several ancestral lineages distinct from the *Shigella* strains (45). This suggests a selective advantage for the *sit* genes in enteroinvasive *E. coli* and *Shigella* rather than the continued presence of the genes in strains derived from a common ancestor. A number of the Sit islands contain phage-like genes, including integrases, and there are T7 promoters in the *S. flexneri* serotype 2a strains, suggesting phage-mediated acquisition of the islands. These T7 promoters are associated with lethality when T7 RNA polymerase is expressed within the cell.

Chen and Schneider (7) had shown that *S. flexneri* 2457T has an additional copy of the T7 promoter sequences in its genome, but

its similarity to the T7 promoter consensus is weak, whereas the repeated sequences in the Sit island are a near-perfect match to the promoter consensus. The T7 polymerase-resistant mutants presumably still have this copy since the Sit island deletion was the only deletion detected by hybridization to microarrays representing all of the genes in the *S. flexneri* 2457T and *E. coli* K-12 genomes. The weaker match to the T7 promoter apparently makes it less deleterious to *S. flexneri* when T7 polymerase is expressed in the cell, and loss of the T7 promoter tandem repeats in the Sit island is sufficient for resistance to the polymerase.

The observation that RecA is required for deletion of the island suggests that the deletions are caused by homologous recombination rather than being the result of T7 polymerase expression. Thus, the presence of the tandem T7 promoters within the island in these *S. flexneri* strains provides a fortuitous marker for the presence of the island, allowing us to use T7 polymerase resistance to select for bacteria from which the Sit PAI has been deleted and to show that the region can be deleted at relatively high frequency.

These data indicate that *S. flexneri* Sit is a ferrous iron transporter that enhances the growth of the bacteria within cultured epithelial cells and the production of disease in a mouse lung model. The ability of the *sit* genes to promote in vivo survival is likely responsible for the retention of these genes in clinical isolates.

ACKNOWLEDGMENTS

This work was supported by grant AI16935 from the National Institutes of Health.

We thank Jia-Wen Jessica Chang for technical assistance, Laura Runyen-Janecky and Enrique Gonzalez for construction of SM160 and pEG3, and Erin Murphy for helpful discussions.

REFERENCES

1. Bagg, A., and J. B. Neilands. 1987. Ferric uptake regulation protein acts as a repressor, employing iron (II) as a cofactor to bind the operator of an iron transport operon in *Escherichia coli*. *Biochemistry* **26**:5471–5477.
2. Boulette, M. L. 2007. *Shigella flexneri* ArcA and FNR regulate iron acquisition and contribute to plaque formation under anaerobic conditions. Ph.D. dissertation. University of Texas, Austin.
3. Boulette, M. L., and S. M. Payne. 2007. Anaerobic regulation of *Shigella flexneri* virulence: ArcA regulates Fur and iron acquisition genes. *J. Bacteriol.* **189**:6957–6967.
4. Carbonetti, N. H., and P. H. Williams. 1984. A cluster of five genes specifying the aerobactin iron uptake system of plasmid ColV-K30. *Infect. Immun.* **46**:7–12.
5. Cartron, M. L., S. Maddocks, P. Gillingham, C. J. Craven, and S. C. Andrews. 2006. Feo—transport of ferrous iron into bacteria. *Biomaterials* **19**:143–157.
6. Chang, A. C., and S. N. Cohen. 1978. Construction and characterization of amplifiable multicopy DNA cloning vehicles derived from the P15A cryptic miniplasmid. *J. Bacteriol.* **134**:1141–1156.
7. Chen, Z., and T. D. Schneider. 2006. Comparative analysis of tandem T7-like promoter containing regions in enterobacterial genomes reveals a novel group of genetic islands. *Nucleic Acids Res.* **34**:1133–1147.
8. Colonna, B., M. Nicoletti, P. Visca, M. Casalino, P. Valenti, and F. Maimone. 1985. Composite IS1 elements encoding hydroxamate-mediated iron uptake in *Flm* plasmids from epidemic *Salmonella* spp. *J. Bacteriol.* **162**:307–316.
9. Darling, A. E., T. J. Treangen, X. Messeguer, and N. T. Perna. 2007. Analyzing patterns of microbial evolution using the mauve genome alignment system. *Methods Mol. Biol.* **396**:135–152.
10. Davanloo, P., A. H. Rosenberg, J. J. Dunn, and F. W. Studier. 1984. Cloning and expression of the gene for bacteriophage T7 RNA polymerase. *Proc. Natl. Acad. Sci. USA* **81**:2035–2039.
11. Earhart, C. F. 2004. Iron uptake via the enterobactin system, p. 133–146. *In* J. H. Crosa, A. R. Mey, and S. M. Payne (ed.), *Iron transport in bacteria*. ASM Press, Washington, DC.
12. Friedman, A. M., S. R. Long, S. E. Briwn, W. J. Buikema, and F. Ausubel. 1982. Construction of a broad host range cosmid cloning vector and its uses in the genetic analysis of *Rhizobium meliloti*. *Gene* **18**:289–296.
13. Hantke, K., G. Nicholson, W. Rabsch, and G. Winkelmann. 2003. Salmochelins, siderophores of *Salmonella enterica* and uropathogenic *Escherichia coli* strains, are recognized by the outer membrane receptor Iron. *Proc. Natl. Acad. Sci. USA* **100**:3677–3682.

14. Headley, V., M. Hong, M. Galko, and S. M. Payne. 1997. Expression of aerobactin genes by *Shigella flexneri* during extracellular and intracellular growth. *Infect. Immun.* **65**:818–821.
15. Headley, V. L., and S. M. Payne. 1990. Differential protein expression by *Shigella flexneri* in intracellular and extracellular environments. *Proc. Natl. Acad. Sci. USA* **87**:4179–4183.
16. Hong, M., Y. Gleason, E. E. Wyckoff, and S. M. Payne. 1998. Identification of two *Shigella flexneri* chromosomal loci involved in intercellular spreading. *Infect. Immun.* **66**:4700–4710.
17. Janakiraman, A., and J. M. Schlauch. 2000. The putative iron transport system SitABCD encoded on SPI1 is required for full virulence of *Salmonella typhimurium*. *Mol. Microbiol.* **35**:1146–1155.
18. Johnson, T. J., K. E. Siek, S. J. Johnson, and L. K. Nolan. 2006. DNA sequence of a ColV plasmid and prevalence of selected plasmid-encoded virulence genes among avian *Escherichia coli* strains. *J. Bacteriol.* **188**:745–758.
19. Kammler, M., C. Schon, and K. Hantke. 1993. Characterization of the ferrous iron uptake system of *Escherichia coli*. *J. Bacteriol.* **175**:6212–6219.
20. Kehres, D. G., A. Janakiraman, J. M. Schlauch, and M. E. Maguire. 2002. SitABCD is the alkaline Mn²⁺ transporter of *Salmonella enterica* serovar Typhimurium. *J. Bacteriol.* **184**:3159–3166.
21. Lawlor, K. M., P. A. Daskaleros, R. E. Robinson, and S. M. Payne. 1987. Virulence of iron transport mutants of *Shigella flexneri* and utilization of host iron compounds. *Infect. Immun.* **55**:594–599.
22. Lawlor, K. M., and S. M. Payne. 1984. Aerobactin genes in *Shigella* spp. *J. Bacteriol.* **160**:266–272.
23. Marlovits, T. C., W. Haase, C. Herrmann, S. G. Aller, and V. M. Unger. 2002. The membrane protein FeoB contains an intramolecular G protein essential for Fe(II) uptake in bacteria. *Proc. Natl. Acad. Sci. USA* **99**:16243–16248.
24. Marolda, C. L., M. A. Valvano, K. M. Lawlor, S. M. Payne, and J. H. Crosa. 1987. Flanking and internal regions of chromosomal genes mediating aerobactin iron uptake systems in enteroinvasive *Escherichia coli* and *Shigella flexneri*. *J. Gen. Microbiol.* **133**:2269–2278.
25. Maurelli, A. T. 2007. Black holes, antivirulence genes, and gene inactivation in the evolution of bacterial pathogens. *FEMS Microbiol. Lett.* **267**:1–8.
26. Maurelli, A. T., R. E. Fernandez, C. A. Bloch, C. K. Rode, and A. Fasano. 1998. “Black holes” and bacterial pathogenicity: a large genomic deletion that enhances the virulence of *Shigella* spp. and enteroinvasive *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **95**:3943–3948.
27. McKenzie, G. J., and N. L. Craig. 2006. Fast, easy and efficient: site-specific insertion of transgenes into enterobacterial chromosomes using Tn7 without need for selection of the insertion event. *BMC Microbiol.* **6**:39.
28. Mellata, M., M. Dho-Moulin, C. M. Dozois, R. Curtiss III, P. K. Brown, P. Arne, A. Bree, C. Desautels, and J. M. Fairbrother. 2003. Role of virulence factors in resistance of avian pathogenic *Escherichia coli* to serum and in pathogenicity. *Infect. Immun.* **71**:536–540.
29. Mey, A. R., E. E. Wyckoff, L. A. Hoover, C. R. Fisher, and S. M. Payne. 2008. *Vibrio cholerae* VciB promotes iron uptake via ferrous iron transporters. *J. Bacteriol.* **190**:5953–5962.
30. Mills, M., and S. M. Payne. 1995. Genetics and regulation of heme iron transport in *Shigella dysenteriae* and detection of an analogous system in *Escherichia coli* O157:H7. *J. Bacteriol.* **177**:3004–3009.
31. Moss, J. E., T. J. Cardozo, A. Zychlinsky, and E. A. Groisman. 1999. The *selC*-associated SHI-2 pathogenicity island of *Shigella flexneri*. *Mol. Microbiol.* **33**:74–83.
32. Neilands, J. B. 1984. Siderophores of bacteria and fungi. *Microbiol. Sci.* **1**:9–14.
33. Oaks, E. V., M. E. Wingfield, and S. B. Formal. 1985. Plaque formation by virulent *Shigella flexneri*. *Infect. Immun.* **48**:124–129.
34. Payne, S. M. 1980. Synthesis and utilization of siderophores by *Shigella flexneri*. *J. Bacteriol.* **143**:1420–1424.
35. Payne, S. M., and R. A. Finkelstein. 1977. Detection and differentiation of iron-responsive avirulent mutants on Congo red agar. *Infect. Immun.* **18**:94–98.
36. Payne, S. M., D. W. Niesel, S. S. Peixotto, and K. M. Lawlor. 1983. Expression of hydroxamate and phenolate siderophores by *Shigella flexneri*. *J. Bacteriol.* **155**:949–955.
37. Payne, S. M., E. E. Wyckoff, E. R. Murphy, A. G. Oglesby, M. L. Boulette, and N. M. Davies. 2006. Iron and pathogenesis of *Shigella*: iron acquisition in the intracellular environment. *Biomaterials* **19**:173–180.
38. Perry, R. D., and C. L. San Clemente. 1979. Siderophore synthesis in *Klebsiella pneumoniae* and *Shigella sonnei* during iron deficiency. *J. Bacteriol.* **140**:1129–1132.
39. Philpott, D. J., J. D. Edgeworth, and P. J. Sansonetti. 2000. The pathogenesis of *Shigella flexneri* infection: lessons from in vitro and in vivo studies. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **355**:575–586.
40. Pupo, G. M., D. K. Karaolis, R. Lan, and P. R. Reeves. 1997. Evolutionary relationships among pathogenic and nonpathogenic *Escherichia coli* strains inferred from multilocus enzyme electrophoresis and *mdh* sequence studies. *Infect. Immun.* **65**:2685–2692.
41. Pupo, G. M., R. Lan, and P. R. Reeves. 2000. Multiple independent origins of *Shigella* clones of *Escherichia coli* and convergent evolution of many of their characteristics. *Proc. Natl. Acad. Sci. USA* **97**:10567–10572.
42. Purdy, G. E., and S. M. Payne. 2001. The SHI-3 iron transport island of *Shigella boydii* 0-1392 carries the genes for aerobactin synthesis and transport. *J. Bacteriol.* **183**:4176–4182.
43. Reeves, S. A. 2001. Iron acquisition in the intracellular environment of the host: multiple iron transport systems in *Shigella dysenteriae*. PhD. dissertation. University of Texas, Austin.
44. Rietsch, A., and J. Beckwith. 1998. The genetics of disulfide bond metabolism. *Annu. Rev. Genet.* **32**:163–184.
45. Rolland, K., N. Lambert-Zechovsky, B. Picard, and E. Denamur. 1998. *Shigella* and enteroinvasive *Escherichia coli* strains are derived from distinct ancestral strains of *E. coli*. *Microbiology* **144**(Pt. 9):2667–2672.
46. Runyen-Janecky, L., E. Dazenski, S. Hawkins, and L. Warner. 2006. Role and regulation of the *Shigella flexneri sit* and MntH systems. *Infect. Immun.* **74**:4666–4672.
47. Runyen-Janecky, L. J., and S. M. Payne. 2002. Identification of chromosomal *Shigella flexneri* genes induced by the eukaryotic intracellular environment. *Infect. Immun.* **70**:4379–4388.
48. Runyen-Janecky, L. J., S. A. Reeves, E. G. Gonzales, and S. M. Payne. 2003. Contribution of the *Shigella flexneri* Sit, Iuc, and Feo iron acquisition systems to iron acquisition in vitro and in cultured cells. *Infect. Immun.* **71**:1919–1928.
49. Sabri, M., S. Leveille, and C. M. Dozois. 2006. A SitABCD homologue from an avian pathogenic *Escherichia coli* strain mediates transport of iron and manganese and resistance to hydrogen peroxide. *Microbiology* **152**:745–758.
50. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
51. Sansonetti, P. J. 2006. The bacterial weaponry: lessons from Shigella. *Ann. N. Y. Acad. Sci.* **1072**:307–312.
52. Schmitt, M. P., and S. M. Payne. 1991. Genetic analysis of the enterobactin gene cluster in *Shigella flexneri*. *J. Bacteriol.* **173**:816–825.
53. Schwyn, B., and J. B. Neilands. 1987. Universal chemical assay for the detection and determination of siderophores. *Anal. Biochem.* **160**:47–56.
54. Torres, A. G., and S. M. Payne. 1997. Haem iron-transport system in enterohaemorrhagic *Escherichia coli* O157:H7. *Mol. Microbiol.* **23**:825–833.
55. Torres, A. G., P. Redford, R. A. Welch, and S. M. Payne. 2001. TonB-dependent systems of uropathogenic *Escherichia coli*: aerobactin and heme transport and TonB are required for virulence in the mouse. *Infect. Immun.* **69**:6179–6185.
56. Turbyfill, K. R., R. W. Kaminski, and E. V. Oaks. 2008. Immunogenicity and efficacy of highly purified in vivo complex vaccine from *Shigella flexneri* 2a. *Vaccine* **26**:1353–1364.
57. Vokes, S. A., S. A. Reeves, A. G. Torres, and S. M. Payne. 1999. The aerobactin iron transport system genes in *Shigella flexneri* are present within a pathogenicity island. *Mol. Microbiol.* **33**:63–73.
58. Wang, R. F., and S. R. Kushner. 1991. Construction of versatile low-copy-number vectors for cloning, sequencing and gene expression in *Escherichia coli*. *Gene* **100**:195–199.
59. Warner, P. J., P. H. Williams, A. Bindereif, and J. B. Neilands. 1981. ColV plasmid-specific aerobactin synthesis by invasive strains of *Escherichia coli*. *Infect. Immun.* **33**:540–545.
60. Wei, J., M. B. Goldberg, V. Burland, M. M. Venkatesan, W. Deng, G. Fournier, G. F. Mayhew, G. Plunkett III, D. J. Rose, A. Darling, B. Mau, N. T. Perna, S. M. Payne, L. J. Runyen-Janecky, S. Zhou, D. C. Schwartz, and F. R. Blattner. 2003. Complete genome sequence and comparative genomics of *Shigella flexneri* serotype 2a strain 2457T. *Infect. Immun.* **71**:2775–2786.
61. Wyckoff, E. E., D. Duncan, A. G. Torres, M. Mills, K. Maase, and S. M. Payne. 1998. Structure of the *Shigella dysenteriae* haem transport locus and its phylogenetic distribution in enteric bacteria. *Mol. Microbiol.* **28**:1139–1152.
62. Wyckoff, E. E., A. R. Mey, A. Leimbach, C. F. Fisher, and S. M. Payne. 2006. Characterization of ferric and ferrous iron transport systems in *Vibrio cholerae*. *J. Bacteriol.* **188**:6515–6523.
63. Zhou, D., W. D. Hardt, and J. E. Galan. 1999. *Salmonella typhimurium* encodes a putative iron transport system within the centisome 63 pathogenicity island. *Infect. Immun.* **67**:1974–1981.