

## *Pseudomonas aeruginosa* Evasion of Phagocytosis Is Mediated by Loss of Swimming Motility and Is Independent of Flagellum Expression<sup>∇†</sup>

Eyal Amiel,<sup>‡</sup> Rustin R. Lovewell,<sup>‡</sup> George A. O'Toole, Deborah A. Hogan, and Brent Berwin\*

Department of Microbiology and Immunology, Dartmouth Medical School, Lebanon, New Hampshire 03756

Received 10 February 2010/Returned for modification 18 March 2010/Accepted 27 April 2010

*Pseudomonas aeruginosa* is a pathogenic Gram-negative bacterium that causes severe opportunistic infections in immunocompromised individuals; in particular, severity of infection with *P. aeruginosa* positively correlates with poor prognosis in cystic fibrosis (CF) patients. Establishment of chronic infection by this pathogen is associated with downregulation of flagellar expression and of other genes that regulate *P. aeruginosa* motility. The current paradigm is that loss of flagellar expression enables immune evasion by the bacteria due to loss of engagement by phagocytic receptors that recognize flagellar components and loss of immune activation through flagellin-mediated Toll-like receptor (TLR) signaling. In this work, we employ bacterial and mammalian genetic approaches to demonstrate that loss of motility, not the loss of the flagellum *per se*, is the critical factor in the development of resistance to phagocytosis by *P. aeruginosa*. We demonstrate that isogenic *P. aeruginosa* mutants deficient in flagellar function, but retaining an intact flagellum, are highly resistant to phagocytosis by both murine and human phagocytic cells at levels comparable to those of flagellum-deficient mutants. Furthermore, we show that loss of MyD88 signaling in murine phagocytes does not recapitulate the phagocytic deficit observed for either flagellum-deficient or motility-deficient *P. aeruginosa* mutants. Our data demonstrate that loss of bacterial motility confers a dramatic resistance to phagocytosis that is independent of both flagellar expression and TLR signaling. These findings provide an explanation for the well-documented observation of nonmotility in clinical *P. aeruginosa* isolates and for how this phenotype confers upon the bacteria an advantage in the context of immune evasion.

*Pseudomonas aeruginosa* is an opportunistic Gram-negative bacterial pathogen that causes severe infections in immunocompromised patients and in the pulmonary compartment of patients suffering from cystic fibrosis (CF) (13, 14). In CF patients, disease severity is positively correlated with colonization by *P. aeruginosa* and the establishment of chronic infection. As part of the colonization process, the bacteria undergo a number of genetic changes that assist in their ability to survive in the mammalian host and to evade detection and clearance by the immune system (9, 21). One such change that has been phenotypically characterized for *P. aeruginosa* is loss of flagellar motility (12, 17). Furthermore, the loss of flagellar gene expression and motility function is associated with increased bacterial burdens and increased disease severity in CF patients (12, 17). While downregulation of flagellar expression has been inferred to confer a survival advantage on *P. aeruginosa* once it colonizes the host by evasion of both phagocytic receptors and TLR5-driven inflammatory signaling, the exact contribution of flagellum downregulation with respect to successful immune evasion is unclear (5, 17, 18).

Nonopsonic phagocytosis of *P. aeruginosa* by murine and human macrophages has previously been reported to require the expression of a flagellum, and the interpretation of these

results concluded that the flagellum is a necessary ligand for triggering phagocytic internalization of the bacteria (18). Furthermore, flagellar expression is reported to be critical for inducing inflammation during *P. aeruginosa* infection, and loss of flagellar gene expression results in impaired inflammatory responses and attenuated bacterial clearance (5). Here, we provide data that challenge the current paradigm that the flagellum functions as a primary phagocytic ligand for *P. aeruginosa* ingestion by immune cells with the formal demonstration that motility, rather than loss of flagellar expression, confers the advantage toward *P. aeruginosa* evasion of phagocytosis.

In these studies, we use *P. aeruginosa* motility-defective mutants to assess the role of bacterial motility in regard to phagocytic recognition by innate immune cells. When present in an aqueous environment, *P. aeruginosa* can swim via rotation of a single polar, monotrichous flagellum (27); there is currently no evidence that *P. aeruginosa* bacteria produce lateral flagella or alter their cell morphology as a direct function of motility. For the purposes of this report, motility refers to flagellum-based bacterial movement in an aqueous environment unless specifically indicated. Of note, *P. aeruginosa* is also capable of a flagellum-independent type of motility termed twitching in which type IV pilus filaments that extend from the cell body adhere to a surface and then retract, thus propelling the bacterium forward (23). Bacterial flagellar motility occurs through a motor complex that provides energy for rotational torque of a helical filament of repeating flagellin subunits that act as a propeller. The rotor, a multimer complex composed of FliG, FliM, and FliN, acts as a molecular switch and determines clockwise or counterclockwise rotation (27). In *P. aeruginosa*, the stator complex, which provides a stationary housing for the

\* Corresponding author. Mailing address: Department of Microbiology and Immunology, Dartmouth Medical School, Lebanon, NH 03756. Phone: (603) 650-6899. Fax: (603) 650-6223. E-mail: berwin@dartmouth.edu.

† Supplemental material for this article may be found at <http://iai.asm.org/>.

‡ E.A. and R.R.L. contributed equally to this report.

∇ Published ahead of print on 10 May 2010.

rotor, is composed of at least four partially redundant integral membrane proteins, MotAB and MotCD. Deletion of all four stators allows for flagellar assembly, but the structure cannot rotate and so the mutant is nonmotile (swimming and swarming defective) (27).

Previous reports have concluded that an intact flagellum is required for phagocytic recognition of *P. aeruginosa* (18). However, here we demonstrate that the phagocytic resistance exhibited by swimming motility-defective bacteria is not due to loss of flagellum-mediated activation of immune cells, since bacteria expressing a nonfunctional flagellum exhibit phagocytic resistance comparable to that of flagellum-deficient bacteria and loss of MyD88 signaling in phagocytic cells does not recapitulate this defect for phagocytosis. Rather, with the use of a variety of *in vitro*, *ex vivo*, and *in vivo* infection models, we show that loss of *P. aeruginosa* motility dramatically alters immune responses to these bacteria compared to those for motile isogenic bacterial strains and that it is the loss of flagellum-mediated motility, but not flagellum expression itself, that results in dramatic bacterial resistance to phagocytosis by murine and human phagocytes. These studies provide an explanation for the clinical observation that *P. aeruginosa* isolates obtained from CF hosts often exhibit a nonmotile phenotype and explain how this phenotype can confer a survival advantage for bacteria that modulate or lose their motility during an active infection.

#### MATERIALS AND METHODS

**Mice and cells.** C57BL/6 wild-type (WT) mice were obtained from The Jackson Laboratory (Bar Harbor, ME). MyD88<sup>-/-</sup> mice were generated by Adachi et al. (1). The bone marrow-derived dendritic cell (BMDC) culture protocol used is a modification of the work of Inaba et al. (11) as previously described (4). For these studies, the *Pseudomonas aeruginosa* clinical isolate PA14 is the parental bacterial strain and wild-type control for all of the isogenic mutants studied. Human peripheral blood monocytes were a generous gift from the lab of Paul Guyre (Dartmouth). Monocytes were differentiated into macrophages by culturing them in the presence of 10 µg/ml granulocyte-macrophage colony-stimulating factor (GM-CSF) for 7 days.

**FACS-based bacterial association assay for GFP<sup>+</sup> bacteria.** Bacterial strains expressing green fluorescent protein (GFP) were generated by transformation of the indicated strains with a multicopy plasmid (pSMC21 Amp<sup>r</sup> Kan<sup>r</sup> Carb<sup>r</sup> GFP<sup>+</sup>) that constitutively expresses GFP under the control of a derivative of the P<sub>tac</sub> promoter (6, 15). C57BL/6 BMDCs were incubated with the indicated GFP<sup>+</sup> bacterial strains for 45 min at 37°C or 4°C, as indicated in the text. For bacterial binding assays, BMDCs were preincubated in 10 µM cytochalasin D (Sigma) in serum-free Hanks balanced salt solution (HBSS) for 60 min at 37°C. Coincubation between BMDCs and the indicated GFP<sup>+</sup> bacterial strains took place in the presence of 10 µM cytochalasin D for 45 min at 37°C. Cells were washed thoroughly in serum-free HBSS and then analyzed by fluorescence-activated cell sorting (FACS) for the acquisition of fluorescence as an indication of BMDC association with the bacteria. The mean fluorescence intensities of the BMDC populations were assessed and graphed to obtain relative efficiency of cellular association with the different bacterial strains.

**Microscopy.** BMDCs from B6.Cg-Tg(CAG-mRFP1) mice (C57BL/6 background; purchased from Jackson Labs), which express red fluorescent protein (RFP) from the actin promoter, were cultured as previously described (5). Cells were coincubated with GFP-expressing *P. aeruginosa* strains as indicated for 45 min at 37°C at a multiplicity of infection (MOI) of ~10. Cells were washed twice in 400 µl of serum-free HBSS prior to a 10-min cytospin onto glass slides at 1,000 rpm. Cells were visualized via fluorescence and differential interference contrast (DIC) microscopy. Microscopy was performed on a Zeiss LSM510 Meta microscope using a 63× lens, followed by image analysis on the LSM5 Image Browser software.

**Bacterial motility swimming assay.** Bacterial motility was examined as described elsewhere (7, 27). Briefly, 0.3% agar plates were poured and the indicated bacterial strains were inoculated from liquid cultures onto the plates by

puncturing inoculates halfway through the depth of the agar. Plates were incubated at room temperature for 48 h and monitored for the generation of bacterial halos as an indication of swimming motility. Plates were subsequently photographed with a digital camera to allow comparison of the relative swimming motilities of the different bacterial strains.

**Western Blot analyses and Coomassie blue staining.** Analysis of total flagellin expression by the indicated bacterial strains was performed by lysis of the whole bacteria in SDS sample buffer containing 5% β-mercaptoethanol. Analysis of the presence of extracellular flagella on the Δ*motAB* Δ*moiCD* mutant was performed as a modification of published methodology (19, 25). Briefly, a 100-ml culture of bacteria was pelleted by centrifugation, resuspended in phosphate-buffered saline (PBS), and sheared for 3 min in a blender. The sheared suspension was centrifuged at 8,000 × *g* for 10 min, and the protein within the supernatant was subsequently precipitated with the addition of trichloroacetic acid (TCA) to 20%. The precipitated pellet was washed with acetone and resuspended in SDS sample buffer containing 5% β-mercaptoethanol. For both the bacterial lysates and the isolated flagellin, the samples were boiled and separated on 12% SDS-PAGE gels. Parallel lanes were used either for Coomassie blue staining or for transfer onto Immobilon-P membranes (Millipore Corp.) for Western analysis. Following blotting with polyclonal anti-FliC antibody (22), Western blots were developed using enhanced chemiluminescence (ECL) (Amersham Biosciences) of horseradish peroxidase (HRP)-conjugated secondary antibodies (Jackson Immunoresearch).

***In vitro* gentamicin protection assays.** Phagocytosis of live *P. aeruginosa* bacteria was performed as a modified version of published protocols (4, 8) and as previously described (2). Briefly, overnight cultures of *P. aeruginosa* were washed and resuspended in serum-free medium and bacterial concentrations were determined. BMDCs of the indicated genotype (2 × 10<sup>5</sup>) were incubated with bacteria at an MOI of ~10 for 45 min at 37°C, followed by incubation in 100 µg/ml gentamicin for 20 min at 37°C. For intracellular killing assays, a modification of the previous protocol (3) was used: following a 1-h coincubation of PA14 bacteria and BMDCs, gentamicin was added and, for the following 60 min, aliquots were assayed as described above for remaining CFU. In all cases, cells were washed and subsequently lysed in 500 µl 0.1% Triton X-100 in PBS. Lysates were plated onto *Pseudomonas* isolation agar and incubated overnight at 37°C. The next day, colonies were counted and relative phagocytosis was determined by CFU counts.

***Ex vivo* peritoneal phagocyte gentamicin protection assay and FACS analyses.** Phagocytosis of live *P. aeruginosa* bacteria was performed as a modified version of previously described protocols (4, 8). Overnight cultures of *P. aeruginosa* were washed twice in 1 ml serum-free HBSS and resuspended in 10 ml HBSS. Bacterial concentrations were determined based on estimated ~1 × 10<sup>9</sup> CFU/ml saturation levels. Naïve C57BL/6 mice were injected intraperitoneally (i.p.) with 1 ml of 4% thioglycolate and subsequently sacrificed 4 days later. The peritoneal cavity was lavaged with 6 ml of serum-free HBSS. The lavage fluid was centrifuged, and pelleted cells were washed twice in serum-free HBSS before being resuspended in 2 ml serum-free HBSS. For the gentamicin protection assay, peritoneal phagocytes and bacteria were subsequently treated as described above for *in vitro* gentamicin protection assays. Alternatively, peritoneal phagocytes were coincubated with GFP-expressing bacteria for 45 min at 37°C. Cells were washed twice in serum-free HBSS and resuspended in 250 µl HBSS. Bacterial association with phagocytes was determined via FACS analysis as described previously.

***Ex vivo* lung phagocyte gentamicin protection assay and FACS analyses.** Phagocytosis of live *P. aeruginosa* bacteria was performed as a modified version of previously described protocols (4, 8). Overnight cultures of *P. aeruginosa* were washed twice in 1 ml serum-free HBSS and resuspended in 10 ml serum-free HBSS. Bacterial concentrations were determined based on estimated ~1 × 10<sup>9</sup> CFU/ml saturation levels. Naïve C57BL/6 mice were treated with 150 µg lipopolysaccharide (Sigma) in PBS via tracheobronchial aspiration and were subsequently sacrificed 3 days later. The lungs were lavaged with 800 µl of PBS with 5 mM EDTA. The bronchoalveolar lavage (BAL) fluid was centrifuged, and pelleted lung cells were washed twice in 400 µl serum-free HBSS before being incubated with the indicated bacteria at an MOI of ~10 for 45 min at 37°C. For the gentamicin protection assay, alveolar phagocytes and bacteria were subsequently treated as described above for *in vitro* gentamicin protection assays. Alternatively, alveolar phagocytes were coincubated with GFP-expressing bacteria for 45 min at 37°C. Cells were washed twice in serum-free HBSS and resuspended in 250 µl HBSS. Bacterial association with lung phagocytes was determined via FACS analysis as described previously.

***In vivo* bacterial uptake assay.** Live *P. aeruginosa* bacteria (5 × 10<sup>6</sup> or bacterial numbers as indicated) either were injected intraperitoneally or were oropharyngeally aspirated into C57BL/6 mice. Mice were sacrificed 1 h postinjection, and

TABLE 1. *P. aeruginosa* genetic mutants used in these studies

| PA14 mutant <sup>a</sup>                           | WT function of gene product       | Mutant phenotype for bacterial motility |
|--|-----------------------------------|---|
| <i>pilB</i> mutant                                 | ATPase for pilus assembly         | Lacks pili, twitch defective            |
| <i>pilG</i> mutant                                 | Pilus transcription factor        | Lacks pili, twitch defective            |
| <i>flgK</i> mutant                                 | Flagellar hook protein            | No flagellum, swimming defective        |
| <i>fliN</i> mutant                                 | Flagellar export/assembly protein | No flagellum, swimming defective        |
| $\Delta$ <i>motAB</i> $\Delta$ <i>motCD</i> mutant | Flagellar stator proteins         | Flagellum intact, swimming defective    |

<sup>a</sup> PA14 is a nonmucoid WT clinical isolate.

peritoneal or BAL fluid lavages, respectively, were collected. Cells were pelleted, washed twice in serum-free HBSS, and resuspended in 750  $\mu$ l of HBSS. Five hundred microliters of each suspension was incubated in the presence of gentamicin for 20 min at 37°C to kill noninternalized bacteria. Cells were washed twice in serum-free HBSS and resuspended in 500  $\mu$ l 0.1% Triton X-100 in PBS. Fifteen microliters of resuspended cells/bacteria was plated on *Pseudomonas* isolation agar plates and incubated overnight at 37°C. Colonies were counted the following morning, and CFU were calculated based on fraction of total sample plated. For peritoneal *in vivo* phagocytosis studies, analyses of total cell numbers and phenotypes were done in parallel. Total cell numbers harvested from peritoneal lavage samples were quantified using trypan blue staining and manual cell counts of viable cells using a hemacytometer.

**Data and statistical analyses.** Sample sizes for each experiment varied and are noted in the text. For all graphs, means and standard deviations are shown. As indicated, one-way analysis of variance (ANOVA) with Tukey's multiple comparison posttest or unpaired Student's *t* test analysis was performed to assess statistical significance of the data. In the figures, statistical significance is represented by an asterisk and indicates  $P \leq 0.05$ .

## RESULTS

***P. aeruginosa* strains lacking a flagellum or flagellar stator proteins are impaired in swimming motility.** In these studies, a variety of *P. aeruginosa* mutants were used to assess and quantitatively compare the effect of either loss of the flagellum or flagellar function on the phagocytic recognition of these bacteria by immune cells (Table 1). PA14 is a nonmucoid wild-type *P. aeruginosa* clinical isolate and is the parental strain for all of the isogenic mutants described in these studies (16, 20). Two mutants (*pilB* and *pilG*) defective for type IV pilus expression and function were used as controls as strains that are impaired in twitching motility but are fully competent at swimming motility. Two different mutations (*flgK*, which codes for a subunit of the flagellar rotor complex) that are phenotypically impaired in proper flagellum expression were used, as well as a quadruple mutant for four flagellar stator proteins ( $\Delta$ *motAB*  $\Delta$ *motCD*) which results in intact flagellum expression but nonfunctional swimming motility (26, 27). To confirm the swimming motility phenotype of the mutant strains analyzed, we performed a standard bacterial swimming assay whereby liquid bacterial cultures are inoculated by stabbing puncture into 0.3% agar plates (see Fig. S1 in the supplemental material). Zones surrounding the point of inoculation were evaluated as an indication of swimming motility (27). In accordance with previous results, we observed complete lack of swimming motility in the flagellum-deficient mutants and the stator-deficient mutant while we observed a swimming phenotype in the

parental PA14 strain and the two pilus-deficient mutant strains (Fig. S1) (26, 27).

**Swimming-deficient *P. aeruginosa* strains are resistant to phagocytosis by BMDCs *in vitro*.** To evaluate the effect of loss of bacterial motility on phagocyte recognition, we transformed parental PA14 as well as flagellum-deficient *flgK* and *fliN* mutants, the  $\Delta$ *motAB*  $\Delta$ *motCD* flagellar stator protein knockout, and the pilus-deficient *pilG* mutant with a GFP expression construct to allow FACS-based analysis of phagocyte-bacterium interactions. The intensity of GFP fluorescence emitted by our transformed strains did not vary significantly among the strains (see Fig. S2 in the supplemental material), allowing us to compare binding and uptake of these different strains directly in relation to each other. Using fluorescence as a quantitative indicator of bacterial association with BMDCs, we observed a marked and significant reduction in the ability of BMDCs to associate with flagellum-deficient strains (*flgK* and *fliN*) compared to the parental PA14 bacteria (Fig. 1A and D), though the pilus-deficient mutant (*pilG*) associated with the BMDCs at levels comparable to those for the WT (Fig. 1A). To more rigorously assess the differential phenotypes observed with the GFP-expressing bacteria, we utilized a gentamicin protection assay to specifically and quantitatively assess *in vitro* phagocytosis of the respective bacterial strains. Both flagellum-deficient strains exhibited a dramatic  $\sim$ 100-fold reduction in their phagocytic recognition by BMDCs *in vitro* compared to the parental PA14 strain (Fig. 1B). To verify that this 100-fold variance was not due to intracellular killing of *P. aeruginosa*, we measured the intracellular killing rate of phagocytosed PA14. Approximately 50% of ingested PA14 bacteria were killed after 2 h of incubation with BMDCs, indicating that the 100-fold differential in phagocytosis observed in a shorter duration (45 min) is not within the margin of the killing rate (Fig. 2C).

With the use of both assays, we next tested if phagocyte recognition and engulfment of *P. aeruginosa* were dependent on a fully assembled flagellum by coincubating BMDCs with the  $\Delta$ *motAB*  $\Delta$ *motCD* flagellar stator mutant and assaying via flow cytometry and gentamicin protection. The  $\Delta$ *motAB*  $\Delta$ *motCD* mutant was substantially and significantly reduced in both its overall association (FACS assay [Fig. 1C, left]) and subsequent engulfment (gentamicin protection assay [Fig. 1C, right]) by BMDCs, similarly to the *flgK* and *fliN* mutants (compare Fig. 1A and B with C). These results were then confirmed by fluorescence microscopy, which similarly demonstrated the dramatic deficit in the association of swimming-defective but flagellum-expressing *P. aeruginosa* with BMDCs *in vitro* (Fig. 1E). This deficit was not due to observable differences in total flagellar expression between the  $\Delta$ *motAB*  $\Delta$ *motCD* mutant and the parental wild type as detected by Coomassie blue staining and Western blot analysis (Fig. 2A), and in biochemical confirmation of a previous report utilizing microscopy (27), we demonstrate that the  $\Delta$ *motAB*  $\Delta$ *motCD* mutant has an intact extracellular flagellum (Fig. 2B). The observation that the  $\Delta$ *motAB*  $\Delta$ *motCD* mutant, but not the *pilG* mutant, recapitulates the dramatic phagocytic deficit seen for the flagellum-deficient strains demonstrates that the resistance to phagocytosis by these strains is not due to loss of an intact flagellum or pili but the loss of functional flagellum-based motility. Additionally, this deficit is specific to flagellum-based swimming




FIG. 1. *P. aeruginosa* mutants deficient for flagellum or flagellar stator proteins are resistant to phagocytosis by BMDCs *in vitro*. (A) C57BL/6 BMDCs were analyzed by FACS for relative association with GFP-transformed PA14, the pilus-deficient *pilG* mutant, or the flagellum-deficient *flgK* and *fliN* mutants. (B) C57BL/6 BMDCs were assayed by gentamicin protection assay for relative phagocytic uptake of PA14 or flagellum-deficient *flgK* and *fliN* mutants. The graph is plotted on a logarithmic scale. (C) FACS analysis (left) and gentamicin protection assay (right) of C57BL/6 BMDCs coincubated with either WT PA14 or the  $\Delta motAB \Delta motCD$  flagellar stator mutant. (D) FACS histogram of total BMDC cellular association with GFP-transformed *P. aeruginosa* strains after 45 min of coincubation. (E) Fluorescence microscopy of RFP-expressing BMDCs coincubated with GFP-transformed PA14 or the  $\Delta motAB \Delta motCD$  mutant viewed with DIC overlay at  $\times 65$  magnification. For all graphs, phagocytic uptake levels were normalized as percentages of the mean WT phagocytosis. For all genotypes,  $n$  is  $\geq 9$ ; means, standard deviations, and statistical significance (asterisks) are shown.

motility, since loss of pilus-based twitching motility did not confer measurable phagocytic evasion.

***P. aeruginosa* mutants deficient for flagellum or flagellar stator proteins are resistant to phagocytosis by human macrophages *in vitro*.** To evaluate whether swimming-defective *P. aeruginosa* strains were also resistant to phagocytosis by human cells, we generated *in vitro* peripheral blood mononuclear cell-derived human macrophages to assay them by gentamicin pro-

tection assay for the relative phagocytic uptake of PA14 and its isogenic *pilB*, *pilG*, *flgK*, *fliN*, and  $\Delta motAB \Delta motCD$  mutants (Fig. 3). The results with human macrophages recapitulated our data with murine BMDCs, whereby we observed a 30- to 100-fold impairment for the phagocytic recognition of swimming-defective *P. aeruginosa* strains by *in vitro*-cultured human macrophages compared to recognition of the wild-type parental bacterial strain (Fig. 3). The recapitulation of the murine




FIG. 2. Phagocytic resistance of nonswimming *P. aeruginosa* is not due to bactericidal activity or differential flagellar expression. (A) WT PA14 (PA14) and the  $\Delta$ *motAB*  $\Delta$ *motCD* (*mot*) strain express similar total levels of flagellin as assessed by Western analysis for the FliC protein (right). Coomassie blue staining of parallel lanes is shown as a control for protein load (left). (B) To confirm that the  $\Delta$ *motAB*  $\Delta$ *motCD* mutant strain has intact extracellular flagella, the flagella were mechanically sheared from the bacteria and assessed by Coomassie blue staining (left) and Western analysis (right). (C) The BMDC killing rate of *P. aeruginosa* was assayed by gentamicin protection assay. Following a 1-h coincubation of PA14 with BMDCs, gentamicin was added, and at the indicated time points following gentamicin addition, aliquots were harvested and lysed to assess the death of internalized bacteria over the course of the assay. CFU were plotted relative to initial recovery. For each time point,  $n \geq 6$ ; standard deviations are shown.

data with human macrophages indicates that the loss of motility by *P. aeruginosa* bacteria is likely a universal evasion strategy from phagocytosis by immune cells that spans multiple cell types and species.

**Loss of MyD88 signaling fails to recapitulate the phagocytic deficit for a flagellum-deficient *P. aeruginosa* strain.** Our data with the  $\Delta$ *motAB*  $\Delta$ *motCD* *P. aeruginosa* strain indicated that the deficit for flagellum-deficient strains is not an intrinsic function of lack of flagellum expression itself or the loss of flagellum-mediated activation of immune cells. To test this hypothesis more stringently, we generated BMDCs from MyD88-deficient (MyD88<sup>-/-</sup>) mice to determine if loss of MyD88 signaling recapitulates the phagocytic deficit observed for flagellum-deficient bacteria. We analyzed the relative association of the GFP-expressing wild-type PA14, *flgK*, and  $\Delta$ *motAB*  $\Delta$ *motCD* strains with WT or MyD88<sup>-/-</sup> BMDCs in parallel. Loss of MyD88 signaling did not change the relative level of bacterial association with BMDCs (Fig. 4A). Furthermore, WT and MyD88<sup>-/-</sup> BMDCs quantitatively assayed in parallel for *in vitro* phagocytosis of wild-type or swimming-defective *P. aeruginosa* supported the FACS analysis in that loss of MyD88 signaling did not recapitulate the phagocytic deficit seen with *flgK* and  $\Delta$ *motAB*  $\Delta$ *motCD* bacteria (Fig. 4B). These results strongly support the idea that the phagocytic




FIG. 3. *P. aeruginosa* mutants deficient for flagellum or flagellar motor proteins are resistant to phagocytosis by human macrophages *in vitro*. Peripheral blood mononuclear cell (PBMC)-derived human macrophages were cultured and assayed by gentamicin protection assay for relative *in vitro* uptake of PA14, pilus-deficient *pilB* and *pilG* mutants, flagellum-deficient *flgK* and *fliN* mutants, and the flagellar stator protein-deficient  $\Delta$ *motAB*  $\Delta$ *motCD* mutant. Phagocytic uptake levels were normalized as percentages of the mean WT phagocytosis. Statistical significance ( $P < 0.05$ ) of differences from wild-type levels is indicated (asterisks). Graphs are plotted on a log scale. For all genotypes,  $n \geq 9$ ; means and standard deviations are shown.

deficit is independent of Toll-like receptor (TLR)/MyD88 signaling during immune recognition of flagellum-deficient bacteria.

**Loss of motility is sufficient to confer *P. aeruginosa* phagocytic resistance to primary peritoneal and alveolar murine cells *ex vivo*.** Having identified a deficit for the phagocytosis of stator protein-deficient *P. aeruginosa* strains by murine BMDCs and human cultured macrophages *in vitro*, we next tested whether or not this observation could be recapitulated by primary cells *ex vivo*. Peritoneal phagocytes were harvested and incubated with the various GFP-transformed bacterial strains, and cellular association with the bacteria was analyzed by FACS (Fig. 5A). Peritoneal phagocytes were also assayed for their ability to phagocytose bacteria, assessed by gentamicin protection (Fig. 5B). Consistent with our data derived from *in vitro*-cultured cells, nonmotile *P. aeruginosa* strains were dramatically resistant to phagocytosis by *ex vivo* primary peritoneal macrophages. As a control to assess whether this massive phagocytic differential was merely reflective of a cell surface binding deficit, we measured cell surface binding of bacteria to murine peritoneal macrophages pretreated with cytochalasin D to inhibit phagocytic uptake. Cytochalasin D treatment of *P. aeruginosa* did not alter its motility (data not shown). No statistical difference in *P. aeruginosa* cell surface binding was observed among the bacterial genotypes (see Fig. S3 in the supplemental material); this is in direct contrast to parallel association experiments using untreated, phagocytically competent macrophages (Fig. 5A). These data support the idea that it is the evasion of phagocytic engulfment and not initial bacterium-phagocyte contact that provides for the phagocytic resistance phenotype of nonmotile *P. aeruginosa*.




FIG. 4. The phagocytic deficiency for nonmotile *P. aeruginosa* strains is independent of MyD88 signaling. (A) WT or MyD88<sup>-/-</sup> BMDCs were assayed for relative *in vitro* cellular association of GFP-PA14, GFP-*flgK*, or GFP- $\Delta$ *motAB*  $\Delta$ *motCD* *P. aeruginosa* strains by FACS analysis. Data are represented as the mean fluorescence intensity (MFI) of each experimental group. (B) WT or MyD88<sup>-/-</sup> BMDCs were assayed for *in vitro* phagocytosis of PA14 or *flgK* or  $\Delta$ *motAB*  $\Delta$ *motCD* *P. aeruginosa* strains by gentamicin protection assay. Phagocytic uptake levels were normalized as percentages of mean WT PA14 phagocytosis. The graph is plotted on a logarithmic scale. For all genotypes,  $n$  is  $\geq 6$  and means, standard deviations, and statistical significance (asterisks) are shown.

The lungs are a major site of clinical infection by *P. aeruginosa* (14, 21), and the prevalence of pulmonary *P. aeruginosa* isolates lacking motility correlates with poorer clinical condition in CF patients (12). Therefore, to address whether our previous results extended to pulmonary cells, we assayed bacterial association and phagocytosis by primary lung phagocytes *ex vivo*. We harvested murine lung cells by bronchial lavage and assessed phagocyte association with GFP-expressing bacteria by FACS analysis (Fig. 5C) and assayed for bacterial uptake by gentamicin protection assay (Fig. 5D). In these assays, flagellum-deficient and stator protein-deficient *P. aeruginosa* strains were highly resistant to phagocytosis and overall association with *ex vivo* alveolar cells. These findings demonstrate that the loss of bacterial motility by *P. aeruginosa* confers resistance to phagocytosis by multiple types of primary murine immune cells.

**Stator protein-deficient *P. aeruginosa* strains are resistant to phagocytosis *in vivo*.** Having determined that loss of motility protects *P. aeruginosa* from phagocytosis *in vitro* and *ex vivo*, we next tested the *in vivo* relevance of our findings. To address whether loss of motility in *P. aeruginosa* enables immune evasion during infection of a host organism, we determined the relative clearance of the PA14, *flgK*, and  $\Delta$ *motAB*  $\Delta$ *motCD*

bacterial strains by both peritoneal and pulmonary phagocytes *in vivo*. With the use of a previously described *in vivo* gentamicin protection assay (2), we observed a dramatic 10-fold reduction in the ability of peritoneal phagocytes to ingest *flgK* and  $\Delta$ *motAB*  $\Delta$ *motCD* mutants *in vivo* compared to uptake of the parental PA14 strain (Fig. 6A). The differences in the phagocytosis of the various bacterial strains were not due to differential recruitment of cells to the peritoneal cavity during the assay, since quantification of total peritoneal cells from treated mice showed no discrepancy in total peritoneal cell numbers between mice injected with the parental *P. aeruginosa* strain and mice injected with the mutant *P. aeruginosa* strains (Fig. 6B, bottom). Extending these findings, and with direct clinical relevance to *P. aeruginosa* colonization of the lung, we assessed the relative clearance of the PA14 and  $\Delta$ *motAB*  $\Delta$ *motCD* bacterial strains by pulmonary phagocytes *in vivo*. Consistent with our findings with peritoneal phagocytes, lung phagocytes ingested motile PA14 10-fold better than did nonmotile PA14 (Fig. 6B, top). From these results, we conclude that downregulation of motility, independent of structural loss of flagella, by *P. aeruginosa* confers a dramatic resistance to recognition by the host immune system *in vivo* and that this provides a mechanism for immune evasion.

## DISCUSSION

*Pseudomonas aeruginosa* is an opportunistic Gram-negative pathogenic bacterium that poses a significant clinical concern for immunocompromised individuals (9, 13, 14, 21). Two prominent clinical correlations that have been identified in CF patients are a strong positive correlation between the extent of the *P. aeruginosa* infection and the severity of the prognosis for the patient (12, 17) and a marked phenotypic shift toward the loss of bacterial motility and downregulation of flagellum expression that correlates with the severity of CF disease (12, 17). Therefore, there is considerable interest in understanding the basis for the latter phenotypic observation, especially in relation to the advantage that it may confer on the bacteria and, consequently, how we may more effectively treat the infection.

Previous reports have shown that loss of flagellar expression in clinical bacterial isolates results in dramatic decreases in nonopsonic phagocytic recognition by murine and human macrophages *in vitro* (17, 18). Other reports have indicated that the flagellum itself, and not the loss of bacterial motility, is the primary determinant in controlling the host immune response to *P. aeruginosa* (5). In these studies we have utilized a bacterial and mammalian genetic approach to show that *P. aeruginosa* strains that are defective for swimming motility are resistant to phagocytosis *in vitro* and *in vivo*. Furthermore, we demonstrate that this resistance to phagocytosis is independent of flagellum expression and not dependent on MyD88 signaling, in response to these bacteria. Our results indicate that it is the functional loss of flagellum-mediated swimming motility in *P. aeruginosa* that confers phagocytic resistance on these bacteria. These data are the first demonstration that the loss of flagellar function, not the loss of the flagellum apparatus as an activation or association ligand, is responsible for conferring resistance to phagocytosis and that this phenotype has an important role in immune evasion by *P. aeruginosa* of both mouse and human immune cells.




FIG. 5. Motor protein-deficient *P. aeruginosa* strains are resistant to phagocytosis by *ex vivo* primary phagocytic cells from the peritoneum and the lung. (A) C57BL/6 peritoneal exudate cells were harvested by lavage and assayed by FACS *ex vivo* for relative cellular association with the GFP-expressing bacterial strains indicated. MFI, mean fluorescence intensity. (B) Peritoneal phagocytes were assayed for relative phagocytosis of WT PA14, *flagK*, and  $\Delta$ *motAB*  $\Delta$ *motCD* bacteria by gentamicin protection assay. (C) C57BL/6 lung phagocytes were harvested via bronchoalveolar lavage and subsequently assayed by FACS *ex vivo* for relative cellular association with GFP-transformed bacteria as indicated. (D) Lung phagocytes were assayed by gentamicin protection assay *ex vivo* for relative uptake of WT, flagellum-deficient mutant *flagK*, and flagellar motor protein-deficient mutant  $\Delta$ *motAB*  $\Delta$ *motCD* bacteria. For all graphs,  $n$  is  $\geq 4$  and means, standard deviations, and statistical significance (asterisks) are shown.

Based on the observations that *P. aeruginosa* tends to down-regulate flagellar gene expression and functional motility in CF patients, it has been proposed that *P. aeruginosa* undergoes these phenotypic changes as part of an immune evasion strategy to avoid detection by phagocytic and cell-activating receptors during chronic infections (12, 17). Furthermore, nonopsonic phagocytosis of *P. aeruginosa* has been reported to require the expression of an intact flagellum, as flagellum-deficient bacteria are reported to be resistant to phagocytosis by murine and human macrophages *in vitro* (18). The  $\Delta$ *motAB*  $\Delta$ *motCD* strain used in these studies is deficient for four stator genes which comprise the stationary components of the *P. aeruginosa* flagellar motor (27). In this strain, flagellar structures are intact on the surface of the bacteria but are not functional and thus these bacteria are impaired in swimming motility (27). Based on previous electron micrograph studies and our confirmatory biochemical studies here, the flagellar structure in the  $\Delta$ *motAB*  $\Delta$ *motCD* mutant is intact and does not appear to be different from its wild-type counterpart (27). By demonstrating that the  $\Delta$ *motAB*  $\Delta$ *motCD* strain is equally as resistant to phagocytosis as the flagellum-deficient strains, we conclude that the flagellum itself is not a critical ligand for nonopsonic phagocytosis of *P. aeruginosa* in our experimental system. Intriguingly, a previous report found that loss of the stator genes *motAB* in a different human clinical *P. aeruginosa* isolate resulted in resistance to phagocytosis by macrophages

*in vitro* (24). However, this did not affect measurable motility and the authors noted that the parental strain used for their studies exhibited an abnormal phenotype which possibly could confound the interpretation of the data (24). In the PA14 *P. aeruginosa* strain used in our studies, only loss of all four bacterial stator proteins ( $\Delta$ *motAB*  $\Delta$ *motCD*) results in swimming motility defects under standard experimental conditions (27) and the isogenic  $\Delta$ *motAB* PA14 mutant did not recapitulate the phagocytic defect observed in the  $\Delta$ *motAB*  $\Delta$ *motCD* mutant (data not shown).

TLR5 has been recognized as a major pattern recognition receptor (PRR) in innate immune cells, binding bacterial flagellin and activating the MyD88 cell signaling pathway (10, 25). We observed only a minor deficit for *P. aeruginosa* phagocytosis in MyD88<sup>-/-</sup> BMDCs that did not recapitulate the dramatic deficit seen for phagocytic recognition of flagellum-deficient bacteria. Similarly, bacterial association with MyD88<sup>-/-</sup> BMDCs was not any lower than that with WT BMDCs. These findings are particularly relevant since previously published hypotheses have argued that flagellum-deficient bacteria escape immune detection primarily through the loss of flagellum-mediated immune activation of TLR5 (5, 17, 18). Because activation via TLR5 is MyD88 dependent, we believe that it is not a major contributor in the phagocytic resistance phenomenon observed in the nonmotile PA14 mutants. This is supported by similar degrees of phagocytic resis-




FIG. 6. Flagellum-deficient and stator protein-deficient *P. aeruginosa* strains are resistant to phagocytosis *in vivo*. (A) (Top) An *in vivo* gentamicin protection assay was used to determine relative uptake of PA14, *flagK*, and  $\Delta$ *motAB*  $\Delta$ *motCD* bacterial strains by peritoneal phagocytes. Phagocytic uptake levels were normalized as percentages of the mean WT phagocytosis levels. (Bottom) Total peritoneal cells were quantified from lavage samples of mice treated with PA14, *flagK*, or  $\Delta$ *motAB*  $\Delta$ *motCD* bacteria, to control for differential immune cell recruitment by the different bacterial strains. (B) Gentamicin protection assay results following *in vivo* oropharyngeal aspiration of WT or  $\Delta$ *motAB*  $\Delta$ *motCD* PA14 are consistent with the data shown in panel A, with nonmotile PA14 being ~10-fold more resistant to phagocytosis than motile PA14. Individual data points (mice), means, and statistical significance (asterisks) are shown.

tance in both flagellated and nonflagellated nonmotile strains. However, it is reasonable to suggest that TLR5 and other MyD88-dependent PRRs likely do play a role in other aspects of the overall immune response to PA14 infection. Indeed, it has been demonstrated with an alternative *P. aeruginosa* strain that the TLR5-flagellin interaction is sufficient to activate alveolar macrophage upregulation of cytokine production (5).

Additionally, we have previously reported that cytokine responses to bacterial challenge are not necessarily directly coupled to the phagocytosis of those bacteria (2). Overall, it is now clear that effective bacterial clearance of nonopsonized PA14 via phagocytosis hinges on motility-based factors outside of simple recognition of flagellin as a stimulating ligand.

In summary, our data demonstrate that loss of swimming motility in *P. aeruginosa* bacteria results in dramatic resistance to phagocytic recognition by innate immune cells *in vitro* and *in vivo*. This phagocytic resistance was observed for both mouse and human cells and was not due to the specific loss of flagellar expression, since bacteria that expressed nonfunctional flagella were able to recapitulate the same phagocytic resistance exhibited by flagellum-deficient bacteria. Furthermore, genetic deletion of MyD88 in phagocytic cells was not able to recapitulate the defect for phagocytosis of flagellum-deficient bacteria, demonstrating that loss of flagellum-mediated TLR5 signaling is not responsible for the impaired phagocytosis of flagellum-deficient *P. aeruginosa* bacteria. We found that motility-deficient *P. aeruginosa* strains are resistant to phagocytosis by both human and mouse phagocytes *in vitro*, as well as primary mouse cells *ex vivo* and *in vivo*. These data provide new insights into important clinical observations about *P. aeruginosa* and the loss of bacterial motility and are the first demonstration that *P. aeruginosa* strains defective for swimming motility are resistant to phagocytosis *in vivo*. These findings provide an important new understanding of the immunological advantage *in vivo* for the loss of motility by *P. aeruginosa* during chronic infections.

#### ACKNOWLEDGMENTS

We thank Shizuo Akira (Osaka University) and Paul Guyre, Harry Higgs, and J. Ma Collins (Dartmouth) for reagents; Julie Acker and Ryan Collins for technical assistance; the Dartmouth Medical School FACS and microscopy core facilities; and Matt Wargo, Laurie Whitaker, and Jenna Allard (UVM) for technical advice.

This research was supported by RO1 AI067405 and a CF Foundation RDP training grant (B.B.) and NIH training grants T32 AI07363 (E.A.) and NIGMS GM008704 (R.R.L.).

#### REFERENCES

- Adachi, O., T. Kawai, K. Takeda, M. Matsumoto, H. Tsutsui, M. Sakagami, K. Nakanishi, and S. Akira. 1998. Targeted disruption of the MyD88 gene results in loss of IL-1- and IL-18-mediated function. *Immunity* 9:143–150.
- Amiel, E., J. L. Acker, R. M. Collins, and B. Berwin. 2009. Uncoupling scavenger receptor A-mediated phagocytosis of bacteria from endotoxic shock resistance. *Infect. Immun.* 77:4567–4573.
- Amiel, E., A. Alonso, S. Uematsu, S. Akira, M. E. Poynter, and B. Berwin. 2009. Pivotal advance: Toll-like receptor regulation of scavenger receptor-A-mediated phagocytosis. *J. Leukoc. Biol.* 85:595–605.
- Amiel, E., S. Nicholson-Dykstra, J. J. Walters, H. Higgs, and B. Berwin. 2007. Scavenger receptor-A functions in phagocytosis of *E. coli* by bone marrow dendritic cells. *Exp. Cell Res.* 313:1438–1448.
- Balloy, V., A. Verma, S. Kuravi, M. Si-Tahar, M. Chignard, and R. Ramphal. 2007. The role of flagellin versus motility in acute lung disease caused by *Pseudomonas aeruginosa*. *J. Infect. Dis.* 196:289–296.
- Bloemberg, G. V., G. A. O'Toole, B. J. Lugtenberg, and R. Kolter. 1997. Green fluorescent protein as a marker for *Pseudomonas* spp. *Appl. Environ. Microbiol.* 63:4543–4551.
- Braun, T. F., S. Poulson, J. B. Gully, J. C. Empey, S. Van Way, A. Putnam, and D. F. Blair. 1999. Function of proline residues of MotA in torque generation by the flagellar motor of *Escherichia coli*. *J. Bacteriol.* 181:3542–3551.
- Duncan, M. J., G. Li, J. S. Shin, J. L. Carson, and S. N. Abraham. 2004. Bacterial penetration of bladder epithelium through lipid rafts. *J. Biol. Chem.* 279:18944–18951.
- Gibson, R. L., J. L. Burns, and B. W. Ramsey. 2003. Pathophysiology and management of pulmonary infections in cystic fibrosis. *Am. J. Respir. Crit. Care Med.* 168:918–951.



10. Hayashi, F., K. D. Smith, A. Ozinsky, T. R. Hawn, E. C. Yi, D. R. Goodlett, J. K. Eng., S. Akira, D. M. Underhill, and A. Aderem. 2001. The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5. *Nature* **410**:1099–1103.
11. Inaba, K., M. Inaba, M. Deguchi, K. Hagi, R. Yasumizu, S. Ikehara, S. Muramatsu, and R. M. Steinman. 1993. Granulocytes, macrophages, and dendritic cells arise from a common major histocompatibility complex class II-negative progenitor in mouse bone marrow. *Proc. Natl. Acad. Sci. U. S. A.* **90**:3038–3042.
12. Luzar, M. A., M. J. Thomassen, and T. C. Montie. 1985. Flagella and motility alterations in *Pseudomonas aeruginosa* strains from patients with cystic fibrosis: relationship to patient clinical condition. *Infect. Immun.* **50**:577–582.
13. Lyczak, J. B., C. L. Cannon, and G. B. Pier. 2000. Establishment of *Pseudomonas aeruginosa* infection: lessons from a versatile opportunist. *Microbes Infect.* **2**:1051–1060.
14. Lyczak, J. B., C. L. Cannon, and G. B. Pier. 2002. Lung infections associated with cystic fibrosis. *Clin. Microbiol. Rev.* **15**:194–222.
15. MacFerrin, K. D., M. P. Terranova, S. L. Schreiber, and G. L. Verdine. 1990. Overproduction and dissection of proteins by the expression-cassette polymerase chain reaction. *Proc. Natl. Acad. Sci. U. S. A.* **87**:1937–1941.
16. Mahajan-Miklos, S., M. W. Tan, L. G. Rahme, and F. M. Ausubel. 1999. Molecular mechanisms of bacterial virulence elucidated using a *Pseudomonas aeruginosa*-*Caenorhabditis elegans* pathogenesis model. *Cell* **96**:47–56.
17. Mahenthalingam, E., M. E. Campbell, and D. P. Speert. 1994. Nonmotility and phagocytic resistance of *Pseudomonas aeruginosa* isolates from chronically colonized patients with cystic fibrosis. *Infect. Immun.* **62**:596–605.
18. Mahenthalingam, E., and D. P. Speert. 1995. Nonopsonic phagocytosis of *Pseudomonas aeruginosa* by macrophages and polymorphonuclear leukocytes requires the presence of the bacterial flagellum. *Infect. Immun.* **63**:4519–4523.
19. Montie, T. C., R. C. Craven, and I. A. Holder. 1982. Flagellar preparations from *Pseudomonas aeruginosa*: isolation and characterization. *Infect. Immun.* **35**:281–288.
20. Rahme, L. G., E. J. Stevens, S. F. Wolfort, J. Shao, R. G. Tompkins, and F. M. Ausubel. 1995. Common virulence factors for bacterial pathogenicity in plants and animals. *Science* **268**:1899–1902.
21. Sadikot, R. T., T. S. Blackwell, J. W. Christman, and A. S. Prince. 2005. Pathogen-host interactions in *Pseudomonas aeruginosa* pneumonia. *Am. J. Respir. Crit. Care Med.* **171**:1209–1223.
22. Sauer, K., and A. K. Camper. 2001. Characterization of phenotypic changes in *Pseudomonas putida* in response to surface-associated growth. *J. Bacteriol.* **183**:6579–6589.
23. Semmler, A. B., C. B. Whitchurch, and J. S. Mattick. 1999. A re-examination of twitching motility in *Pseudomonas aeruginosa*. *Microbiology* **145**:2863–2873.
24. Simpson, D. A., and D. P. Speert. 2000. RpmA is required for nonopsonic phagocytosis of *Pseudomonas aeruginosa*. *Infect. Immun.* **68**:2493–2502.
25. Smith, K. D., and A. Ozinsky. 2002. Toll-like receptor-5 and the innate immune response to bacterial flagellin. *Curr. Top. Microbiol. Immunol.* **270**:93–108.
26. Toutain, C. M., N. C. Caizza, M. E. Zegans, and G. A. O'Toole. 2007. Roles for flagellar stators in biofilm formation by *Pseudomonas aeruginosa*. *Res. Microbiol.* **158**:471–477.
27. Toutain, C. M., M. E. Zegans, and G. A. O'Toole. 2005. Evidence for two flagellar stators and their role in the motility of *Pseudomonas aeruginosa*. *J. Bacteriol.* **187**:771–777.

---

Editor: A. Camilli