

Adherence of *Streptococcus mutans* to Dextran Synthesized in the Presence of Extracellular Dextranucrase

HOWARD K. KURAMITSU

Department of Microbiology, Northwestern University Medical-Dental School, Chicago, Illinois 60611

Received for publication 15 November 1973

Live or heat-killed cells of *Streptococcus mutans* specifically adhere to dextran previously synthesized on glass surfaces by the action of extracellular dextranucrase.

Certain strains of *Streptococcus mutans* capable of synthesizing insoluble dextran appear to initiate the formation of dental plaque on smooth surfaces (8). Dextranucrase (EC 2.4.1.5.) from these strains catalyzes the formation of insoluble dextran from sucrose (4). This enzyme is primarily detected as an extracellular protein but has also been demonstrated in a cell-bound form (4). Recently, Mukasa and Slade (6) directly demonstrated that cell-bound dextranucrase can mediate the attachment of *S. mutans* to smooth surfaces. Previously, Gibbons and Fitzgerald (3) demonstrated that *S. mutans* could adhere to teeth coated with dextran synthesized by *Leuconostoc* ATCC 14935. This suggested that dextran synthesized on tooth surfaces by the action of the extracellular dextranucrase from *S. mutans* might also play a role in cellular adherence. This report demonstrates that *S. mutans* can specifically adhere to dextran formed on smooth surfaces by the action of the extracellular enzyme of the organism.

Human cariogenic *S. mutans* GS-5 and *S. salivarius* GS-15 were supplied by R. J. Gibbons, Harvard University Dental School. *S. mutans* strains HS-6, OMZ-176, FA-1 and *S. sanguis* 10556 were kindly supplied by H. D. Slade. All organisms were maintained and grown as previously described (5) except that *Bacillus stearothermophilus* was grown at 55 C. Dextran-coated glass surfaces were prepared by incubating 0.012 units of partially purified dextranucrase (0.12 units/mg) with 2% sucrose and saline-0.04% sodium azide (total volume 2.0 ml) in glass tubes (13 by 100 mm) inclined at a 30° angle. The enzyme was prepared after precipitation of the culture medium of *S. mutans* GS-5 with ammonium sulfate and passage of the enzyme through a Bio Gel A-15 column (2). After incubation for 18 h at 37 C, the tubes were decanted and gently washed

three times with saline. Visual examination of the tubes revealed a thin film of dextran as noted previously (6). Glucose-grown cells (approximately 3×10^8 per tube), washed three times with saline and suspended in saline-sodium azide, were added to the dextran-coated tubes in a total volume of 2.0 ml. The tubes were again incubated for 18 h at 37 C at an inclined angle. Adhered cells were gently washed three times with saline and suspended vigorously in 3.0 ml of 0.5 N NaOH, and the turbidity was determined at 540 nm (7).

When washed cell suspensions of *S. mutans* GS-5 were incubated with glass surface-coated dextran synthesized in the presence of the extracellular enzyme, significant cellular adherence was observed (Fig. 1). Cellular adherence was shown to be dependent on the prior incubation of the extracellular dextranucrase together with sucrose (Table 1). The omission of either component resulted in turbidity measurements




FIG. 1. Adherence of *S. mutans* GS-5 to dextran-coated glass surfaces as a function of cell concentration. Adherence assays were carried out as described in the text. Absorbance values have been corrected for blank tubes lacking sucrose. The cell suspension used contained approximately 2×10^9 cells per ml.

TABLE 1. Requirements for cellular adherence^a

Dextranucrase	Sucrose	Cells	A ₅₄₀
+	+	+	0.268
-	+	+	0.048
Boiled ^b	+	+	0.056
+	-	+	0.055
+	+	-	0.059
+	+	Boiled ^b	0.320

^a Adherence assays were carried out as described in the text.

^b Heated for 10 min at 100 C.

TABLE 2. Specificity of cellular adherence^a

Organism	Group ^b	A ₅₄₀ ^c
<i>S. mutans</i> GS-5	c	0.126
<i>S. mutans</i> HS-6	a	0.131
<i>S. mutans</i> FA-1	b	0.030
<i>S. mutans</i> OMZ-176	d	0.055
<i>S. salivarius</i> GS-15		0.013
<i>S. sanguis</i> 10556		0
<i>B. stearothermophilus</i> 1503-4R		0.001

^a Adherence assays were carried out as described in the text by using equal numbers of cells (quantitated by turbidity measurements in a Klett colorimeter).

^b Bratthall (1) classification of *S. mutans*.

^c Absorbance corrected for blank tubes lacking sucrose.

approximating that of NaOH solutions. Furthermore, heat-killed cells could absorb equally well to the dextran-coated tubes. This indicated that dextranucrase activity associated with the cells did not play a polymerizing role in the observed adherence.

Cellular adherence did not appear to be the result of a nonspecific trapping effect since cells of *S. salivarius*, *S. sanguis*, and *B. stearothermophilus* did not adhere significantly to the dextran-coated surfaces (Table 2). In contrast, three other strains of *S. mutans*, HS-6, FA-1, and OMZ-176, demonstrated varying degrees of adherence to the surfaces. These later

differences might be the result of variations in the cell-surface recognition sites of the organisms.

These results suggest that the cell surface of *S. mutans* contains sites which specifically interact with insoluble dextran molecules. The nature of these receptor sites is of interest and currently under investigation in several laboratories. Extrapolating from these results, it seems likely that *S. mutans* can adhere specifically to tooth surfaces coated with dextran produced by the action of its extracellular dextranucrase on dietary sucrose. Furthermore, these results indicate that dextran polymer formation need not be restricted to the cell surface of *S. mutans* for cellular adherence to occur as has been suggested previously (6).

This investigation was supported by Public Health Service grant DE-03258 from the National Institute of Dental Research.

LITERATURE CITED

1. Bratthall, D. 1970. Demonstration of five serological groups of Streptococcal strains resembling *Streptococcus mutans*. *Odontol. Revy* **21**:143-152.
2. Gibbons, R. J. 1972. Presence of an invertase-like enzyme and a sucrose permeation system in strains of *Streptococcus mutans*. *Caries Res.* **6**:122-131.
3. Gibbons, R. J., and R. J. Fitzgerald. 1969. Dextran-induced agglutination of *Streptococcus mutans* and its potential role in the formation of microbial dental plaque. *J. Bacteriol.* **98**:341-346.
4. Gibbons, R. J., and M. Nygaard. 1968. Synthesis of insoluble dextran and its significance in the formation of gelatinous deposits by plaque-forming streptococci. *Arch. Oral Biol.* **13**:1249-1262.
5. Kuramitsu, H. K. 1973. Characterization of invertase activity from cariogenic *Streptococcus mutans*. *J. Bacteriol.* **115**:1003-1010.
6. Mukasa, H., and H. D. Slade. 1973. Mechanism of adherence of *Streptococcus mutans* to smooth surfaces. *Infect. Immunity* **8**:555-562.
7. Olson, G. A., A. S. Bleiweis, and P. A. Small, Jr. 1972. Adherence inhibition of *Streptococcus mutans*: an assay reflecting a possible role of antibody in dental caries prophylaxis. *Infect. Immunity* **5**:419-427.
8. Scherp, H. W. 1971. Dental caries, prospects for prevention. *Science* **173**:1199-1205.